ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ ORANGE ZEST- aspirin, chlorpheniramine maleate, phenylephrine bitartrate tablet, effervescent Bayer HealthCare LLC.

AS Severe Cold PowerFast Fizz Orange Zest UI1614457

Drug Facts

Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)*

Chlorpheniramine maleate 2 mg

Phenylephrine bitartrate 7.8 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Antihistamine

Nasal decongestant

Uses

Uses

 \cdot temporarily relieves these symptoms due to a cold:

- \cdot minor aches and pains \cdot headache \cdot runny nose
- \cdot nasal and sinus congestion \cdot sneezing \cdot sore throat
- \cdot temporarily reduces fever

Directions

Directions

• adults and children 12 years and over: take 2 tablets fully dissolved

in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours

or as directed by a doctor.

• children under 12 years: do not use

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

 \bullet hives \bullet facial swelling \bullet asthma (wheezing) \bullet shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

• if you are allergic to aspirin or any other pain reliever/fever reducer

if you are now taking a prescription monoamine oxidase inhibitor
(MAOI) (certain drugs for depression, psychiatric, or emotional

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

 if you have ever had an allergic reaction to this product or any of its ingredients

• in children under 12 years of age

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis,

or kidney disease

- you are taking a diuretic
- you have
- asthma diabetes
- thyroid disease glaucoma

- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- \cdot taking a prescription drug for
- · gout · diabetes · arthritis
- · taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint vomit blood have bloody or black stools
- have stomach pain that does not get better
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness, or sleeplessness occur

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Other information

Other information

- each tablet contains: potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

Inactive ingredients

Inactive ingredients anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #6, flavor, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

Questions or Comments

Questions or comments? 1-800-986-0369 (Mon - Fri 9AM - 5PM EST)

Carton 24 count



Alka-Seltzer PLUS® Severe Cold Orange Zest

PowerFast Fizz™

NEW Improved Flavor

ASPIRIN (NSAID) / Pain Reliever-Fever Reducer

Chlorheniramine Maleate / Antihistamine

Phenylephrine Bitartrate / Nasal Decongestant

- Nasal Congestion
- Runny Nose

- Sore Throat
- Headache + Body AcheSinus Pressure

24 EFFERVESCENT TABLETS

ALKA-SELTZEI ZEST aspirin, chlorphenira Product Informa Product Type Route of Administr	amine male		e bitartra			-	ΞE	
Product Informa Product Type Route of Administr	ation			ate tablet, e	effervescent			
Product Type Route of Administr		HUMAN OTC DRUG						
Product Type Route of Administr		HUMAN OTC DRUG						
Route of Administr	ation	HUMAN OTC DRUG						
	ation		Iten	ltem Code (Source)		NDC:0280-0020		
		ORAL						
Active Ingredien	t/Active	Moiety						
	Ingre	edient Name			Basis of S	Strength	Strength	
PHENYLEPHRINE BITARTRATE (UNII: 2703Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)PHENYLEPHRIN BITARTRATE						NE	7 mg	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U) CHLORPHENIRAMINE - CHLORPH						AMINE	2 mg	
ASPIRIN (UNII: R16C05Y76E) (ASPIRIN - UNII:R16C05Y76E) ASPIRIN							325 mg	
Inactive Ingredie	ents							
		Ingredient Name	•			Stre	Strength	
FD&C YELLOW NO. 6	(UNII: H77V	EI93A8)						
SUCRALOSE (UNII: 96K6UQ3ZD4)								
MANNITOL (UNII: 30WL								
POTASSIUM BICARBO								
SODIUM BICARBONAT		DF5V39QO)						
POVIDONE (UNII: FZ98	•							
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) FD&C RED NO. 40 (UNII: WZB9127XOA)								
CALCIUM SILICATE (UNII: S4255P4G5M)								
DIMETHICONE (UNII: 92RU3N3Y10)								
	21105115110	7						
Product Charact	eristics							
Color	white (SPO	TTED)	Score		no	score		
Shape	ROUND		Size		25	25mm		
Flavor	ORANGE		Imprint Code		AS	ASP;ORANGE		
Contains								
Packaging								
# Item Code	Pac	kage Description			ng Start Ite		ing End ite	

10 in 1 CARTON	07/01/2020	
2 in 1 POUCH; Type 0: Not a Combination Product		
12 in 1 CARTON	03/30/2023	
2 in 1 POUCH; Type 0: Not a Combination Product		
nformation		
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	2 in 1 POUCH; Type 0: Not a Combination Product 12 in 1 CARTON 2 in 1 POUCH; Type 0: Not a Combination Product	2 in 1 POUCH; Type 0: Not a Combination 12 in 1 CARTON 03/30/2023 2 in 1 POUCH; Type 0: Not a Combination Product

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 1/2025

Bayer HealthCare LLC.