

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**WinCo Foods, LLC**

-----  
**Drug Facts**

**Active ingredient (in each capsule)**

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- pain gets worse or last more than 10 days
- fever gets worse or last more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

### **Inactive ingredients**

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

### **Questions?**

Or to report an adverse event call **800-824-1706** Monday-Friday 9am-4pm MST

### **Principal Display Panel**

†Compare to the active ingredient in Advil® Liqui-Gels®

### **Ibuprofen**

Ibuprofen Capsules 200 mg (NSAID)/

Pain Reliever-Fever Reducer

SOFTGELS\*\*

\*\*Liquid-Filled Capsules

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.**

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

PRODUCT OF USA

### **Package Label**

PLD-AD7 F0020457 WFLH11  
 Rev# 03/12  
 Lot No.:  
 Exp. Date:



**Drug Facts (continued)**  
**Directions**  
 ■ do not take more than directed  
 ■ the smallest effective dose should be used  
 ■ adults and children 12 years and over:  
 ■ take 1 capsule every 4 to 6 hours while symptoms persist  
 ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used  
 ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor  
 ■ children under 12 years: ask a doctor  
**Other information**  
 ■ each capsule contains: **potassium 20 mg**  
 ■ read all warnings and directions before use.  
 ■ Keep carton.  
 ■ store at 20° to 25°C (68° to 77°F)  
 ■ avoid excessive heat above 40°C (104°F). Protect from light.  
**Inactive ingredients** FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol  
**Questions?** Or to report an adverse event call 800-924-1706 Monday – Friday 9am – 4pm MST

**DISTRIBUTED BY:**  
 WINCO FOODS, LLC, BOISE, ID 83704  
**PRODUCT OF U.S.A.**

†This product is not manufactured or distributed by PFIZER CONSUMER HEALTHCARE, distributor of ADVIL® Liqui-Gels®.

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING**

**Drug Facts (continued)**  
**Ask a doctor before use if**  
 ■ the stomach bleeding warning applies to you  
 ■ you have problems or serious side effects from taking pain relievers or fever reducers  
 ■ you have a history of stomach problems, such as heartburn  
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma  
 ■ you are taking a diuretic  
**Ask a doctor or pharmacist before use if you are**  
 ■ under a doctor's care for any serious condition  
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
 ■ taking any other drug  
**When using this product**  
 ■ take with food or milk, if stomach upset occurs  
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed  
**Stop use and ask a doctor if**  
 ■ you experience any of the following signs of stomach bleeding:  
 ■ have bloody or black stools ■ feel faint  
 ■ have stomach pain that does not get better  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ redness or swelling is present in the painful area  
 ■ any new symptoms appear  
**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts**  
**Active ingredient (in each capsule) Purpose**  
 Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*.....Pain reliever/fever reducer  
 †(present as the free acid and potassium salt)  
 \*nonsteroidal anti-inflammatory drug  
**Uses**  
 ■ temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches  
 ■ minor pain of arthritis ■ toothache  
 ■ backache ■ the common cold  
 ■ menstrual cramps  
 ■ temporarily reduces fever  
**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
 ■ hives ■ facial swelling ■ asthma (wheezing)  
 ■ shock ■ skin reddening ■ rash ■ blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 ■ are age 60 or older  
 ■ have had stomach ulcers or bleeding problems  
 ■ take a blood thinning (anticoagulant) or steroid drug  
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ take more or for a longer time than directed  
**Do not use**  
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
 ■ right before or after heart surgery

†Compare to the active ingredient in ADVIL® Liqui-Gels®

Relieves: Minor Aches & Pains, Fever



**80**  
Softgels\*\*

**WinCo FOODS**  
**Ibuprofen**  
 IBUPROFEN CAPSULES  
 200 mg (NSAID)/  
 PAIN RELIEVER - FEVER REDUCER

\*\*Liquid Filled Capsules



**WinCo Ibuprofen Softgels**

**IBUPROFEN**

ibuprofen capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-141
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

**Inactive Ingredients**

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDRO XIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

**Product Characteristics**

<b>Color</b>	BLUE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	IB200
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-141-80	1 in 1 BOX	12/31/2014	
1		80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	12/31/2014	

**Labeler** - WinCo Foods, LLC (056098817)

Revised: 12/2018

WinCo Foods, LLC