

EFFERVESCENT COLD RELIEF- aspirin, chlorpheniramine maleate, phylephrine bitartrate tablet, effervescent
Topco Associates LLC

TopCare Effervescent Cold Relief

Active ingredients

Active ingredients

(in each effervescent tablet) Purpose

Aspirin 325 mg (NSAID)*. Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg Antihistamine

Phenylephrine Bitartrate 7.8 mg Nasal decongestant

*Nonsteroidal anti-inflammatory drug

Purpose

(in each effervescent tablet) Purpose

Aspirin 325 mg (NSAID)*. Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg Antihistamine

Phenylephrine Bitartrate 7.8 mg Nasal decongestant

*Nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves these symptoms due to the common cold
- sneezing ■ nasal congestion ■ sore throat
- headache ■ minor aches and pains ■ runny nose
- sinus congestion and pressure
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do Not Use

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer

■ if you are allergic to aspirin ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Ask a doctor before use if you have

Ask a doctor before use if you have ■ asthma ■ glaucoma ■ diabetes ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ been placed on a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are ■ you are taking sedatives or tranquilizers ■ presently taking a prescription drug

■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

When using this product ■ **do not exceed recommended dosage**

■ you may get drowsy

■ avoid alcoholic drinks

- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

Keep out of reach of children.

In case of overdose

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours
- do not take more than 8 tablets in 24 hours
- *children under 12 years: ask a doctor*

Other information

each tablet contains: sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

Inactive ingredients

acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Questions or Comments?

Call 1-888-423-0139

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



Drug Facts (continued)

for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer if you are allergic to aspirin or if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients, ask a doctor before use. If stomach bleeding warning applies to you, you have a history of stomach problems such as heartburn or you have high blood pressure, heart disease, liver cirrhosis, or kidney disease or you are taking a diuretic.

Ask a doctor before use if you have asthma, glaucoma, diabetes, thyroid disease, trouble breathing due to an enlarged prostate gland, or a breathing problem such as emphysema or chronic bronchitis. If you are taking a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers, or presently taking a prescription drug or taking sedatives or tranquilizers for anticoagulation (thinning the blood), diabetes, gout or arthritis.

When using this product do not exceed recommended dosage.

You may get drowsy if you get drowsy or avoid alcoholic drinks.

Exhaustion may occur, especially in children if you get drowsy or avoid alcoholic drinks.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The stomach bleeding problems may increase drowsiness if you get drowsy or avoid alcoholic drinks.

Be careful when driving a motor vehicle or operating machinery if you get drowsy or avoid alcoholic drinks.

STOP use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel faint
- have bloody or black stools
- have stomach pain that does not get better

Drug Facts

Active Ingredients
(in each effervescent tablet)

Aspirin 325 mg (NSAID)
Chlorpheniramine maleate 2 mg
Phenylephrine bitartrate 7.5 mg
Nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/fever reducer
Sore throat

Uses
temporarily relieves these symptoms due to the common cold:
sore throat
nasal congestion
runny nose
headache
fever
temporarily reduces fever

Warnings
Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor before these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives or facial swelling.

TopCare health

Effervescent Cold Relief

ASPIRIN (NSAID)/PAIN RELIEVER-FEVER REDUCER
CHLORPHENIRAMINE MALEATE/ANTIHISTAMINE
PHENYLEPHRINE BITARTRATE/NASAL DECONGESTANT

20 EFFERVESENT TABLETS ORIGINAL FLAVOR

TopCare health

NDC 36800-768-20

COMPARE TO ALKA-SELTZER PLUS® COLD ACTIVE INGREDIENTS*

Effervescent Cold Relief

ASPIRIN (NSAID)/PAIN RELIEVER-FEVER REDUCER
CHLORPHENIRAMINE MALEATE/ANTIHISTAMINE
PHENYLEPHRINE BITARTRATE/NASAL DECONGESTANT

FAST RELIEF OF:

- Nasal Congestion
- Runny Nose
- Headache & Body Aches
- Sore Throat
- Sinus Pressure

20 EFFERVESENT TABLETS ORIGINAL FLAVOR

Drug Facts (continued)

- an allergic reaction occurs. Seek medical help right away
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
(see overdose warning)

- do not use more than directed
- adults and children 12 years and over: take 2 tablets completely dissolved in 4 oz of water every 4 hours
- do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

Other information
each tablet contains: sodium 464 mg

- phenylketonurics: contains phenylalanine 9 mg per tablet
- store at room temperature (59°-86°F)

Inactive ingredients
acesulfame potassium, aspartame, citric acid, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

Questions or comments?
Call 1-888-423-0139

DISTRIBUTED BY TOPCOO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
© TOPCOO TOWARZ21 QUESTIONS? 1-888-423-0139
topcare@topcoo.com www.topcarebrand.com

Visit here for more information:
<http://topbrnds.com/48zqf>

This Top Care® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.



PN: 87360 TC C3
REV: 05/21

*This product is not manufactured, distributed or licensed by Bayer, the maker of Alka-Seltzer Plus® Cold.

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

EFFERVESENT COLD RELIEF

aspirin, chlorpheniramine maleate, phylephrine bitartrate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-768
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ASPARTAME (UNII: Z0H242BBR1)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
DOCUSATE SODIUM (UNII: F05Q2T2JA0)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-768-20	10 in 1 CARTON	07/01/2010	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2010	

Labeler - Topco Associates LLC (006935977)

Revised: 11/2024

Topco Associates LLC