

GOOD SENSE TUSSIN CF- dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution
L. Perrigo Company

Perrigo Tussin CF Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Phenylephrine HCl, USP 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic

bronchitis or emphysema

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).
Do not refrigerate.

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

Principal Display Panel

Non-Drowsy

Tussin CF

Cough Suppressant (Dextromethorphan HBr)

Expectorant (Guaifenesin)

Nasal Decongestant (Phenylephrine HCl)

Multi-Symptom Cold

Relieves:

Nasal Congestion

Cough

Mucus

Peak Cold

Adult

For Ages 12 & Over

Compare to active ingredients of Robitussin® Multi-Symptom Cold

4 FL OZ (118 mL)

NDC 0113-0703-26

GOODSENSE®

Non-Drowsy

Tussin

CF

Cough Suppressant
(Dextromethorphan HBr)
Expectorant (Guaifenesin)
Nasal Decongestant
(Phenylephrine HCl)

Multi-Symptom Cold

Relieves:

- Nasal Congestion
- Cough • Mucus



Peak Cold

Adult

For Ages 12 & Over

Compare to active ingredients of
Robitussin® Multi-Symptom
Cold

100%
SATISFACTION
GUARANTEED

4 FL OZ (118 mL)

NDC 0113-0703-26

GOODSENSE®

Non-Drowsy

Tussin

CF

Cough Suppressant
(Dextromethorphan HBr)
Expectorant (Guaifenesin)
Nasal Decongestant
(Phenylephrine HCl)

Multi-Symptom Cold

Relieves:

- Nasal Congestion
- Cough • Mucus



Peak Cold


Adult

For Ages 12 & Over

Compare to active ingredients of
Robitussin® Multi-Symptom
Cold

100%
SATISFACTION
GUARANTEED

4 FL OZ (118 mL)



CF Multi-Symptom Tussin

GOODSENSE®

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr,
USP 20 mg.....Cough suppressant
Guaifenesin, USP 200 mg.....Expectorant
Phenylephrine HCl,
USP 10 mg.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition. ►

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

- each 10 mL contains: sodium 5 mg
- store at 20-25°C (68-77°F).
- Do not refrigerate.

Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

1-800-719-9260

Gluten Free

Dosage Cup Included

**DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING**

Distributed By **Perrigo®** Allegan, MI 49010

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



LOT NO.

EXP.

: 51626 C2 C7

GOOD SENSE TUSSIN CF

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-0703
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics				
Color		RED (clear)	Score	
Shape			Size	
Flavor		FRUIT	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0703-26	1 in 1 CARTON	07/29/2014	08/31/2025
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0113-0703-34	1 in 1 CARTON	07/29/2014	06/30/2024
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	07/29/2014	08/31/2025

Labeler - L. Perrigo Company (006013346)

Revised: 12/2024

L. Perrigo Company