

**DAYTIME SINUS NIGHTTIME SINUS MAXIMUM STRENGTH- acetaminophen,
diphenhydramine hcl, guaifenesin, phenylephrine hcl
Target Corporation**

Target 44-615694-SM

Active ingredients (in each caplet) (Sinus Day)

Acetaminophen 325 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever

Expectorant

Nasal decongestant

Active ingredients (in each caplet) (Sinus Night)

Acetaminophen 325 mg

Diphenhydramine HCl 12.5 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever

Antihistamine/cough suppressant

Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
 - cough (***Nighttime only***)
 - runny nose and sneezing (***Nighttime only***)
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive (***Daytime only***)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other product containing diphenhydramine, even one used on skin
(Nighttime only)

Ask a doctor before use if you have

- heart disease
- liver disease
- diabetes
- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis **(Nighttime only)**
- glaucoma **(Nighttime only)**

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers **(Nighttime only)**

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children **(Nighttime only)**
- marked drowsiness may occur **(Nighttime only)**
- alcohol, sedatives, and tranquilizers may increase drowsiness **(Nighttime only)**
- avoid alcoholic beverages **(Nighttime only)**

- be careful when driving a motor vehicle or operating machinery (**Nighttime only**)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If taking NIGHTTIME and DAYTIME products, carefully read each section to ensure correct dosing. Do not take DAY & NIGHT at the same time.

Directions

- **do not use more than directed**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients (Daytime only)

corn starch, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Inactive ingredients (Nighttime only)

corn starch, croscarmellose sodium, crospovidone, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, iron oxide yellow, magnesium stearate, methacrylic acid and

ethyl acrylate copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium bicarbonate, stearic acid, talc, titanium dioxide

Questions?

Call 1-800-910-6874

Principal Display Panel

NDC 11673-694-09

**Compare to active ingredients in
Maximum Strength Mucinex®
SINUS-MAX® Day***

maximum strength

daytime sinus

acetaminophen

guaifenesin, phenylephrine HCl

pain reliever, expectorant,

nasal decongestant

ACTUAL

SIZE

AGES

12+

YEARS

12

CAPLETS

up&up™

Compare to active ingredients in

Maximum Strength Mucinex®

SINUS-MAX® Night*

maximum strength

nighttime sinus

acetaminophen

diphenhydramine HCl,

phenylephrine HCl

pain reliever,

antihistamine/cough suppressant,

nasal decongestant

ACTUAL

SIZE

AGES

12+

YEARS

8

CAPLETS

up&up™

12 DAY AND 8 NIGHT

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of the registered trademark Maximum Strength Mucinex® SINUS-MAX® Day & Night.

50844 REV0517B61569409

Distributed by Target Corp., Minneapolis, MN 55403

TM & ©2021 Target Brands, Inc.

Questions? Call 1-800-910-6874



<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> with any other product containing diphenhydramine, even one used on skin (<i>Nighttime only</i>) <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> heart disease thyroid disease difficulty in urination due to enlargement of the prostate gland asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus) a breathing problem such as emphysema or chronic bronchitis (<i>Nighttime only</i>) glaucoma (<i>Nighttime only</i>) <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin taking sedatives or tranquilizers (<i>Nighttime only</i>) <p>When using this product</p> <ul style="list-style-type: none"> do not exceed recommended dosage excitability may occur, especially in children (<i>Nighttime only</i>) marked drowsiness may occur (<i>Nighttime only</i>) alcohol, sedatives, and tranquilizers may increase drowsiness (<i>Nighttime only</i>) avoid alcoholic beverages (<i>Nighttime only</i>) 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> be careful when driving a motor vehicle or operating machinery (<i>Nighttime only</i>) <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> nervousness, dizziness, or sleeplessness occur pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>If taking NIGHTTIME and DAYTIME products, carefully read each section to ensure correct dosing. Do not take DAY & NIGHT at the same time.</p>
---	--

Target 44-615694

DAYTIME SINUS NIGHTTIME SINUS MAXIMUM STRENGTH

acetaminophen, diphenhydramine hcl, guaifenesin, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-694
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-694-09	1 in 1 CARTON; Type 0: Not a Combination Product	07/01/2017	03/27/2026

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	8

Part 1 of 2

DAYTIME SINUS MAXIMUM STRENGTH

acetaminophen, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;615
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2013	

Part 2 of 2

NIGHTTIME SINUS MAXIMUM STRENGTH

acetaminophen, diphenhydramine hcl, phenylephrine hcl tablet, film coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;694
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2017	03/27/2026

Labeler - Target Corporation (006961700)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(11673-694)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11673-694) , pack(11673-694)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(11673-694)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(11673-694)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11673-694)

Revised: 7/2024

Target Corporation