ZHONG HUA FENG SHI DIE DA GAO EXTERNAL ANALGESIC PLASTER- menthol, methyl salicylate patch

Albert Max, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ZHONG HUA FENG SHI DIE DA GAO External Analgesic Plaster

Drug Facts

Active Ingredients

Menthol 4% Methyl Salicylate 1%

Purpose

External Analgesic

Uses

For the temporary relief of minor aches and pains of muscles and joints due to: •arthritis •strains •bruises •sprains •simple backache

Warnings

For external use only

Do not use

- •on wounds •irritated or damaged skin •sensitive skin
- •with a heating pad •if pregnant
- •with, or as the same time as, other external analgesic products
- •if allergic to aspirin or salicylates
- •if allergic to any ingredients of this product

When using this product

- •avoid contact with the eyes
- do not bandage tightly
- •do not apply to wounds or damaged skin
- •do not use more than 3 plasters a day
- •do not use 1 hour before or after bathing

Stop use and ask a doctor if

- condition worsens
- •symptoms persist for more than 7 days
- •symptoms clear up and occur again within a few days
- •excessive irritation of the skin develops
- •rashes, itching or redness are present
- •side effects occur.

If pregnant, ask a health professional before use.

Keep out of reach of children to avoid accidental poisoning.

In case of overdose, get medical help or contact a Poison Control Center right away. Do not use otherwise than as directed.

Directions

- •Adults and children 12 years of age and older: clean and dry affected area. Remove plaster from the protective layer, apply to the affected area, rub the area with your hand to secure the patch.
- •Plaster will remain effective for 8 hours.
- •Avoid using the same patch on the same area for more than 2 days.
- •Do not apply to area with excessive hair. Adhesive plaster may hurt skin upon removal.
- •Wet patch with warm water before removing from skin
- •Children under 12 years of age: Do not use or consult a doctor.

Other Information

- •Store unused patch in the pouch
- •Store at room temperature, 20 to 25° C(68 to 77° F)

Inactive ingredients

Chinese Clematidis Root & Rhizome, Yellow wax, Momordica cochinchinensis seeds, Castor Seeds, Myrrh, Frankincense, Sesame Oil, Resin, FD&C Blue No. 2, FD&C Yellow No. 4, Zinc Oxide


Questions or Comments Call: 1-800-796-1888

Distributed by:

A&A Medical Supplies, Inc. South El Monte, CA 91733 Made in Taiwan

Cooling Plaster FOR EXTERNAL USE ONLY Contains 5 (7.5×10cm 2.9 inch×3.9inch) Patches

Packaging



Methyl Salicylate 1%. External Analogues For the temporary relief of minor aches and pains of muscles and joints due to: arthritis * strains * bruises * sprains * simple backache Warnings For external use only Do not use on wounds * irritated or damaged skin * sensitive skin with a heating pad * if pregnant with or as the same time as, other external analgesic products with or as the same time as, other external analgesic products if allergic to any ingredients of this product When using this product avoid contact with the eyes avoid contact with the eyes avoid contact with the eyes do not apply to wounds or damaged skin do not apply to wounds or damaged skin do not use more than 3 plasters a day do not use 1 hour before or after bathing Stop use and ask a doctor if condition worsens symptoms persist for more than 7 days symptoms clear up and occur again within a few days excessive irritation of the skin develops symptoms clear up and occur again within a few days excessive irritation of the skin develops symptoms clear up and occur again within a few days excessive irritation of the skin develops successive irritations of the skin develops symptoms or the skin develops successive irritations of the skin develops successive irritations of the skin develops successive irritations are present side effects occur. if pregnant, ask a health professional before use. Keep out of reach of children to avoid accidental poisoning. In case of overdose, get medical help or contact a Poison Control Centers with your hand to secure the patch. Directions Advilts and children 12 years of age and older; clean and dry affected area. Remove plaster from the protective layer, apply to the affected area, rub the with your hand to secure the patch. Do not apply to area with excessive hair. Adhesive plaster may hurt skin up removal. Wet patch with warm water before removing from sk	Active Ingredients Menthol 4%	Purpose External Analge
and joints due to: *arthritis strains *bruises *sprains *simple backache *Warnings* For external use only Do not use *on wounds *irritated or damaged skin *sensitive skin *with a heating pad *if pregnant *with, or as the same time as, other external analgesic products *if allergic to aspirin or salicylates *on or bandage tightly when using this product *avoid contact with the eyes *do not bandage tightly or damaged skin *do not use more than 3 plasters a day *do not use a hour before or after bathing Stop use and ask a doctor if *condition worsens *symptoms persist for more than 7 days *symptoms clear up and occur again within a few days *excessive irritation of the skin develops *symptoms clear up and occur again within a few days *excessive irritation of the skin develops *side effects occur. *If pregnant, ask a health professional before use. Keep out of reach of children to avoid accidental poisoning. In case of overdose, get medical help or contact a Poison Control Centeraway. Do not use otherwise than as directed. *Prections *Adults and children 12 years of age and older: clean and dry affected area. *Remove plaster from the protective layer, apply to the affected area, rub the with your hand to secure the patch. *Plaster will remain effective for 8 hours. *Po not apply to alea with excessive hair. Adhesive plaster may hurt skin up removal. *Wet patch with warm water before removing from skin. *Children under 12 years of age: Do not use or consult a doctor. *Other Information *Store unused patch in the pouch.	Methyl Salicylate 1%	External Analge
For external use only Do not use on wounds * irritated or damaged skin * sensitive skin with a heating pad * if pregnant with, or as the same time as, other external analgesic products if allergic to aspirin or salicylates if allergic to aspirin or salicylates if allergic to aspirin or salicylates with product avoid contact with the eyes do not bandage tightly of on the same of the same o	and joints due to:	
on wounds * irritated or damaged skin * sensitive skin * with a heating pad * if pregnant * with, or as the same time as, other external analgesic products * if allergic to aspirin or salicylates * if allergic to aspiring or damaged skin * if asp	<i>Warnings</i> For external use only	
avoid confact with the eyes do not bandage tightly do not apply to wounds or damaged skin do not bandage tightly do not apply to wounds or damaged skin do not use more than 3 plasters a day do not use 1 hour before or after bathing Stop use and ask a doctor if condition worsens sometimes before more than 7 days symptoms clear up and occur again within a few days excessive irritation of the skin develops rashes; fiching or redness are present side effects occur. If pregnant, ask a health professional before use. If pregnant is ask of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the professional before use. If pregnant is a stop of the pr	 on wounds * irritated or damaged skin * se with a heating pad * if pregnant with, or as the same time as, other external if allergic to aspirin or salicylates 	
condition worsens symptoms persist for more than 7 days symptoms persist for more than 8 days rashes, tiching or redness are present side effects occur. If pregnant, ask a health professional before use. Keep out of reach of children to avoid accidental poisoning. In case of overdose, get medical help or contact a Poison Control Center Do not use otherwise than as directed. Directions Adults and children 12 years of age and older: clean and dry affected area. When the profestive layer, apply to the affected area, not the with your hand to secure the patch. Plaster will remain effective for 8 hours. Avoid using the same patch on the same are for more than 2 days. Well patch with warm water before removing from skin. Children under 12 years of age: Do not use or consult a doctor. Other Information Store unused patch in the pouch.	avoid contact with the eyes do not bandage tightly do not apply to wounds or damaged skin do not use more than 3 plasters a day	
In case of overdose, get medical help or contact a Poison Control Center away. Do not use otherwise than as directed. Pirections Adults and children 12 years of age and older: clean and dry affected area. Remove plaster from the protective layer, apply to the affected area, rub the with your hand to secure the patch. Plaster will remain effective for 8 hours. Plaster will remain effective for 8 hours. Do not apply to area with excessive hair. Adhesive plaster may hurt skin upremoval. Wet patch with warm water before removing from skin. Children under 12 years of age: Do not use or consult a doctor. Other Information Store unusued patch in the pouch	condition worsens symptoms persist for more than 7 days symptoms clear up and occur again within excessive irritation of the skin develops	a few days
Adults and children 12 years of age and older: clean and dry affected area. Remove plaster from the protective layer, apply to the affected area, rub the with your hand to secure the patch. Plaster will remain effective for 8 hours. Plaster will remain effective for 8 hours. Plaster will remain effective for 8 hours. On ond apply to area with excessive hair. Adhesive plaster may hurt skin upremoval. Wet patch with warm water before removing from skin. Children under 12 years of age: Do not use or consult a doctor. Other Information Store unused patch in the pouch.	In case of overdose, get medical help or c	e use. idental poisoning. ontact a Poison Control Center
Store unused patch in the pouch	Adults and children 12 years of age and old Remove plaster from the protective layer, a with your hand to secure the patch. Plaster will remain effective for 8 hours. Avoid using the same patch on the same a Do not apply to area with excessive hair. A removal.	oply to the affected area, rub the a ea for more than 2 days. dhesive plaster may hurt skin upo
	Store unused patch in the pouch) 77°F)



A&A Medical Supplies, Inc.
Distributed by: A&A Medical Supplies, Inc. South El Monte, CA 91733

Questions or Comments Call:1-800-796-1888

ZHONG HUA FENG SHI DIE DA GAO EXTERNAL ANALGESIC PLASTER

menthol, methyl salicylate patch

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68233-804
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10 EIP3A) (MENTHOL - UNII:L7T10 EIP3A)	MENTHOL, UNSPECIFIED FORM	0.04 g in 1 g	
METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ)	METHYL SALICYLATE	0.01g in 1g	

Inactive Ingredients		
Ingredient Name	Strength	
CLEMATIS CHINENSIS ROOT (UNII: 8Z18N528CU)		
YELLOW WAX (UNII: 2ZA36H0S2V)		
MOMORDICA COCHINCHINENSIS SEED (UNII: 2T87O1UPVD)		
RICINUS COMMUNIS SEED (UNII: 7EK4SFN1TX)		
MYRRH (UNII: JC71GJ1F3L)		
FRANKINCENSE (UNII: R9 XLF1R1WM)		
SESAME O IL (UNII: QX10 HYY4QV)		

PINUS MASSONIANA RESIN (UNII: 64S07U83T7)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
YELLOW OB (UNII: QE535CBH7S)	
ZINC OXIDE (UNII: SOI2LOH54Z)	

Product Characteristics			
Color	green	Score	
Shape	RECTANGLE	Size	
Flavor		Imprint Code	
Contains			

	Packaging			
7	# Item Code	Package Description	Marketing Start Date	Marketing End Date
:	NDC:68233-804-01	5 in 1 BAG	11/01/2005	
:	1	1.7 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	11/0 1/20 0 5	

Labeler - Albert Max, Inc. (149445798)

Revised: 12/2018 Albert Max, Inc.