

KEEPAN A- benzalkonium chloride, lidocaine hydrochloride spray
Sato Pharmaceutical Co., Ltd.

Keepan A

Active ingredients

Benzalkonium chloride 0.1%

Lidocaine hydrochloride 1%

Purpose

Benzalkonium chloride First aid antiseptic

Lidocaine hydrochloride External analgesic

Uses First aid for minor cuts, scrapes and burns:

- to help reduce the risk of infection
- for the temporary relief of pain and itching

Warnings

For external use only

Do not use

- in the eyes ■ in large quantities
- over large areas of the body ■ over raw surfaces or blistered areas
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep or puncture wounds ■ animal bites ■ serious burns

When using this product

- avoid contact with the eyes

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days, or clear up and occur again within a few days
- irritation, redness, swelling or pain persists or increases or a rash or infection develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 2 years of age and older:

- clean the affected area
- spray a small amount directly on the affected area or apply with clean gauze or cotton saturated with KEEPAN A 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged, let dry first

Children under 2 years of age: Do not use, consult a doctor.

Inactive ingredients alcohol, fragrance, hydrochloric acid, purified water, sodium

chloride, sodium citrate

PUSH

KEEPAN A

Drug Facts

| Active ingredients | Purpose |
|---------------------------------|----------------------|
| Benzalkonium chloride 0.1%..... | First aid antiseptic |
| Lidocaine hydrochloride 1%..... | External analgesic |

Uses First aid for minor cuts, scrapes and burns:
 to help reduce the risk of infection
 for the temporary relief of pain and itching

Warnings
For external use only

Do not use
 in the eyes in large quantities
 over large areas of the body over raw surfaces or blistered areas
 longer than 1 week unless directed by a doctor

Ask a doctor before use if you have
 deep or puncture wounds animal bites serious burns

When using this product
 avoid contact with the eyes

Stop use and ask a doctor if
 condition worsens
 symptoms persist for more than 7 days, or clear up and occur again within a few days
 irritation, redness, swelling or pain persists or increases or a rash or infection develops

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
 Adults and children 2 years of age and older:
 clean the affected area
 spray a small amount directly on the affected area or apply with clean gauze or cotton saturated with KEEPAN A 1 to 3 times daily
 may be covered with a sterile bandage
 if bandaged, let dry first
 Children under 2 years of age: Do not use, consult a doctor

Inactive ingredients alcohol, fragrance, hydrochloric acid, purified water, sodium chloride, sodium citrate

Manufactured by
SATO PHARMACEUTICAL CO.,LTD.
 1-5-27 MOTOAKASAKA, MINATO-KU, TOKYO, JAPAN
 Reports of serious side effects associated with use of this product can be sent to:
SATO PHARMACEUTICAL, INC
 20501 Earl Street, Suite 3, Torrance, CA 90503



KEEPAN A

**Benzalkonium Chloride 0.1%
Lidocaine Hydrochloride 1%**

**FIRST AID ANTISEPTIC
EXTERNAL ANALGESIC**
For minor cuts, scrapes and burns

キーパンA
救急殺菌剤、外用鎮痛剤
きり傷、すり傷に

2 FL OZ (60 mL)

KEEPAN A

FIRST AID ANTISEPTIC EXTERNAL ANALGESIC For minor cuts, scrapes and burns

キーパンA 救急殺菌剤、外用鎮痛剤 きり傷、すり傷に



3 49873 01902 4

キーパンA

EXP. DATE
使用期限

LOT
製造記号

US-B

KEEPAN A

benzalkonium chloride, lidocaine hydrochloride spray

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:49873-019 |
| Route of Administration | TOPICAL | | |

| Active Ingredient/Active Moiety | | | | |
|----------------------------------------------------------------------------|------------------------------------------|-------------------------------------------------------------|----------------------|--------------------|
| Ingredient Name | | Basis of Strength | Strength | |
| BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII: 7N6JUD5X6Y) | | BENZALKONIUM CHLORIDE | 0.1 g in 100 mL | |
| LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII: 98PI200987) | | LIDOCAINE HYDROCHLORIDE ANHYDROUS | 1 g in 100 mL | |
| Inactive Ingredients | | | | |
| Ingredient Name | | | Strength | |
| ALCOHOL (UNII: 3K9958V90M) | | | | |
| HYDROCHLORIC ACID (UNII: QTT17582CB) | | | | |
| WATER (UNII: 059QF0KO0R) | | | | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | | | | |
| TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K) | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:49873-019-01 | 1 in 1 CARTON | 07/15/1981 | |
| 1 | | 60 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| OTC Monograph Drug | M003 | 07/15/1981 | | |

Labeler - Sato Pharmaceutical Co., Ltd. (690575642)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------------|---------|-----------|-------------------------------------------------------------|
| Sato Pharmaceutical Co., Ltd. | | 715699133 | manufacture(49873-019) , label(49873-019) , pack(49873-019) |

Revised: 12/2024

Sato Pharmaceutical Co., Ltd.