DAYTIME AND NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride / acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride Walgreens

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### Walgreens Daytime & Nighttime Cold and Flu

#### **DAYTIME COLD AND FLU**

### **Drug Facts**

#### 

#### Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains headache
- fever sore throat reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

### Warnings

**Liver warning** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Drug Facts** (continued)

**Allergy alert** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

**Drug Facts** (continued)

### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days
- redness or swelling is present new symptoms occur
- cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

Keep out of reach of children.

### **Overdose warning**

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- take only as directed see Overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 years & over	2 softgels with water every 4 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

# ■ when using other Nighttime or Daytime products, carefully read each label to ensure correct dosing

#### Other information

■ store at room temperature

### **Inactive ingredients**

FD&C Red #40, FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

NIGHTTIME COLD AND FLU

### **Drug Facts**

Active ingredients (in each softgel)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

#### Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep minor aches & pains headache
- fever sore throat runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

### **Warnings**

**Liver warning** This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 4 doses in 24 hours, which is the maximum daily

amount for this product

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### **NIGHTTIME COLD AND FLU**

**Drug Facts** (continued)

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleep

### Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- thyroid disease
   diabetes
   glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

■ alcohol, sedatives, and tranquilizers may increase drowsiness

#### **NIGHTTIME SEVERE COLD AND FLU SOFTGELS**

**Drug Facts** (continued)

#### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7

days **■** fever gets worse or lasts more than 3 days

- redness or swelling is present
   new symptoms occur
- cough comes back or occurs with rash or headache that lasts These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

Keep out of reach of children.

#### Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- take only as directed see Overdose warning
- do not exceed 4 doses per 24 hours

adults & children 12 years & over	2 softgels with water every 4 hours	
children 4 to under 12 years	ask a doctor	
children under 4 years	do not use	

#### Other information

store at room temperature

### **Inactive ingredients**

D&C Yellow #10, FD&C Blue #1, gelatin, glycerin, polyethylene glycol 400, povidone K30, propylene glycol, purified water, shellac, sorbitol sorbitan, sodium hydroxide, titanium dioxide

#### Questions or comments?

Call toll free: 1-888-333-9792

#### PRINCIPAL DISPLAY PANEL

DAYTIME COLD AND FLU

ACETAMINOPHEN /
PAIN RELIEVER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCI / NASAL DECONGESTANT

MAXIMUM STRENGTH

**ACTUAL SIZE** 

16SOFTGELS

NIGHTTIME COLD AND FLU

ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE / ANTIHISTAMINE
PHENYLEPHRINE HCI / NASAL DECONGESTANT

MAXIMUM STRENGTH

**ACTUAL SIZE** 

8SOFTGELS

**TOTAL 24 SOFTGELS** 



#### DAYTIME AND NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride /

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride kit

#### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0363-2218

ı	P	ackaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0363- 2218-24	1 in 1 PACKAGE, COMBINATION; Type 0: Not a Combination Product	02/06/2020	

Quant	Quantity of Parts		
Part #	Part # Package Quantity Total Product Quantity		
Part 1	2 BLISTER PACK	16	
Part 2	1 BLISTER PACK	8	

### Part 1 of 2

#### **DAYTIME COLD AND FLU**

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled

#### **Product Information**

Route of Administration ORAL

#### **Active Ingredient/Active Moiety Ingredient Name Basis of Strength** Strength ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) **ACETAMINOPHEN** 325 mg **DEXTROMETHORPHAN HYDROBROMIDE** (UNII: 9D2RTI9KYH) **DEXTROMETHORPHAN** 10 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) **HYDROBROMIDE** PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE -**PHENYLEPHRINE** 5 mg **HYDROCHLORIDE** UNII:1WS297W6MV) GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) 200 mg **GUAIFENES IN**

Inactive Ingredients		
Ingredient Name	Strength	
FD&C RED NO. 40 (UNII: WZB9127XOA)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
GELATIN (UNII: 2G86QN327L)		
GLYCERIN (UNII: PDC6A3C0OX)		

POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K30 (UNII: U725QWY32X)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6092ICV9RU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL (OBLONG)	Size	20mm
Flavor		Imprint Code	341
Contains			

ı	Pac	ckaging			
	# Item Package Description Marketing Start Marketing End Code Date Date				Marketing End Date
	1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
OTC monograph final	part341	02/06/2020		

### Part 2 of 2

### **NIGHTTIME COLD AND FLU**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

### **Product Information**

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients		
Ingredient Name	Strength	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
GELATIN (UNII: 2G86QN327L)		
GLYCERIN (UNII: PDC6A3C0OX)		
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)		
POVIDONE K30 (UNII: U725QWY32X)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SHELLAC (UNII: 46N107B710)		
SODIUM HYDROXIDE (UNII: 55X04QC32I)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics			
Color	green	Score	no score
Shape	OVAL (OBLONG)	Size	20mm
Flavor		Imprint Code	116
Contains			

F	Packaging			
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1	L	8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/06/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/04/2020	

## Labeler - Walgreens (008965063)

# **Registrant -** Spirit Pharmaceuticals LLC (179621011)

Establishment				
Name	Address	ID/FEI	<b>Business Operations</b>	
MEDGEL PVT LTD		677385498	manufacture(0363-2218)	

Revised: 8/2021 Walgreens