

**ANTI-DIARRHEAL- loperamide hcl tablet**  
**BI-MART**

-----

***Active ingredient (in each caplet)***

Loperamide HCl 2 mg

***Purpose***

Anti-diarrheal

***Use***

Controls symptoms of diarrhea, including Traveler's Diarrhea

***Warnings***

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

- if you have bloody or black stool.

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **drink plenty of clear liquids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

<b>adults and children 12 years and over</b>	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
<b>children 9-11 years (60-95 lbs)</b>	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
<b>children 6-8 years (48-59 lbs)</b>	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
<b>children 2-5 years (34-47 lbs)</b>	ask a doctor
<b>children under 2 years (up to 33 lbs)</b>	do not use

**Other information**

- do not use if seal around caplet is broken or damaged
- store between 20° and 25°C (68° and 77°F)

**Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, dicalcium phosphate dihydrate, FD&C blue #1 brilliant blue lake, magnesium stearate, microcrystalline cellulose, silica gel

**Questions?**

Call toll-free **1-844-912-4012**

**Principal Display Panel-12 Caplets**

**NDC 37835-131-12**

Compare to active ingredient in Imodium<sup>®</sup> A-D †

**ANTI-DIARRHEAL**

**CAPLETS\***

## ANTI-DIARRRHEAL

Distributed by **BI-MART**, Eugene, OR 97402



ANTI-DIARRHEAL

CAPLETS\*

Loperamide HCl Tablets, 2 mg

ANTI-DIARRHEAL

Controls the symptoms of diarrhea

Original Prescription Strength

100 Caplets\*

Each caplet (\*capsule shaped tablet contains Loperamide HCl, 2 mg)

†This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Imodium® A-D.

Distributed by **BI-MART**, Eugene, OR 97402

0 71357 91154 7

**BI-MART** NDC 37835-131-01

Compare to the active ingredient in Imodium® A-D†

**ANTI-DIARRHEAL**  
LOPERAMIDE HYDROCHLORIDE CAPLETS, 2 mg

Each caplet contains Loperamide HCl, 2 mg

**ORIGINAL PRESCRIPTION STRENGTH**

**CONTROLS SYMPTOMS OF DIARRHEA**

**100 Caplets**

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

**Drug Facts**

**Active ingredient**  
(in each caplet)  
Loperamide HCl 2 mg.....Anti-diarrheal

**Purpose**  
Use controls symptoms of diarrhea, including Traveler's Diarrhea

**Warnings**  
**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.  
**Heart alert:** Taking more than directed can cause serious heart problems or death.  
**Do not use:** ■ if you have bloody or black stool  
**Ask a doctor before use if you have:** ■ fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm  
**Ask a doctor before use if you are taking a** prescription drug. Loperamide may interact with certain prescription drugs.  
**When using this product** tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.  
**Stop use and ask a doctor if:** ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition.

PEEL FOR DIRECTIONS 131-01-BM 02/25  
LIFT HERE

<b>ANTI-DIARRHEAL</b>			
loperamide hcl tablet			
<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-131
Route of Administration	ORAL		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name		Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII: 6X9OC3H4II)		LOPERAMIDE HYDROCHLORIDE	2 mg
<b>Inactive Ingredients</b>			
Ingredient Name			Strength
STARCH, CORN (UNII: O8232NY3SJ)			
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)			

DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)				
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
Product Characteristics				
Color	green (light)	Score	2 pieces	
Shape	OVAL	Size	10mm	
Flavor		Imprint Code	44;375	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-131-12	2 in 1 CARTON	05/01/2025	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:37835-131-24	4 in 1 CARTON	05/01/2025	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:37835-131-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2025	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA076497		05/01/2025	

**Labeler -** BI-MART (027630078)

**Registrant -** BI-MART (027630078)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	manufacture(37835-131)