


BENZONATATE- benzonatate capsule
RedPharm Drug, Inc.

Benzonatate 100mg

DESCRIPTION

Benzonatate capsules, USP, a non-narcotic oral antitussive agent, is 2, 5, 8, 11, 14, 17, 20, 23, 26- onaoxaoctacosan-28-ylp-(butylamino) benzoate; with a molecular weight of 603.7.



Each benzonatate capsules, USP for oral administration contains 100 mg or 200 mg benzonatate, USP. In addition, each capsule contains the following inactive ingredients: D&C Yellow #10, gelatin, glycerin, lecithin, light mineral oil, propylene glycol, purified water, shellac glaze, titanium dioxide, and white edible ink.

CLINICAL PHARMACOLOGY

Benzonatate capsules act peripherally by anesthetizing the stretch receptors located in the respiratory passages, lungs, and pleura by dampening their activity and thereby reducing the cough reflex at its source. It begins to act within 15 to 20 minutes and its effect lasts for 3 to 8 hours. Benzonatate capsules have no inhibitory effect on the respiratory center in recommended dosage.

INDICATIONS AND USAGE

Benzonatate capsules, USP are indicated for the symptomatic relief of cough.

CONTRAINDICATIONS

Hypersensitivity to benzonatate or related compounds.

WARNINGS

Hypersensitivity

Severe hypersensitivity reactions (including bronchospasm, laryngospasm and cardiovascular collapse) have been reported which are possibly related to local anesthesia from sucking or chewing the capsule instead of swallowing it. Severe

reactions have required intervention with vasopressor agents and supportive measures.

Psychiatric Effects

Isolated instances of bizarre behavior, including mental confusion and visual hallucinations, have also been reported in patients taking Benzonatate capsules in combination with other prescribed drugs.

Accidental Ingestion and Death in Children

Keep benzonatate capsules out of reach of children. Accidental ingestion of benzonatate capsules resulting in death has been reported in children below age 10. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. If accidental ingestion occurs, seek medical attention immediately (see **OVERDOSAGE**).

PRECAUTIONS

Benzonatate is chemically related to anesthetic agents of the para-amino-benzoic acid class (e.g. procaine; tetracaine) and has been associated with adverse CNS effects possibly related to a prior sensitivity to related agents or interaction with concomitant medication.

Information for Patients:

Swallow benzonatate capsules whole. Do not break, chew, dissolve, cut, or crush benzonatate capsules. Release of benzonatate from the capsule in the mouth can produce a temporary local anesthesia of the oral mucosa and choking could occur. If numbness or tingling of the tongue, mouth, throat, or face occurs, refrain from oral ingestion of food or liquids until the numbness has resolved. If the symptoms worsen or persist, seek medical attention.

Keep benzonatate capsules out of reach of children. Accidental ingestion resulting in death has been reported in children. Signs and symptoms of overdose have been reported within 15-20 minutes and death has been reported within one hour of ingestion. Signs and symptoms may include restlessness, tremors, convulsions, coma and cardiac arrest. If accidental ingestion occurs, seek medical attention immediately.

Overdosage resulting in death may occur in adults.

Do not exceed a single dose of 200 mg and a total daily dosage of 600 mg. If you miss a dose of benzonatate capsules, skip that dose and take the next dose at the next scheduled time. Do not take 2 doses of benzonatate capsules at one time.

Usage in Pregnancy

PREGNANCY CATEGORY C

Animal reproduction studies have not been conducted with benzonatate capsules. It is also not known whether benzonatate capsules can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Benzonatate capsules should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk caution should be exercised when benzonatate capsules are administered to a nursing woman.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity, mutagenicity, and reproduction studies have not been conducted with benzonatate capsules..

Pediatric Use

Safety and effectiveness in children below the age of 10 have not been established. Accidental ingestion resulting in death has been reported in children below age 10. Keep out of reach of children.

ADVERSE REACTIONS

Potential Adverse Reactions to benzonatate capsules may include:

Hypersensitivity reactions including bronchospasm, laryngospasm, cardiovascular collapse possibly related to local anesthesia from chewing or sucking the capsule.

CNS

sedation; headache; dizziness; mental confusion; visual hallucinations.

GI

constipation; nausea; GI upset.

Dermatologic

pruritus; skin eruptions.

Other

nasal congestion; sensation of burning in the eyes; vague “chilly” sensation; numbness of the chest; hypersensitivity.

Deliberate or accidental overdose has resulted in death, particularly in children.

Hypersensitivity reactions

including bronchospasm, laryngospasm, cardiovascular collapse possibly related to local anesthesia from chewing or sucking the capsule.

OVERDOSAGE

Intentional and unintentional overdose may result in death, particularly in children.

The drug is chemically related to tetracaine and other topical anesthetics and shares various aspects of their pharmacology and toxicology. Drugs of this type are generally well absorbed after ingestion.

Signs and Symptoms

The signs and symptoms of overdose of benzonatate capsules have been reported within 15-20 minutes. If capsules are chewed or dissolved in the mouth, oropharyngeal anesthesia will develop rapidly, which may cause choking and airway compromise.

CNS stimulation may cause restlessness and tremors which may proceed to clonic convulsions followed by profound CNS depression. Convulsions, coma, cerebral edema and cardiac arrest leading to death have been reported within 1 hour of ingestion.

Treatment

In case of overdose, seek medical attention immediately. Evacuate gastric contents and administer copious amounts of activated charcoal slurry. Even in the conscious patient, cough and gag reflexes may be so depressed as to necessitate special attention to protection against aspiration of gastric contents and orally administered materials. Convulsions should be treated with a short-acting barbiturate given intravenously and carefully titrated for the smallest effective dosage. Intensive support of respiration and cardiovascular-renal function is an essential feature of the treatment of severe intoxication from overdosage.

Do not use CNS stimulants.

DOSAGE AND ADMINISTRATION

Adults and Children over 10 years of age: Usual dose is one 100 mg or 200 mg capsule three times a day as needed for cough. If necessary to control cough, up to 600 mg daily in three divided doses may be given. **Benzonatate capsules should be swallowed whole.** Benzonatate capsules are not to be broken, chewed, dissolved, cut or crushed.

HOW SUPPLIED

Benzonatate Capsules USP, 100 mg are clear yellow, oval-shaped softgel capsules imprinted with "PC14" and are supplied as follows:

NDC 42806-714-01 in bottle of 100 capsules

NDC 42806-714-05 in bottle of 500 capsules

Benzonatate Capsules USP, 200 mg are clear yellow, oval-shaped softgel capsules imprinted with "PC15" and are supplied as follows:

NDC 42806-715-01 in bottle of 100 capsules

NDC 42806-715-05 in bottle of 500 capsules

Store at 20° to 25° C (68° to 77° F) [see USP Controlled Room Temperature].

PROTECT FROM LIGHT

Dispense in tight (USP), child-resistant containers.

Rx Only

Manufactured by:

Humanwell PuraCap Pharmaceutical

Wuhan, Hubei 430206, China

Distributed by:

Epic Pharma, LLC

Laurelton, NY 11413

Rev. 10-2018-00

PI-EBEN-00

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL - 100 mg 500ct

No Varnish	DOSAGE AND USE: See accompanying prescribing information.	NDC 42806-714-01	EACH SOFTGEL (SOFT GELATIN CAPSULE) CONTAINS: Benzonatate, USP ... 100 mg
	Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature] Protect from light.	Benzonatate Capsules, USP 100 mg Rx Only 100 CAPSULES	Dispense in a tight (USP), child-resistant container. CAUTION: KEEP OUT OF THE REACH OF CHILDREN.
	EBEN01-00 Rev. 10-2018-00		 N 3 42806-714-01 0
	Distributed by: Epic Pharma, LLC Laurelton, NY 11413	Manufactured by: Humanwell PuraCap Pharmaceutical Wuhan, Hubei 430206, China	

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL - 200 mg 500ct

DOSAGE AND USE: See accompanying prescribing information.

Store at 20° to 25°C (68° to 77°F).
[See USP Controlled Room Temperature]
Protect from light.

EBEN05-00

Rev. 10-2018-00

NDC 42806-715-05

Benzonatate Capsules, USP

200 mg

Rx Only

500 CAPSULES

EACH SOFTGEL (SOFT GELATIN CAPSULE) CONTAINS:

Benzonatate, USP 200 mg

Dispense in a tight (USP), child-resistant container.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

Distributed by:
Epic Pharma, LLC
Laurelton, NY 11413




Manufactured by:
Humanwell PuraCap Pharmaceutical
Wuhan, Hubei 430206, China

No Varnish



42806-715-05 5

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL




NDC: 67296-1782-7

BENZONATATE

100MG

Rx Only 15 Capsules

Lot: 005623X1 Exp: 11/21



67296 17827 2

Usual adult dosage: See package insert

Store at controlled room temperature: 20-25 C (68-77 F)

Mfg: Humanwell PuraCap Pharmaceuticals
Wuhan Hubei 430206 China
42806-714-05

Dist. by: Redpharm Drug Eden Prairie, MN 55344

SIN 69229

(01)00367296178272
(21)1000000000000371
(17)211130
(10)005623X1

BENZONATATE

benzonatate capsule

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67296-1782(NDC:42806-714)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZONATATE (UNII: 5P4DHS6ENR) (BENZONATATE - UNII:5P4DHS6ENR)	BENZONATATE	100 mg

Inactive Ingredients	
Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics			
Color	yellow	Score	no score
Shape	OVAL	Size	7mm
Flavor		Imprint Code	PC14
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67296-1782-7	15 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206948	01/01/2020	

Labeler - RedPharm Drug, Inc. (828374897)

Establishment			
Name	Address	ID/FEI	Business Operations
RedPharm Drug, Inc.		828374897	repack(67296-1782)

Revised: 7/2024

RedPharm Drug, Inc.