

SUDAFED PE SINUS CONGESTION- diphenhydramine hydrochloride and phenylephrine hydrochloride
Johnson & Johnson Consumer Inc.

SUDAFED PE SINUS CONGESTION

SUDAFED PE[®] DAYTIME

Drug Facts

Active ingredient (in each tablet)

Phenylephrine HCl 10 mg

Purpose

Nasal decongestant

Uses

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

adults and children 12 years and over	<ul style="list-style-type: none">take 1 tablet every 4 hoursdo not take more than 6 tablets in 24 hours
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F)
- do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, D&C yellow no. 10 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

Questions or comments?

call **1-888-217-2117** (toll-free) or **215-273-8755** (collect)

SUDAFED PE[®] NIGHTTIME

Drug Facts

Active ingredients (in each tablet)

	Purpose
Diphenhydramine HCl 25 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
 - nasal congestion
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - nasal congestion

- temporarily relieves sinus congestion and pressure

Warnings

Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not exceed recommended dose**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

adults and children 12 years and over	<ul style="list-style-type: none"> take 1 tablet every 4 hours do not take more than 6 tablets in 24 hours
children under 12	ask a doctor

years

ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **do not use if blister unit is torn or broken**

Inactive ingredients

carnauba wax, FD&C blue no. 1 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

Questions or comments?

call **1-888-217-2117** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

PREVIOUSLY SUDAFED PE[®] DAY + NIGHT SINUS CONGESTION
NDC 50580-239-01

SUDAFED PE[®]

SINUS CONGESTION

Phenylephrine HCl
Nasal Decongestant

DAYTIME

- NASAL
CONGESTION

actual size

12 TABLETS 10 mg each

Diphenhydramine HCl, Phenylephrine HCl
Antihistamine, Nasal Decongestant

NIGHTTIME

- NASAL CONGESTION
- RUNNY NOSE

actual size

8 TABLETS | TOTAL: 20 TABLETS

The makers of the SUDAFED® family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

PREVIOUSLY SUDAFED PE® DAY + NIGHT SINUS CONGESTION NDC 50580-239-01

SUDAFED^{PE}

SINUS CONGESTION

Phenylephrine HCl
Nasal Decongestant

Diphenhydramine HCl, Phenylephrine HCl
Antihistamine, Nasal Decongestant



DAYTIME

• NASAL CONGESTION

actual size

12 TABLETS 10 mg each

NIGHTTIME

• NASAL CONGESTION
• RUNNY NOSE

actual size

8 TABLETS | TOTAL: 20 TABLETS

30042428



SUDAFED^{PE}

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&JCI 2018
Daytime: Active ingredient made in Germany | Nighttime: Made in Italy

Does not contain Pseudoephedrine

SUDAFED^{PE} NIGHTTIME

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional for use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222)

Directions

adults and children	take 1 tablet every 4 hours
children 12 years and over	do not take more than 6 tablets in 24 hours
children under 12 years	ask a doctor

Other information

- store between 20-25° C (68-77° F)
- do not use if blister unit is torn or broken

Inactive ingredients - camphor, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, xylitol, titanium dioxide

Questions or comments?
call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

DO NOT USE IF BLISTER UNIT IS TORN OR BROKEN

SUDAFED^{PE} SINUS CONGESTION

NDC 50580-239-01

DAYTIME

Phenylephrine HCl
Nasal Decongestant

12 TABLETS 10 mg each

NIGHTTIME

Diphenhydramine HCl, Phenylephrine HCl
Antihistamine, Nasal Decongestant

8 TABLETS

TOTAL: 20 TABLETS

SUDAFED^{PE}

Stop use and ask a doctor if

- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Symptoms do not improve within 7 days or occur with a fever

▲

Questions or comments?
call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, xylitol, titanium dioxide

<p>Drug Facts (continued)</p> <p>When using this product do not exceed recommended dose</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ nervousness, dizziness, or sleepiness occur ■ symptoms do not improve within 7 days or occur with a fever <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions</p> <table border="1"> <tr> <td>adults and children</td> <td>■ take 1 tablet every 4 hours</td> </tr> <tr> <td>12 years and over</td> <td>■ do not take more than 6 tablets in 24 hours</td> </tr> <tr> <td>children under 12 years</td> <td>ask a doctor</td> </tr> </table> <p>Other information</p> <ul style="list-style-type: none"> ■ store between 20-25°C (68-77°F) ■ do not use if blister units torn or broken <p>Inactive ingredients carnauba wax, D & C yellow no. 10 aluminum</p>	adults and children	■ take 1 tablet every 4 hours	12 years and over	■ do not take more than 6 tablets in 24 hours	children under 12 years	ask a doctor	<p>Drug Facts (continued)</p> <p>If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ do not exceed recommended dose ■ marked drowsiness may occur ■ avoid alcoholic drinks
adults and children	■ take 1 tablet every 4 hours						
12 years and over	■ do not take more than 6 tablets in 24 hours						
children under 12 years	ask a doctor						

Important: Read all product information before using. Keep this box for important information.

Do not take the day and night tablets at the same time. Do not take more than a total of 6 tablets in a 24-hour period. Take only as directed.

<p>SUDAFED PE® DAYTIME</p> <p>Drug Facts</p> <table border="1"> <tr> <th>Active ingredient (in each tablet)</th> <th>Purpose</th> </tr> <tr> <td>Phenylephrine HCl 10 mg.....</td> <td>Nasal decongestant</td> </tr> </table> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves sinus congestion and pressure ■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies <p>Warnings</p> <p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland 	Active ingredient (in each tablet)	Purpose	Phenylephrine HCl 10 mg.....	Nasal decongestant	<p>SUDAFED PE® NIGHTTIME</p> <p>Drug Facts</p> <table border="1"> <tr> <th>Active ingredients (in each tablet)</th> <th>Purpose</th> </tr> <tr> <td>Diphenhydramine HCl 25 mg.....</td> <td>Antihistamine</td> </tr> <tr> <td>Phenylephrine HCl 10 mg.....</td> <td>Nasal decongestant</td> </tr> </table> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: <ul style="list-style-type: none"> ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ■ nasal congestion ■ temporarily relieves these symptoms due to the common cold: <ul style="list-style-type: none"> ■ runny nose ■ sneezing ■ nasal congestion ■ temporarily relieves sinus congestion and pressure <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ to make a child sleepy ■ with any other product containing diphenhydramine, even one used on skin 	Active ingredients (in each tablet)	Purpose	Diphenhydramine HCl 25 mg.....	Antihistamine	Phenylephrine HCl 10 mg.....	Nasal decongestant
Active ingredient (in each tablet)	Purpose										
Phenylephrine HCl 10 mg.....	Nasal decongestant										
Active ingredients (in each tablet)	Purpose										
Diphenhydramine HCl 25 mg.....	Antihistamine										
Phenylephrine HCl 10 mg.....	Nasal decongestant										

LIFT HERE

SUDAFED PE SINUS CONGESTION

diphenhydramine hydrochloride and phenylephrine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-239
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-239-01	1 in 1 PACKAGE	06/17/2019	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
--------	------------------	------------------------

Part 1	2 BLISTER PACK	12
Part 2	2 BLISTER PACK	8

Part 1 of 2

SUDAFED PE SIINUS CONGESTION

phenylephrine hydrochloride tablet, film coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	WL;80;PE
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 CARTON		

1	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
---	--	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/17/2019	

Part 2 of 2

SUDAFED PE SINUS CONGESTION PLUS ALLERGY

diphenhydramine hydrochloride and phenylephrine hydrochloride tablet, film coated

Product Information

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	PE;WL95
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 CARTON		
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/17/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/17/2019	

Labeler - Johnson & Johnson Consumer Inc. (878046358)

Revised: 1/2024

Johnson & Johnson Consumer Inc.