

**ADULT LOW DOSE PAIN RELIEVER- aspirin tablet, delayed release
TWIN MED LLC**

Adult Low Dose Aspirin Delayed Release Tablets, 81 mg

Drug Facts

Active ingredients (in each tablet)

Aspirin 81 mg (NSAID)**

**nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or symptoms needing immediate relief.**

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

- This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - redness or swelling present
 - new symptoms occur
 - ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- **adults and children 12 years and over:**
 - take 4 to 8 tablets every 4 hours, not to exceed 48 tablets in 24 hours unless directed by a doctor
- **children under 12 years:**consult a doctor

Other information

- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-68°F).
- **DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED**

Inactive ingredients

Starch, Povidone, Microcrystalline Cellulose, Croscarmellose Sodium, Stearic Acid, Colloidal Silicon Dioxide, Methacrylic Acid and Ethyl acrylate Co-polymer, Triethyl Citrate, Talc, Fumed Silica, Titanium Dioxide, D&C Yellow #10, FD&C Yellow #6.

Questions or comments?

call toll-free **1-844-912-4012**

Distributed by:**Twin Med LLC**

Santa Fe Springs,

CA 90670

www.procureproducts.com

Made in the USA**PRINCIPAL DISPLAY PANEL****NDC 55681-403-03**

*Compare to active ingredients in Bayer[®] Aspirin Low Dose 81 mg

Adult Low Dose Pain Reliever Aspirin Delayed Release Tablets, 81 mg

Pain Reliever (NSAID)

Enteric Coated**300 Tablets**

*This product is not manufactured or distributed by the owner of the registered trademark Bayer[®].

0 71357 91112 7



PROCURE
HEALTHCARE PRODUCTS

*Compare to active ingredient in
BAYER® ASPIRIN LOW DOSE 81 MG

ASPIRIN
ADULT LOW DOSE
PAIN RELIEVER (NSAID)**



300 TABLETS | 81 MG
ENTERIC COATED

NDC 55681-403-03

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 81mg (NSAID)* Pain reliever
**nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

PEEL FOR DIRECTIONS

LIFT HERE

Drug Facts, (cont'd)

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ drink a full glass of water with each dose
■ **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours, not to exceed 48 tablets in 24 hours unless directed by a doctor
■ **children under 12 years:** consult a doctor

Other information ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients
Starch, Povidone, Microcrystalline Cellulose, Croscarmellose Sodium, Stearic Acid, Colloidal Silicon Dioxide, Methacrylic acid and ethyl acrylate co-polymer, Triethyl Citrate, Talc, Fumed Silica, Titanium Dioxide, D&C Yellow #10, FD&C Yellow #6

Questions? call toll-free 1-844-912-4012

*This product is not manufactured or distributed by the owner of the registered trademark Bayer®

Distributed by: Twin Med LLC
Santa Fe Springs, CA 90670
www.procureproducts.com

MADE IN THE USA
ITEM# PC0TC190

GLUE
0.75"

ADULT LOW DOSE PAIN RELIEVER

aspirin tablet, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55681-403
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients		
Ingredient Name		Strength
STARCH, CORN (UNII: O8232NY3SJ)		
POVIDONE (UNII: FZ989GH94E)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		

METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	A1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55681-403-03	300 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/01/2024	

Labeler - TWIN MED LLC (009579330)

Establishment

Name	Address	ID/FEI	Business Operations
ELYSIUM PHARMACEUTICALS LIMITED		915664486	manufacture(55681-403)

Revised: 7/2025

TWIN MED LLC