

IBUPROFEN- ibuprofen tablet, coated
BETTER LIVING BRANDS LLC.

Ibuprofen
Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)*
See New Warnings

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

■ you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions (Caplets)

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F). [See USP Controlled Room Temperature].
- do not use if foil inner seal is broken or missing

Inactive ingredients

colloidal silicon dioxide, FD&C Yellow #6, maize starch, polyethylene glycol, polyvinyl alcohol, povidone k30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide


Questions or comments?

call 1-877-770-3183 weekdays 9:00 AM to 4:30 PM EST.


Caplets

Kaj
Pre
nt

55821 RD 19183




3 21130 79862 0



Scan here for more information

DISTRIBUTED BY
BETTER LIVING BRANDS LLC
PO BOX 95
PLEASANTON, CA 94686-0095
1-800-723-3929
www.betterlivingbrandsllc.com

OUR PROMISE
100% GUARANTEED



Quality Guaranteed

Ibuprofen

TABLETS USP, 200 mg

Pain Reliever/
Fever Reducer
(NSAID)*


Contains no ingredient made from a gluten-containing grain (wheat, barley, or rye)

See New Warnings

Compare to active ingredient of Motrin® IB†

NDC 2130-358-05

Actual Size



50 COATED CAPLETS**
(*Capsule-Shaped Tablets)

Drug Facts (continued)

Questions or comments?
Call 1-877-770-3183 weekdays 9:00 AM to 4:30 PM EST

†This product is not manufactured or distributed by Michael Consumer Healthcare Division, the owner of the registered trademark Motrin® IB

MADE IN INDIA


Lot _____ Exp. _____

COATING FREE AREA

COATING FREE AREA


R-01

300000449
547000004



COATING FREE AREA


COATING FREE AREA



COATING FREE AREA

31

090909 0909




300000449
547000004

R-01


COATING FREE AREA

COATING FREE AREA




0 1 0 9 9 6 3 1 3

COATING FREE AREA



COATING FREE AREA

COATING FREE AREA



300000449
547000004

R-01

COATING FREE AREA

Important: Read all product information before using. Keep this box for important information. Do not use if the inner seal is broken or missing.

Drug Facts

Active ingredient (in each caplet): Purpose
Ibuprofen 200 mg.....Pain reliever/fever reducer (NSAID)*

Uses

- Temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - backache
 - backache
 - the common cold
 - menstrual cramps
- Temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, or skin redness/itching or rash in babies

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older

- Have had stomach ulcers or bleeding problems
- Take other drugs containing aspirin or steroid drug
- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, rofecoxib, or others)
- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed

Drug Facts (continued)

If you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better

If you have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling or pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area

Directions

- Do not take more than directed
- Take 1 caplet every 4 to 6 hours while symptoms persist
- If pain or fever does not respond to 1 caplet, 2 caplets may be used
- Do not exceed 6 caplets in 24 hours, unless directed by a doctor
- Ask a doctor

Other information

- Children under 12 years:
- Children 12 years and older:
- Ask a doctor

Directions

- Do not take more than directed
- Take 1 caplet every 4 to 6 hours while symptoms persist
- If pain or fever does not respond to 1 caplet, 2 caplets may be used
- Do not exceed 6 caplets in 24 hours, unless directed by a doctor
- Ask a doctor

Other information

- Children under 12 years:
- Children 12 years and older:
- Ask a doctor

Directions

- Do not take more than directed
- Take 1 caplet every 4 to 6 hours while symptoms persist
- If pain or fever does not respond to 1 caplet, 2 caplets may be used
- Do not exceed 6 caplets in 24 hours, unless directed by a doctor
- Ask a doctor

Other information

- Children under 12 years:
- Children 12 years and older:
- Ask a doctor

Drug Facts (continued)

Heart attack and stroke warnings (NSAIDs), except aspirin

These can be fatal. The risk is higher if you use more than directed or for longer than directed

Do not use

- If you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- Right before or after heart surgery

Ask a doctor before use if

- You have problems or serious side effects from taking pain relievers or fever reducers
- You have a history of stomach problems, such as heartburn, indigestion, ulcers, or acid reflux
- You have high blood pressure, heart disease, liver disease, kidney disease, asthma, or had a stroke
- You are taking a diuretic

Ask a doctor or pharmacist before use if you are

- Taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- Under a doctor's care for any serious condition
- Taking any other drug

When using this product

- Take with food or milk if stomach upset occurs
- Stop use and ask a doctor if
- You experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- You have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling or pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	orange (Caplet shaped film coated tablets)	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-358-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	07/12/2019	
2	NDC:21130-358-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/12/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202312	07/12/2019	

Labeler - BETTER LIVING BRANDS LLC. (009137209)