

**IBUPROFEN- ibuprofen tablet, coated**  
**BETTER LIVING BRANDS LLC.**

-----  
**Ibuprofen**  
**Tablets USP, 200 mg**  
**Pain Reliever/Fever Reducer (NSAID)\***  
**See New Warnings**

**Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**Allergy alert:**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

### **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack and stroke warning:**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition

- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of the reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions (Tablets)**

- do not take more than directed
- the smallest effective dose should be used

Adults and children 12 years and older

- take 1 Tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

### **Other information**

- store between 20-25°C (68-77°F). [See USP Controlled Room Temperature].
- do not use if foil inner seal is broken or missing

### **Inactive ingredients**

colloidal silicon dioxide, FD&C Yellow #6, maize starch, polyethylene glycol, polyvinyl alcohol, povidone k30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

### **Questions or comments?**

call **1-877-770-3183** weekdays 9:00 AM to 4:30 PM EST.

### **Tablets**

end user : GRANULES  
 item : 1250 CHANGE TAB 100CG  
 code # :  
 size : 147/8 X 147/8 X 3+13/16  
 ref # : 1P158068  
 material : 016 S35

**OUR PROMISE**  
 100% GUARANTEED  
 SOLIDITY & COMPLIANCE  
 QUALITY CONTROLLED

DISTRIBUTED BY  
 BETTER LIVING BRANDS LLC.  
 P.O. BOX 99  
 PLEASANTON, CA 94669-0099  
 1-888-723-3829  
 www.betterlivingbrandsllc.com



Important: Read all product information before using. Keep this box for important information. Do not use if full review sheet is broken or missing.

**Drug Facts**  
**Active ingredient** **Purpose**  
 (in each tablet) **Ibuprofen 200 mg** Pain reliever/fever reducer  
**NSAID**†  
 †Nonsteroidal anti-inflammatory drug

**Uses**  
 temporarily relieves minor aches and pains due to:  
 • headache • muscular aches • minor pain of arthritis • toothache • backache • the common cold • menstrual cramps  
 temporarily reduces fever

**Warnings**  
 Always alert: Ibuprofen may cause severe allergic reaction, especially in people allergic to aspirin.  
 Symptoms may include:  
 • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters  
 If an allergic reaction occurs, stop use and seek medical help right away.  
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
 • are age 60 or older  
 • have had stomach ulcers or bleeding problems  
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
 • have 3 or more alcoholic drinks every day while using this product  
 • take more or for a longer time than directed  
 Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke.

**Ibuprofen**  
 TABLETS USP, 200 mg  
 Pain Reliever/  
 Fever Reducer  
 (NSAID)<sup>†</sup>  
 Contains no harmful metals  
 from a gluten-containing grain  
 (wheat, barley, or rye)  
 See New Warnings

Actual Size

Compare to  
 Motrin® IB  
 (NSAID) 200 mg

100 COATED TABLETS

**Drug Facts (continued)**  
 directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
 Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**  
 do not take more than directed  
 at the smallest effective dose should be used

**adults and children 12 years and older:**  
 • take 1 tablet every 4 to 6 hours while symptoms persist  
 • if pain or fever does not respond to 1 tablet, 2 tablets may be used  
 • do not exceed 6 tablets in 24 hours, unless directed by a doctor  
 • ask a doctor

**children under 12 years:**  
 • ask a doctor

**Other information**  
 • store between 20-25°C (68-77°F)

**Inactive ingredients** croscellose, silicon dioxide, FD&C yellow #6, maize starch, polyethylene glycol, polyethylene glycol, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Questions or comments?**  
 call 1-877-776-5365  
 weekdays 9:00 AM to 4:00 PM EST

†This product is not manufactured or distributed by Janssen Consumer Healthcare Division, the owner of the registered trademark Motrin® IB

MADE IN INDIA

**Drug Facts (continued)**  
 These can be fatal. The risk is higher if you use more than directed or for longer than directed

**Do not use**  
 • if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer  
 • if right before or after head surgery

**Ask a doctor before use if**  
 • you have problems or serious side effects from taking pain relievers or fever reducers  
 • in the stomach bleeding warning: applies to you  
 • you have a history of stomach problems, such as heartburn  
 • you have high blood pressure, heart disease, liver disease, kidney disease, asthma or had a stroke  
 • you are taking a diuretic  
 • you are taking or planning to take aspirin if you are taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin  
 • you are under a doctor's care for any serious condition  
 • taking any other drug

**When using this product**  
 • take with food or milk if stomach upset occurs  
 • stop use and ask a doctor if:  
 • you experience any of the following signs of stomach bleeding:  
 • feel faint • vomit blood • have bloody or black stools  
 • have stomach pain that does not get better  
 • you have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling  
 • pain gets worse or lasts more than 10 days  
 • never gets worse or lasts more than 3 days  
 • redness or swelling is present in the painful area  
 • any new symptoms appear  
 • if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely

300000444  
 547000001  
 R-02

300000444  
 547000001  
 R-02

Scan here for more information

Lot

Exp.



# IBUPROFEN

ibuprofen tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-359
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

### Product Characteristics

<b>Color</b>	orange (round shaped biconvex film coated tablets)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G;2
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-359-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	
2	NDC:21130-359-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202312	07/12/2019	

**Labeler** - BETTER LIVING BRANDS LLC. (009137209)

Revised: 3/2023

BETTER LIVING BRANDS LLC.