

FAST PAIN RELIEF- ibuprofen tablet, coated
HealthLife of USA LLC

Drug Facts

Active ingredient(in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- **do not take more than directed**
- **the smallest effective dose should be used**
- **adults and children 12 years and over:** take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- **children under 12 years:** ask a doctor

Other information

- each tablet contains: sodium 0.030 mg
- store at 20° to -25°C (68° to -77°F)

- avoid excessive heat above 40°C (104°F)
- read all product information before using
- **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, iron oxide red, lactose monohydrate, povidone k30, sodium starch glycolate, stearic acid, titanium dioxide, triacetin

Questions or Comments

Call toll free 1-844-832-1138 Monday through Friday 9AM – 5PM EST or www.healthlifeofusa.com

Principal Display Panel

COMPARE TO ACTIVE INGREDIENT IN Advil®†

Ibuprofen 200 mg

Pain Reliever/Fever Reducer (NSAID)

†This product is not manufactured or distributed by Pfizer Healthcare, owner of the registered trademark of Advil®



FAST PAIN RELIEF

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69517-111
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69517-111-25	25 in 1 CARTON	04/06/2016	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:69517-111-50	50 in 1 CARTON	04/06/2016	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:69517-111-02	2 in 1 POUCH	04/06/2016	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:69517-111-10	1000 in 1 BOTTLE	04/06/2016	
4	NDC:69517-111-05	500 in 1 BOTTLE		
4	NDC:69517-111-30	30 in 1 BOTTLE		
4		1 in 1 CARTON; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/01/2011	

Labeler - HealthLife of USA LLC (079656178)

Establishment			
Name	Address	ID/FEI	Business Operations
Granules India Limited		918609236	manufacture(69517-111)

Revised: 6/2017

HealthLife of USA LLC