

**TUKOL COUGH AND COLD HONEY FLAVOR- dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride syrup**  
**Genomma Lab USA**

-----  
**Tukol® Cough & Cold Honey Flavor Syrup**

**Active ingredients (in each 5 mL or 1 teaspoon) Purpose**

Dextromethorphan HBr, 10 mg.....Cough Suppressant  
Guaifenesin, 100 mg.....Expectorant  
Phenylephrine HCl, 5 mg.....Nasal Decongestant

**Purpose**

Cough Suppressant

Expectorant

Nasal Decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- nasal congestion
- cough due to minor throat and bronchial irritation

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are** taking any other oral nasal decongestant or stimulant

**Do not exceed recommended dosage**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product
- TSP = teaspoon

age	dose
adults and children 12 years and over	2 teaspoons (10 mL) every 4 hours
children under 12 years	do not use

**Other information**

- **each teaspoon (5 mL) contains:** sodium 3 mg
- store at 15-30°C (59-86°F). Do not refrigerate

**Inactive ingredients**

anhydrous citric acid, FD&C yellow # 6, flavor, glycerin, honey, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

**Questions or comments?**

1-877-994-3666 Monday to Friday from 8 am to 6 pm, Central time.

# Tukol<sup>®</sup>

## HONEY / MIEL

### COUGH & CONGESTION

#### TOS Y CONGESTIÓN

Dextromethorphan HBr / Guaifenesin / Phenylephrine HCl  
Dextrometorfano HBr / Guaifenesina / Fenilefrina HCl

Ages/Edades  
12+

SABOR A MIEL

NATURAL  
HONEY  
FLAVOR

Relieves/Alivia:

- COUGH / TOS
- NASAL CONGESTION / CONGESTIÓN NASAL
- CHEST CONGESTION / CONGESTIÓN DE PECHO

4 FL OZ (118 ml)

#### Drug Facts

**Active ingredients (in each 5 mL or 1 teaspoon) Purposes**  
Dextromethorphan HBr, 10 mg.....Cough Suppressant  
Guaifenesin, 100 mg.....Expectorant  
Phenylephrine HCl, 5 mg.....Nasal Decongestant

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
  - nasal congestion
  - cough due to minor throat and bronchial irritation

#### Warnings

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.**

#### Do not exceed recommended dosage.

#### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- do not take more than 6 doses in any 24 hour period
- this adult strength product is not intended for use in children under 12 years of age
- measure only with dosing cup provided
- keep dosing cup with product

age	dose
adults and children 12 years and over	2 teaspoons (10 mL) every 4 hours
children under 12 years	do not use

#### Drug Facts (continued)

#### Other information

- each teaspoon (5 mL) contains: sodium 3 mg
- store at 15-30°C (59-86°F). Do not refrigerate.

#### Inactive ingredients

anhydrous citric acid, FD&C yellow #6, flavor, glycerin, honey, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

#### Questions or comments?

1-877-994-3666 Monday to Friday from 8 am to 6 pm, Central Time.

#### Información del Medicamento

#### Ingredientes activos

(en cada 5 mL o 1 cucharadita) Propósitos  
Dextrometorfano HBr, 10 mg.....Supresor de la Tos  
Guaifenesina, 100 mg.....Expectorante  
Fenilefrina HCl, 5 mg.....Descongestionante Nasal

#### Usos

- ayuda a desprender las flemas (mucosidad) y a adelgazar las secreciones bronquiales para drenar las vías respiratorias.
- alivia temporalmente los siguientes síntomas que se presentan con el resfriado:
  - congestión nasal
  - tos debida a irritación menor de garganta y bronquios

#### Advertencias

**No utilizarlo** si usted está tomando un inhibidor de la monoamino oxidasa (IMAO) de prescripción (ciertos medicamentos para la depresión, condiciones psiquiátricas o emocionales, o enfermedad de Parkinson), o durante 2 semanas después de suspender el medicamento IMAO. Si usted desconoce si su medicamento de prescripción contiene un IMAO, consulte a su médico o farmacéutico antes de tomar este producto.

#### Consulte a su médico si usted padece

- enfermedad cardíaca
- hipertensión
- enfermedad de la tiroides
- diabetes
- dificultad al orinar debido a dilatación de la glándula prostática
- tos que se presenta con exceso de flema (mucosidad)
- tos persistente o crónica tal como ocurre con el tabaquismo, asma, bronquitis crónica o enfisema

#### Información del Medicamento (continuación)

**Consulte a su médico o farmacéutico antes de utilizarlo si usted está tomando cualquier otro descongestionante o estimulante nasal oral.**

#### No exceder la dosis recomendada.

#### Suspenda su uso y consulte a su médico si

- se siente nervioso, mareado o con insomnio
- los síntomas no mejoran en los siguientes 7 días o se acompañan de fiebre
- la tos persiste por más de 7 días, regresa, o se presenta acompañada por fiebre, erupción o dolor de cabeza persistente. Estos podrían ser signos de una condición grave.

**En caso de embarazo o lactancia**, consulte a un profesional médico antes de su uso.

**Manténgase fuera del alcance de los niños.** En caso de sobredosis, obtenga ayuda médica o contacte al Centro de Control de Envenenamiento de manera inmediata.

#### Indicaciones

- no tomar más de 6 dosis en un periodo de 24 horas
- este producto de concentración para adultos no está previsto para su uso en niños menores de 12 años de edad
- medir solo con la copa dosificadora incluida
- mantenga la copa dosificadora con este producto
- TSP = cucharadita

edad	dosis
adultos y niños 12 años de edad y mayores	2 cucharaditas (10 mL) cada 4 horas
niños menores de 12 años	no utilizar

#### Información adicional

- cada cucharadita (5 mL) contiene: sodio 3 mg
- almacenar entre 15-30 °C (59-86 °F). No refrigerar.

#### Ingredientes inactivos

ácido cítrico anhidro, FD&C amarillo #6, sabor, glicerina, miel, propilenglicol, agua purificada, benzoato de sodio, solución de sorbitol, sucralosa, goma xantana.

#### ¿Preguntas o comentarios?

1-877-994-3666 Lunes a Viernes de 8 am a 6 pm, Tiempo del Centro.

Distributed by / Distribuido por:  
Genomma Lab USA Inc.,  
Houston, TX 77098



## TUKOL COUGH AND COLD HONEY FLAVOR

dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride syrup

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50066-534
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>HONEY</b> (UNII: Y9H1V576FH)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3Z D4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>WATER</b> (UNII: 059QF0KOOR)	

### Product Characteristics

<b>Color</b>	yellow (Amber to yellow)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50066-534-04	1 in 1 CARTON	09/20/2019	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/20/2019	

**Labeler** - Genomma Lab USA (832323534)

**Registrant** - Genomma Lab USA (832323534)

Revised: 11/2024

Genomma Lab USA