

CYSTEX MAX UTI PAIN RELIEF- phenazopyridine hydrochloride tablet
Clarion Brands, LLC

Drug Facts

Active ingredient

Phenazopyridine hydrochloride 99.5 mg

Purpose

Urinary tract analgesic

Use fast relief from urinary pain, burning, urgency and frequency associated with urinary tract infections. Treatment should not exceed 2 days without consulting a doctor; see Directions.

Warnings

- Do not exceed recommended dosage
- Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.

Do not use

if you have Glucose-6-Phosphare Dehydrogenase (G6PD) deficiency unless approved by your doctor

Ask a doctor before use if you have

- kidney disease
- allergies to foods, preservatives or dyes
- had a hypersensitive reaction to phenazopyridine hydrochloride

When using this product

- do not exceed recommended dosage
- stomach upset may occur, taking this product with or after meals may reduce stomach upset
- your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.
- do not use for more than 2 days (12 tablets) without consulting a doctor

Stop use and ask a doctor if

- your symptoms last for more than 2 days

- you suspect you are having an adverse reaction to the medication

If pregnant or breast-feeding, ask a health professional before use. **A pregnancy test and consultation with a healthcare professional if pregnancy is confirmed is recommended prior to use**

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 12 years of age and older:take 2 tablets 3 times a day with a full glass of water or after meals as needed

- **children under 12 years:**
- ask a doctor

Other information

- **WARNING:** This product contains Phenazopyridine Hydrochloride, known to the state of California to cause cancer.
- this product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests
- this product may stain soft contact lenses and other items if handled after touching tablets
- store at room temperature 59°F-86°F (15°C-30°C) in a dry place and protect from light

Tamper evident:Product is sealed within blisters. Do not use if any part of the blister is torn, open or damaged.

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, silicon dioxide, sodium starch glycolate, talc, triacetin

Questions

call **1-844-297-8394**

PRINCIPAL DISPLAY PANEL

MAX STRENGTH

Cystex[®]
Urinary Health

Phenazopyridine Hydrochloride 99.5 mg

MAX UTI PAIN RELIEF

FAST-ACTING UTI PAIN RELIEF

Strongest UTI Pain Relief Available
for Fast-Acting Relief of Pain & Burning *

✓Relief of **Pain, Burning, & Urgency**

✓ **Targets Pain** at the Source of the UTI

STARTS WORKING

IN AS LITTLE AS
15 MINUTES

Cystex is not intended to replace a doctor's care.

24 Tablets

TAMPER EVIDENT: TABLETS SEALED IN BLISTER. DO NOT USE IF BLISTER IS OPENED
OR DAMAGED.

1000284

REV XX/XX REESECODE

For more information on
Cystex Max Strength UTI Pain
Relief or other products, please
visit www.cystex.com or call
844-297-8394

Distributed by: Cystex LLC
811 Broad Street, Suite 600
Chattanooga, TN 37402 ©2024

*without a prescription



CYSTEX MAX UTI PAIN RELIEF

phenazopyridine hydrochloride tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69693-514 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|----------------------------------|----------|
| PHENAZOPYRIDINE HYDROCHLORIDE (UNII: 0EWG668W17) (PHENAZOPYRIDINE - UNII:K2J09EMJ52) | PHENAZOPYRIDINE HYDROCHLORIDE | 99.5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| STARCH, CORN (UNII: O8232NY3SJ) | |

| |
|--|
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) |
| LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G) |
| MAGNESIUM STEARATE (UNII: 70097M6I30) |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A) |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) |
| SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D) |
| TALC (UNII: 7SEV7J4R1U) |
| TRIACETIN (UNII: XHX3C3X673) |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | red | Score | no score |
| Shape | ROUND | Size | 10mm |
| Flavor | | Imprint Code | CYSTEK |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:69693-514-24 | 1 in 1 CARTON | 08/02/2024 | |
| 1 | | 24 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | 505G(a)(3) | 08/02/2024 | |

Labeler - Clarion Brands, LLC (079742703)

Revised: 1/2026

Clarion Brands, LLC