

SODIUM CHLORIDE- sodium chloride injection
Nephron Pharmaceuticals Corporation

0.9% Sodium Chloride Injection, USP

DESCRIPTION

Sodium Chloride Injection, USP is a sterile, nonpyrogenic solution for fluid and electrolyte replenishment in single dose containers for intravenous administration. It contains no antimicrobial agents. The nominal pH is 5.6 (4.5 to 7.0). Composition, osmolarity, and ionic concentration are shown below:

0.9% Sodium Chloride Injection, USP contains 9 g/L Sodium Chloride, USP (NaCl) with an osmolarity of 308 mOsmol/L (calc). It contains 154 mEq/L sodium and 154 mEq/L chloride.

The plastic container is not made with PVC or DEHP.

The plastic container is made of polypropylene formulated and developed for parenteral drugs. The suitability of the plastic container has been established through biological evaluations, which have shown the container passes Class VI testing as specified in the U.S. Pharmacopeia for Biological Tests — Plastic Containers. These tests confirm the biological safety of the container closure system.

The container has two ports, one is for the intravenous administration set and the other is a medication addition site. Refer to the Directions for Use of the container to properly identify the ports.

No vapor barrier is necessary.

CLINICAL PHARMACOLOGY

Sodium Chloride Injection, USP has value as a source of water and electrolytes. It is capable of inducing diuresis depending on the clinical condition of the patient.

INDICATIONS AND USAGE

Sodium Chloride Injection, USP is indicated as a source of water and electrolytes.

0.9% Sodium Chloride Injection, USP is also indicated for use as a priming solution in hemodialysis procedures.

CONTRAINDICATIONS

None known.

WARNINGS

Hypersensitivity

Hypersensitivity and infusion reactions, including hypotension, pyrexia, tremor, chills, urticaria, rash, and pruritus have been reported with 0.9% Sodium Chloride Injection, USP.

Stop the infusion immediately if signs or symptoms of a hypersensitivity reaction develop, such as tachycardia, chest pain, dyspnea and flushing. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Electrolyte Imbalances

Fluid Overload

Depending on the volume and rate of infusion, and the patient's underlying clinical condition, intravenous administration of Sodium Chloride Injection, USP can cause fluid disturbances such as overhydration/ hypervolemia and congested states, including pulmonary congestion and edema.

Avoid 0.9% Sodium Chloride Injection, USP in patients with or at risk for fluid and/or solute overloading. If use cannot be avoided, monitor fluid balance, electrolyte concentrations, and acid base balance, as needed and especially during prolonged use.

Hyponatremia

Sodium Chloride Injection, USP may cause hyponatremia. Hyponatremia can lead to acute hyponatremic encephalopathy characterized by headache, nausea, seizures, lethargy, and vomiting. Patients with brain edema are at particular risk of severe, irreversible and life-threatening brain injury.

The risk of hospital-acquired hyponatremia is increased in patients with cardiac or pulmonary failure, and in patients with non-osmotic vasopressin release (including SIADH) treated with high volume of Sodium Chloride Injection, USP.

The risk for hyponatremia is increased in pediatric patients, elderly patients, postoperative patients, those with psychogenic polydipsia, and in patients treated with medications that increase the risk of hyponatremia (such as diuretics, certain antiepileptic and psychotropic medications). See **DRUG INTERACTIONS** .

Patients at increased risk for developing complications of hyponatremia such as hyponatremic encephalopathy, include pediatric patients, women (in particular premenopausal women), patients with hypoxemia, and patients with underlying central nervous system disease. Avoid Sodium Chloride Injection, USP in patients with or at risk for hyponatremia. If use cannot be avoided, monitor serum sodium concentrations.

Rapid correction of hyponatremia is potentially dangerous with risk of serious neurologic complications. Brain adaptations reducing risk of cerebral edema make the brain vulnerable to injury when chronic hyponatremia is too rapidly corrected, which is known as osmotic demyelination syndrome (ODS). To avoid complications, monitor serum sodium and chloride concentrations, fluid status, acid-base balance, and signs of neurologic complications.

Hypernatremia

Hypernatremia may occur with Sodium Chloride Injection, USP. Conditions that may increase the risk of hypernatremia, fluid overload and edema (central and peripheral), include patients with: primary hyperaldosteronism; secondary hyperaldosteronism associated with, for example, hypertension, congestive heart failure, liver disease (including cirrhosis), renal disease (including renal artery stenosis, nephrosclerosis); and pre-eclampsia.

Certain medications, such as corticosteroids or corticotropin, may also increase risk of sodium and fluid retention, see **DRUG INTERACTIONS** .

Avoid Sodium Chloride Injection, USP in patients with, or at risk for, hypernatremia. If use cannot be avoided, monitor serum sodium concentrations.

Rapid correction of hypernatremia is potentially dangerous with risk of serious neurologic complications. Excessively rapid correction of hypernatremia is also associated with a risk for serious neurologic complications such as osmotic demyelination syndrome (ODS) with risk of seizures and cerebral edema.

PRECAUTIONS

Patients with Severe Renal Impairment

Administration of Sodium Chloride Injection, USP in patients with or at risk of severe renal impairment, may result in hypernatremia and/or fluid overload (see **WARNINGS**). Avoid Sodium Chloride Injection, USP in patients with severe renal impairment or conditions that may cause sodium and/or potassium retention, fluid overload, or edema. If use cannot be avoided, monitor patients with severe renal impairment for development of these adverse reactions.

Drug Interactions

Other Products that Affect Fluid and/or Electrolyte Balance

Administration of Sodium Chloride Injection, USP to patients treated concomitantly with drugs associated with sodium and fluid retention may increase the risk of hypernatremia and volume overload. Avoid use of Sodium Chloride Injection, USP in patients receiving such products, such as corticosteroids or corticotropin. If use cannot be avoided, monitor serum electrolytes, fluid balance and acid-base balance.

Lithium

Renal sodium and lithium clearance may be decreased during administration of 0.45% Sodium Chloride Injection, USP. Monitor serum lithium concentrations during concomitant use.

Renal sodium and lithium clearance may be increased during administration of 0.9% Sodium Chloride Injection, USP. Monitor serum lithium concentrations during concomitant use.

Other Drugs that Increase the Risk of Hyponatremia

Administration of Sodium Chloride Injection, USP in patients treated concomitantly with medications associated with hyponatremia may increase the risk of developing hyponatremia.

Avoid use of Sodium Chloride Injection, USP in patients receiving products, such as diuretics, and certain antiepileptic and psychotropic medications. Drugs that increase the vasopressin effect reduce renal electrolyte free water excretion and may also increase the risk of hyponatremia following treatment with intravenous fluids. If use cannot be avoided, monitor serum sodium concentrations.

Pregnancy

There are no adequate and well controlled studies with Sodium Chloride Injection, USP in pregnant women and animal reproduction studies have not been conducted with this drug. Therefore, it is not known whether Sodium Chloride Injection, USP can cause fetal harm when administered to a pregnant woman. Sodium Chloride Injection, USP should be given during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

It is not known whether this drug is present in human milk. Because many drugs are present in human milk, caution should be exercised when Sodium Chloride Injection, USP is administered to a nursing woman.

Pediatric Use

The use of Sodium Chloride Injection, USP in pediatric patients is based on clinical practice. (See **DOSAGE AND ADMINISTRATION**).

Closely monitor plasma electrolyte concentrations in pediatric patients who may have impaired ability to regulate fluids and electrolytes. In very low birth weight infants, excessive or rapid administration of Sodium Chloride Injection, USP may result in increased serum osmolality and risk of intracerebral hemorrhage.

Children (including neonates and older children) are at increased risk of developing hyponatremia as well as for developing hyponatremic encephalopathy.

Geriatric Use

Geriatric patients are at increased risk of developing electrolyte imbalances. Sodium Chloride Injection, USP is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Therefore, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy. Consider monitoring renal function in elderly patients.

ADVERSE REACTIONS

Post-Marketing Adverse Reactions

The following adverse reactions have been identified during postapproval use of Sodium Chloride Injection, USP. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

The following adverse reactions have been reported in the post-marketing experience during use of Sodium Chloride Injection, USP and include the following:

General disorders and administration site conditions : Infusion site erythema, injection site streaking, burning sensation, and infusion site urticaria

Hypersensitivity reactions : Hypotension, pyrexia, tremor, chills, urticaria, rash, and pruritus.

Metabolism and nutrition disorders : Hyponatremia*, hyponatremia, hyperchloremic metabolic acidosis.

Nervous System Disorders : Hyponatremic encephalopathy

*Adverse reaction of hyponatremia is only related to 0.9% Sodium Chloride Injection, USP.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate therapeutic countermeasures and save the remainder of the fluid for examination if deemed necessary.

OVERDOSAGE

Excessive administration of:

- 0.9% Sodium Chloride Injection, USP can cause hyponatremia.
- Sodium Chloride Injection, USP can cause fluid overload (which can lead to pulmonary and/or peripheral edema). See **WARNINGS** and **ADVERSE REACTIONS** .

When assessing an overdose, any additives in the solution must also be considered. The effects of an overdose may require immediate medical attention and treatment.

Interventions include discontinuation of Sodium Chloride Injection, USP administration, dose reduction, and other measures as indicated for the specific clinical constellation (e.g., monitoring of fluid balance, electrolyte concentrations and acid base balance).

DOSAGE AND ADMINISTRATION

Important Administration Instructions

- Sodium Chloride Injection, USP is intended for intravenous administration using sterile equipment.
- Do not connect flexible plastic containers in series in order to avoid air embolism due to possible residual air contained in the primary container.
- Set the vent to the closed position on a vented intravenous administration set to prevent air embolism.
- Use a dedicated line without any connections to avoid air embolism.
- Do not pressurize intravenous solutions contained in flexible plastic containers to increase flow rates in order to avoid air embolism due to incomplete evacuation of residual air in the container.
- Prior to infusion, visually inspect the solution for particulate matter and discoloration.
- The solution should be clear and there should be no precipitates. Do not administer unless solution is clear, and container is undamaged.

Dosing Information

The choice of product, dosage, volume, rate, and duration of administration is dependent upon the age, weight and clinical condition of the patient and concomitant therapy, and administration should be determined by a physician experienced in intravenous fluid therapy.

Introduction of Additives

Additives may be incompatible.

Evaluate all additions to the plastic container for compatibility and stability of the resulting preparation. Consult with a pharmacist, if available.

If, in the informed judgment of the physician, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced. After addition, if there is a discoloration and/or the appearance of precipitates, insoluble complexes or crystals, do not use. Do not store solutions containing additives. Discard any unused portion.

HOW SUPPLIED

The available sizes of 0.9% Sodium Chloride Injection, USP are shown below:

- NDC 0487-4301-05 - 50 mL, 64 units per shipper
- NDC 0487-4301-10 - 100 mL, 64 units per shipper
- NDC 0487-4301-25 - 250 mL, 35 units per shipper
- NDC 0487-4301-50 - 500 mL, 18 units per shipper

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended the product be stored at room temperature (25° C/ 77° F); brief exposure up to 40° C/ 104°F does not adversely affect the product.

DIRECTIONS FOR USE

For Information on Risk of Air Embolism - see **DOSAGE AND ADMINISTRATION** .

Inspection Before Preparation for Administration

Visually inspect the container. If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired. Check for minute leaks by squeezing bag firmly. If leaks are found, discard solution as sterility may be impaired. If supplemental medication is desired, follow directions below. Check solution for clarity and absence of foreign matter. If solution is not clear or contains foreign matter, discard the solution.

Preparation for Administration

1. Suspend container from eyelet support.
2. Remove plastic protector from outlet port at bottom of container.
 - Grip the small wing on the neck of the port with one hand.
 - Grip the large wing on the cap with the other hand and twist.
 - The cap will pop off.
3. Attach administration set. Refer to complete directions accompanying set.

To Add Medication

Additives may be incompatible.

To add medication before solution administration

1. Remove additive port closure: hold container below additive port and grasp cap between thumb and forefinger then flip cap upward. Swab exposed additive port using aseptic technique.
2. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
3. Mix solution and medication thoroughly. For high density medication such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.

To add medication during solution administration

1. Close clamp on the set.
2. Prepare medication site.
3. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
4. Remove container from IV pole and/or turn to an upright position.
5. Evacuate both ports by squeezing them while container is in the upright position.
6. Mix solution and medication thoroughly.
7. Return container to in-use position and continue administration.

Rev 02-09-21
Manufactured by:
Nephron Pharmaceuticals Corporation (Company Logo)
Nephron Pharmaceuticals Corporation
4500 12th Street Extension
West Columbia, SC 29172
1-844-224-2225

PACKAGE LABELING - PRINCIPLE DISPLAY PANEL

Principal Display - 50 mL Container

NDC 0487-4301-05

NDC: 0487-4301-05

0.9% SODIUM CHLORIDE INJECTION, USP

50 mL Single-Dose Container

Each 50 mL contains: 450 mg Sodium Chloride USP, pH 5.6 (4.5 to 7.0), mEq/50 mL Sodium 8 Chloride 8, Osmolarity 0.308 mOsmol/mL (calc).
STERILE NONPYROGENIC. FOR INTRAVENOUS USE. Read Package Insert for Full Information. Additives may be incompatible. Dosage Intravenously as directed by a physician. Cautions: Do not use unless solution is clear.

R_x Only

IC2091 Rev 04-15-19



Nephron Pharmaceuticals Corporation
4500 12th St. Extension
W. Columbia, SC 29172
1-844-224-2225

EXP: MMM YYYY

LOT: XXXXXX



Principal Display - 50 mL Case

NDC 0487-4301-05



Nephron Pharmaceuticals Corporation
 4500 12th Street Extension
 West Columbia, SC 29172
 1-844-224-2225

0000



0.9% Sodium Chloride Injection, USP
50 mL Single-Dose Container
 Read Package Insert for Full Information
 Store at 25°C/77°F. Avoid excessive heat.
 QTY 64 x 50 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-05



(01)0487430105(21)012345678901

Principal Display - 100 mL Container

NDC 0487-4301-10



Nephron Pharmaceuticals Corporation
 4500 12th Street Extension
 West Columbia, SC 29172
 1-844-224-2225

0000



0.9% Sodium Chloride Injection, USP
50 mL Single-Dose Container
 Read Package Insert for Full Information
 Store at 25°C/77°F. Avoid excessive heat.
 QTY 64 x 50 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-05



(01)0487430105(21)012345678901

NDC: 0487-4301-10

0.9% SODIUM CHLORIDE INJECTION, USP

100 mL Single-Dose Container

Each 100 mL contains: 900 mg Sodium Chloride USP, pH 5.6 (4.5 to 7.0), mEq/100 mL Sodium 15 Chloride 15, Osmolarity 308 mOsmol/L (calc).
STERILE NONPYROGENIC. FOR INTRAVENOUS USE. Read Package Insert for Full Information. Additives may be incompatible. Dosage Intravenously as directed by a physician. Cautions: Must not be used in series connections. Do not use unless solution is clear.



Nephron Pharmaceuticals Corporation
4500 12th St. Extension
W. Columbia, SC 29172
1-844-224-2225

Rx Only

IC2092 Rev 04-12-19

EXP: MMM YYYY
LOT: XXXXXX



Principal Display - 100 mL Case

NDC 0487-4301-10



Nephron Pharmaceuticals Corporation
4500 12th Street Extension
West Columbia, SC 29172
1-844-224-2225

0000



0.9% Sodium Chloride Injection, USP
100 mL Single-Dose Container
Read Package Insert for Full Information
Store at 25°C/77°F. Avoid excessive heat.
QTY 64 x 100 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-10



(01)0487430110(21)012345678901



Nephron Pharmaceuticals Corporation
4500 12th Street Extension
West Columbia, SC 29172
1-844-224-2225

0000



0.9% Sodium Chloride Injection, USP
100 mL Single-Dose Container
Read Package Insert for Full Information
Store at 25°C/77°F. Avoid excessive heat.
QTY 64 x 100 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-10



(01)0487430110(21)012345678901

Principal Display - 250 mL Container

NDC 0487-4301-25

0.9%

NDC: 0487-4301-25

SODIUM CHLORIDE INJECTION, USP

250 mL

 Single-Dose Container

Each 100 mL contains 900 mg Sodium Chloride USP, pH 5.6 (4.5 to 7.0), mEq/L Sodium 154 Chloride 154, Osmolarity 308 mOsmol/L (calc).

STERILE NONPYROGENIC. FOR INTRAVENOUS USE. Additives may be incompatible. Consult with pharmacist if available. When introducing additives use aseptic technique. Mix thoroughly. Do not store. Dosage intravenously as directed by a physician. Read Package Insert for Full Information. Cautions: Squeeze and inspect bag which maintains product sterility. Discard if leaks are found. Must not be used in series connections. Do not use unless solution is clear. Store at room temperature (25°C/77°F). Avoid excessive heat.

R_x Only

IC2093 Rev 04-15-19



Nephron Pharmaceuticals Corporation
4500 12th Street Extension
West Columbia, SC 29172
1-844-224-2225

EXP: MMM YYYY

LOT: XXXXXX



Principal Display - 250 mL Case

NDC 0487-4301-25



0.9% Sodium Chloride Injection, USP
250 mL Single-Dose Container
Read Package Insert for Full Information
Store at 25°C/77°F. Avoid excessive heat.
QTY 35 x 250 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-25



(01)0487430125(21)012345678901



0.9% Sodium Chloride Injection, USP
250 mL Single-Dose Container
Read Package Insert for Full Information
Store at 25°C/77°F. Avoid excessive heat.
QTY 35 x 250 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-25



(01)0487430125(21)012345678901

Principal Display - 500 mL Container

NDC 0487-4301-50

0.9%

NDC: 0487-4301-50

SODIUM CHLORIDE INJECTION, USP

500 mL Single-Dose Container

Each 100 mL contains: 900 mg Sodium Chloride USP, pH 5.6 (4.5 to 7.0), mEq/L Sodium 154, Chloride 154, Osmolarity 308 mOsmol/L (calc).

STERILE NONPYROGENIC. FOR INTRAVENOUS USE. Additives may be incompatible. Consult with pharmacist if available. When introducing additives use aseptic technique. Mix thoroughly. Do not store. Dosage intravenously as directed by a physician. Read Package Insert for Full Information. Cautions: Squeeze and inspect bag which maintains product sterility. Discard if leaks are found. Must not be used in series connections. Do not use unless solution is clear. Store at room temperature (25°C/77°F). Avoid excessive heat.

Rx Only

IC2094 Rev 04-15-19



Nephron Pharmaceuticals Corporation
4500 12th Street Extension
West Columbia, SC 29172
1-844-224-2225

EXP: MMM YYYY

LOT: XXXXXX



Principal Display - 500 mL Case

NDC 0487-4301-50



Nephron Pharmaceuticals Corporation
4500 12th Street Extension
West Columbia, SC 29172
1-844-224-2225

0000



0.9% Sodium Chloride Injection, USP
500 mL Single-Dose Container
Read Package Insert for Full Information
Store at 25°C/77°F. Avoid excessive heat.
QTY 35 x 500 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-50



(01)0487430150(21)012345678901



Nephron Pharmaceuticals Corporation
4500 12th Street Extension
West Columbia, SC 29172
1-844-224-2225

0000



0.9% Sodium Chloride Injection, USP
500 mL Single-Dose Container
Read Package Insert for Full Information
Store at 25°C/77°F. Avoid excessive heat.
QTY 35 x 500 mL

EXP: MMM YYYY LOT: A1234B



(17)000000(10)XXXXXX(37)00

NDC: 0487-4301-50



(01)0487430150(21)012345678901

SODIUM CHLORIDE

sodium chloride injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0487-4301
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	9 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0487-4301-25	35 in 1 CASE	06/04/2020	
1		250 mL in 1 BAG; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
2	NDC:0487-4301-10	64 in 1 CASE	06/04/2020	
		100 mL in 1 BAG; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

2		100 mL in 1 BAG; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
3	NDC:0487-4301-05	64 in 1 CASE	06/04/2020	
3		50 mL in 1 BAG; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
4	NDC:0487-4301-50	18 in 1 CASE	06/04/2020	
4		500 mL in 1 BAG; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211968	06/04/2020	

Labeler - Nephron Pharmaceuticals Corporation (079160190)

Establishment

Name	Address	ID/FEI	Business Operations
Nephron SC, LLC		079160190	manufacture(0487-4301) , analysis(0487-4301)

Revised: 1/2026

Nephron Pharmaceuticals Corporation