

**IBUPROFEN- ibuprofen tablet**  
**RETAIL BUSINESS SERVICES,LLC.**

-----  
**Ibuprofen 200 mg Tablets, USP**  
**Pain Reliever/Fever Reducer (NSAID)\***  
**See New Warnings**

**Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**Allergy alert:**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

### **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack and stroke warning:**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition

- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of the reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- do not take more than directed
- the smallest effective dose should be used

Adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

### **Other information**

- store between 20-25°C (68-77°F). [See USP Controlled Temperature]
- avoid excessive heat above 40°C (104°F)

### **Inactive ingredients**

colloidal silicon dioxide, ferric oxide red, maize starch, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide.

### **Questions or comments?**

call **1-877-770-3183** weekdays 9:00 AM to 4:30PM EST.

### **Principal Display Panel**

code # :		vi
size :	1+3/4 X 1+3/4 X 3+3/8	da
ref # :	PP180604B	
material :	.016 SBS	

COATING FREE AREA

R-00  
551000002  
300000524

3 4152031045 4

1219

S-2577

For patient updates or concerns, contact us at 1-877-848-8448. © 2007 Abbott Laboratories. All rights reserved. Daily peroxide or per meter test.

This product is not manufactured or distributed by Wyeth Consumer Healthcare, distributor of Advil®.

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
- you have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • big swelling • pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

**Drug Facts (continued)**

- any new symptoms appear
- pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older	take 1 tablet every 4 to 6 hours while symptoms persist
	if pain or fever does not respond to 1 tablet, 2 tablets may be used
	do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

**Other information**

- store between 20-25°C (68-77°F). [See USP Controlled Room Temperature].
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients** colloidal silicon dioxide, iron oxide red, maize starch, polyethylene glycol, povidone I-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

**Important: Read all the product information before using. Keep this box for important information.**

**DO NOT USE IF FOIL INNER SEAL IS BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet) Purpose**

Ibuprofen 200 mg Pain reliever/fever reducer (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** ibuprofen may cause severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock • skin redness • rash • blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

COATING FREE AREA

300000524  
551000002  
R-00

Actual Size



COATING FREE AREA

**Drug Facts (continued)**

**Questions or comments?**  
call 1-877-770-3183 weekdays 9:00 AM to 4:30 PM EST

Lot

Exp.

COATING FREE AREA

Principal Display Panel







Principal Display Panel



ibuprofen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72476-365
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

### Product Characteristics

<b>Color</b>	brown (Reddish Brown color Tablets)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	G2
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-365-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	07/26/2020	
2	NDC:72476-365-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/26/2020	
3	NDC:72476-365-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	07/26/2020	
4	NDC:72476-365-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	07/26/2020	
5	NDC:72476-365-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/26/2020	
6	NDC:72476-365-99	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/26/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	07/26/2020	

**Labeler** - RETAIL BUSINESS SERVICES,LLC. (967989935)

Revised: 12/2025

RETAIL BUSINESS SERVICES,LLC.