NYSTATIN- nystatin tablet, coated Bryant Ranch Prepack

Nystatin Tablets, USP (Oral) Rx only

DESCRIPTION

Nystatin is an antimycotic polyene antibiotic obtained from *Streptomyces noursei* . Its structural formula:

$$\begin{array}{c} HO \\ \\ HOOC \\ \\ \\ CH_3 \\ \\ CH_3$$

 $C_{47}H_{75}NO_{17}$

M.W. 926.13

Nystatin tablets are for oral administration and contain 500,000 units of nystatin per tablet.

Nystatin tablets contain the inactive ingredients: corn starch, confectioner sugar, dibasic calcium phosphate, FD&C yellow #6, FD&C red #40, FD&C blue # 2, hydroxypropyl cellulose, hypromellose, microcrystalline cellulose, magnesium stearate, polyethylene glycol, polysorbate 80, talc and titanium dioxide.

CLINICAL PHARMACOLOGY

Pharmacokinetics

Gastrointestinal absorption of nystatin is insignificant. Most orally administered nystatin is passed unchanged in the stool. In patients with renal insufficiency receiving oral therapy with conventional dosage forms, significant plasma concentrations of nystatin may occasionally occur.

Microbiology

Nystatin is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi. *Candida albicans* demonstrates no significant resistance to nystatin *in vitro* on repeated subculture in increasing levels of nystatin; other *Candida* species become quite resistant. Generally, resistance does not develop *in vivo*. Nystatin acts by binding to sterols in the cell membrane of susceptible *Candida* species with a resultant change in membrane permeability allowing leakage of intracellular components. Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

INDICATIONS AND USAGE

Nystatin Tablets are intended for the treatment of non-esophageal mucus membrane gastrointestinal candidiasis.

CONTRAINDICATIONS

Nystatin tablets are contraindicated in patients with a history of hypersensitivity to any of their components.

PRECAUTIONS

General

This medication is not to be used for the treatment of systemic mycoses. Discontinue treatment if sensitization or irritation is reported during use.

Carcinogenesis, Mutagenesis, Impairment of Fertility

No long-term animal studies have been performed to evaluate carcinogenic potential. There also have been no studies to determine mutagenicity or whether this medication affects fertility in males or females.

Pregnancy

Teratogenic Effects

Category C

Animal reproduction studies have not been conducted with nystatin. It is also not known whether nystatin can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Nystatin should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether nystatin is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when nystatin is administered to a nursing woman.

ADVERSE REACTIONS

Nystatin is well tolerated even with prolonged therapy. Oral irritation and sensitization have been reported. (See **PRECAUTIONS: General.**)

Gastrointestinal

Diarrhea (including one case of bloody diarrhea), nausea, vomiting, gastrointestinal upset/disturbances.

Dermatologic

Rash, including urticaria has been reported rarely. Stevens-Johnson syndrome has been reported very rarely.

Other

Tachycardia, bronchospasm, facial swelling, and nonspecific myalgia have also been rarely reported.

To report SUSPECTED ADVERSE REACTIONS, contact Avet Pharmaceuticals Inc. at 1-866-901-DRUG (3784) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Oral doses of nystatin in excess of five million units daily have caused nausea and gastrointestinal upset. There have been no reports of serious toxic effects of superinfections (see **CLINICAL PHARMACOLOGY**, **Pharmacokinetics**).

DOSAGE AND ADMINISTRATION

The usual therapeutic dosage is one to two tablets (500,000 to 1,000,000 units nystatin) three times daily. Treatment should generally be continued for at least 48 hours after clinical cure to prevent relapse.

HOW SUPPLIED

Nystatin Tablets USP, 500,000 Units are round brown, film-coated tablets debossed "HP51" on one side and plain on the other side are packaged in:

- NDC: 63629-8350-1: 28 Tablets in a BOTTLE
- NDC: 63629-8350-2: 56 Tablets in a BOTTLE
- NDC: 63629-8350-3: 30 Tablets in a BOTTLE
- NDC: 63629-8350-4: 120 Tablets in a BOTTLE

Store at 20° to 25°C (68° to 77°F). [See USP Controlled Room Temperature].

Repackaged/Relabeled by: Bryant Ranch Prepack, Inc. Burbank, CA 91504

Nystatin Tablets USP, 500,000 Units



Each tablet contains: Nystatin USP 500,000 Units.

Usual Adult Dosage: Scan Package Insert QR Code for prescribing literature.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container as defined in the USP.

Keep this and all drugs out of the reach of children.

NDC 63629-8350-1

Nystatin Tablets, USP (oral)

500,000 units

BRP

Repackaged by:

Bryant Ranch Prepack, Inc. Burbank, CA 91504 USA Rx only

28 Tablets
Manufactured by:
Strides Pharma Science





NYSTATIN

nystatin tablet, coated

Droduc	t Inform	ation
Produc	сиптоги	iation

Product Type

HUMAN PRESCRIPTION DRUG

HUMAN PRESCRIPTION (Source)

NDC:63629-8350(NDC:23155-051)

Route of Administration ORAL

Active Ingredient/Active Moiety

POLYETHYLENE GLYCOL 8000 (UNII: Q662QK8M3B)

Ingredient Name
Basis of Strength
NYSTATIN (UNII: BDF101C72E) (NYSTATIN - UNII:BDF101C72E)
NYSTATIN 500000 [USP'U]

Inactive Ingredients

Ingredient Name
Strength
STARCH, CORN (UNII: 08232NY3SJ)
SUCROSE (UNII: C151H8M554)
HYDROXYPROPYL CELLULOSE (90000 WAMW) (UNII: UKE75GEA7F)
ISOPROPYL ALCOHOL (UNII: ND2M416302)
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
TALC (UNII: 7SEV7J4R1U)
MAGNESIUM STEARATE (UNII: 70097M6I30)
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ 8WG20P6)
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	

Product Characteristics			
Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	HP;51
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63629- 8350-1	28 in 1 BOTTLE; Type 0: Not a Combination Product	04/29/2022	
2	NDC:63629- 8350-2	56 in 1 BOTTLE; Type 0: Not a Combination Product	09/02/2020	
3	NDC:63629- 8350-3	30 in 1 BOTTLE; Type 0: Not a Combination Product	09/02/2020	
4	NDC:63629- 8350-4	120 in 1 BOTTLE; Type 0: Not a Combination Product	06/07/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA062474	10/31/2011	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment			
Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(63629-8350), RELABEL(63629-8350)

Revised: 8/2025 Bryant Ranch Prepack