

NIGHTTIME COLD AND FLU RELIEF- nighttime cold and flu relief capsule

Grocery Outlet

Nighttime Cold and Flu Relief

Active ingredient(s)

Acetaminophen 325mg

Dextromethorphan HBr 15mg

Doxylamine succinate 6.25mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Use(s)

temporarily relieves common cold and flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever,

headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- to make a child sleep

Ask a doctor before use if you

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drink
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor before use if

- pain gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition

If pregnant or breastfeeding,

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children,

even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 4 doses per 24 hours
- **Adults and children 12 years and over:** 2 softgels with water every 6 hours
- **Children 4 to under 12 years of age:** ask a doctor
- **Children under 4 years of age:** do not use

Other information

store at room temperature 20-25 °C (68-77°F)

do not use if blister unit is torn or open

Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, white edible ink

Questions/Comments

Call **1-877-290-4008**

Principal Display Panel

Drug Facts		Drug Facts (continued)	
Active ingredients (in each softgel)	Purposes	Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin	
Acetaminophen 325mg.....	Pain reliever/fever reducer	When using this product ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drink ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness	
Dextromethorphan HBr 15mg.....	Cough suppressant	Stop use and ask a doctor if ■ pain gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
Doxylamine succinate 6.25mg.....	Antihistamine	If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.	
Uses temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing		Directions ■ take only as directed ■ do not exceed 4 doses per 24 hours	
Warnings		Adults and children 12 years and over 2 softgels with water every 6 hours	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4 doses in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product		Children 4 to under 12 years of age ask a doctor	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.		Children under 4 years of age do not use	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		Other information ■ store at room temperature 20-25 °C (68-77° F)	
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drugs. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product ■ to make a child sleep		Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, white edible ink	
Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland		Questions or comments? Call 1-877-290-4008	

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Vicks® NyQuil™ Cold & Flu Relief LiquiCaps®

DISTRIBUTED BY: Grocery Outlet, Inc. | Emeryville, CA 94608
www.groceryoutlet.com
PRODUCT OF CHINA.
LB2428 RT125

100% MONEY BACK GUARANTEE

GO Nighttime Cold and Flu Relief

ACETAMINOPHEN, 325 mg (Pain Reliever / Fever Reducer)
DEXTROMETHORPHAN HBr, 15 mg (Cough Suppressant)
DOXYLAMINE SUCCLNATE, 6.25 mg (Antihistamine)

GO Nighttime Cold and Flu Relief

Relief of:

- Aches
- Sore Throat
- Runny Nose
- Fever
- Sneezing
- Cough

Actual Size

24 SOFTGELS

DOXYLAMINE SUCCLNATE, 6.25 mg (Antihistamine)
DEXTROMETHORPHAN HBr, 15 mg (Cough Suppressant)
ACETAMINOPHEN, 325 mg (Pain Reliever / Fever Reducer)

GO Nighttime Cold and Flu Relief

ACETAMINOPHEN, 325 mg (Pain Reliever / Fever Reducer)
DEXTROMETHORPHAN HBr, 10 mg (Cough Suppressant)
DOXYLAMINE SUCCLNATE, 6.25 mg (Antihistamine)

Lat: _____
Exp: _____

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

DO NOT USE IF BUBBLER UNITS IS TORN OR OPEN

8 10159 83561 9

Blister

NIGHTTIME COLD AND FLU RELIEF

nighttime cold and flu relief capsule

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:85828-418
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
WATER (UNII: 059QF0KO0R)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Product Characteristics

Color	green (Clear)	Score	no score
Shape	CAPSULE (Oblong)	Size	20mm
Flavor		Imprint Code	PC10
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:85828-418-93	2 in 1 CARTON	03/25/2026	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/25/2026	

