

**MUCUS RELIEF SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH-**  
**dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid**  
**Rite Aid Corporation**

-----

**Drug Facts**

**Active ingredients (in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCL 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal Decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

**Warnings**

**Do not use**

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product,  
do not use more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with a fever
- cough comes back, or occurs with fever, rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in a 24-hour period
- measure only with dosing cup provided Do not use any other dosing device.
- keep dosing cup with product
- mL=milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

**Other information**

- **each 20 mL contains:** sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum

**Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

Compare to the active ingredients in Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough\*

### **MAXIMUM STRENGTH**

**MUCUS** RELIEF

**SEVERE COUGH & CONGESTION**

**MULTI-SYMPTOM RELIEF**

COUGH SUPPRESSANT

EXPECTORANT

NASAL DECONGESTANT

DEXTROMETHORPHAN HBr 20 mg

GUAIFENESIN 400 mg

PHENYLEPHRINE HCL 10 MG

Controls cough

Relieves nasal & chest congestion

Thins & loosens mucus

For ages 12 & over

FL OZ (mL)

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

**DISTRIBUTED BY:**

RITE AID, 30 HUNTER LANE

CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

**Package Label**



### Drug Facts (continued)

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product, do not use more than directed.**

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.



DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 [www.riteaid.com](http://www.riteaid.com)

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

PLD-A283D LB007073

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

### Drug Facts

#### Active ingredients (in each 20 mL)

|                            |                    |
|----------------------------|--------------------|
| Dextromethorphan HBr 20 mg | .....              |
| .....                      | Cough suppressant  |
| Guaifenesin 400 mg         | .....              |
| .....                      | Expectorant        |
| Phenylephrine HCl 10 mg    | .....              |
| .....                      | Nasal decongestant |

#### Purposes

PEEL CORNER FOR MORE DRUG FACTS

### Drug Facts (continued)

#### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

#### Other information

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

#### Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### Drug Facts (continued)

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

### Warnings

#### Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

### Drug Facts (continued)

#### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PEEL CORNER FOR MORE DRUG FACTS

**RITE AID Maximum Strength Mucus Relief Severe Cough & Congestion**

## MUCUS RELIEF SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

| Product Information                                                                      |                  |                                                                |                                  |                    |
|------------------------------------------------------------------------------------------|------------------|----------------------------------------------------------------|----------------------------------|--------------------|
| Product Type                                                                             |                  | HUMAN OTC DRUG                                                 | Item Code (Source)               | NDC:11822-3722     |
| Route of Administration                                                                  |                  | ORAL                                                           |                                  |                    |
|                                                                                          |                  |                                                                |                                  |                    |
| Active Ingredient/Active Moiety                                                          |                  |                                                                |                                  |                    |
| Ingredient Name                                                                          |                  |                                                                | Basis of Strength                | Strength           |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)<br>(DEXTROMETHORPHAN - UNII:7355X3ROTS) |                  |                                                                | DEXTROMETHORPHAN<br>HYDROBROMIDE | 20 mg<br>in 20 mL  |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)                           |                  |                                                                | GUAIFENESIN                      | 400 mg<br>in 20 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)         |                  |                                                                | PHENYLEPHRINE<br>HYDROCHLORIDE   | 10 mg<br>in 20 mL  |
|                                                                                          |                  |                                                                |                                  |                    |
| Inactive Ingredients                                                                     |                  |                                                                |                                  |                    |
| Ingredient Name                                                                          |                  |                                                                |                                  | Strength           |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)                                                 |                  |                                                                |                                  |                    |
| EDETATE DISODIUM (UNII: 7FLD91C86K)                                                      |                  |                                                                |                                  |                    |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD)                                                       |                  |                                                                |                                  |                    |
| FD&C RED NO. 40 (UNII: WZB9127XOA)                                                       |                  |                                                                |                                  |                    |
| GLYCERIN (UNII: PDC6A3C0OX)                                                              |                  |                                                                |                                  |                    |
| PROPYL GALLATE (UNII: 8D4SNN7V92)                                                        |                  |                                                                |                                  |                    |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3)                                                      |                  |                                                                |                                  |                    |
| WATER (UNII: 059QF0KO0R)                                                                 |                  |                                                                |                                  |                    |
| SODIUM BENZOATE (UNII: OJ245FE5EU)                                                       |                  |                                                                |                                  |                    |
| TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)                                           |                  |                                                                |                                  |                    |
| SORBITOL (UNII: 506T60A25R)                                                              |                  |                                                                |                                  |                    |
| SUCRALOSE (UNII: 96K6UQ3ZD4)                                                             |                  |                                                                |                                  |                    |
| XANTHAN GUM (UNII: TTV12P4NEE)                                                           |                  |                                                                |                                  |                    |
|                                                                                          |                  |                                                                |                                  |                    |
| Packaging                                                                                |                  |                                                                |                                  |                    |
| #                                                                                        | Item Code        | Package Description                                            | Marketing Start Date             | Marketing End Date |
| 1                                                                                        | NDC:11822-3722-2 | 177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 09/17/2021                       | 02/01/2027         |
|                                                                                          |                  |                                                                |                                  |                    |
| Marketing Information                                                                    |                  |                                                                |                                  |                    |
| Marketing Category                                                                       |                  | Application Number or Monograph Citation                       | Marketing Start Date             | Marketing End Date |
| OTC Monograph Drug                                                                       |                  | M012                                                           | 09/17/2021                       | 02/01/2027         |

**Labeler** - Rite Aid Corporation (014578892)