

**PHENAGIL- chlorpheniramine maleate, dextromethorphan hbr and phenylephrine hcl tablet**  
**Gil Pharmaceutical Corp.**

-----  
**PHENAGIL® CH**

**ANTI-HISTAMINE, NASAL DECONGESTANT AND ANTITUSSIVE**

**DYE FREE**

**PRESERVATIVE FREE**

**SODIUM FREE**

**SUGAR FREE**

***Drug Facts***

***Active ingredients***

(in each tablet)

Chlorpheniramine maleate 3.5 mg

Dextromethorphan HBr 29 mg

Phenylephrine HCl 10 mg

***Purposes***

Antihistamine

Cough suppressant

Nasal decongestant

***Uses***

Temporarily relieves:

- cough due to minor throat and bronchial irritation associated with a cold or inhaled irritants.
- nasal congestion due to the common cold, hay fever or other upper respiratory allergies.
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
  - runny nose.
  - sneezing.
  - itchy, watery eyes.
  - itching of the nose or throat.
  - temporarily restores freer breathing through the nose.

## **Warnings**

### **Do not use**

- to sedate a child or to make a child sleepy.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- heart disease.
- high blood pressure.
- thyroid disease.
- diabetes.
- trouble urinating due to an enlarged prostate gland.
- glaucoma.
- cough that occurs with too much phlegm (mucus).
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema.

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers.
- taking any other oral nasal decongestant or stimulant.

### **When using this product**

- **do not use more than directed.**
- may cause drowsiness.
- avoid alcoholic beverages.
- alcohol, sedatives, and tranquilizers may increase drowsiness.
- be careful when driving a motor vehicle or operating machinery.
- excitability may occur, especially in children.

### **Stop use and ask a doctor if**

- you get nervous, dizzy, or sleepless.
- symptoms do not get better within 7 days or are accompanied by fever.
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions

- **Adults and Children 12 years and over.:** Take one tablet every 6-8 hours. Do not take more than 4 tablets in a 24 hour period.
- **Children 6 to 12 years of age:** Take 1/2 tablet every 6-8 hours. Do not take more than 2 tablets in a 24 hour period.
- **Children under 6 years:** Consult a doctor.

## Other information

- store at room temperature.
- do not use if safety seal is broken or missing.

## Inactive ingredients

Hydroxypropyl Methylcellulose, Lactose Monohydrate, Microcrystalline Cellulose, Magnesium Stearate and Polyethylene Glycol.

## Questions or comments?

Call **1-787-848-9114**, Monday–Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

## MANUFACTURED FOR:

GIL PHARMACEUTICAL CORP.  
PONCE, PUERTO RICO 00716

Label revised: 04/22

## PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

**PHENAGIL<sup>®</sup> CH - NDC 58552-337-01 - 100's Bottle Label.**

**PHENAGIL<sup>®</sup> CH**  
NDC 58552-337-01  
ANTIHISTAMINE, NASAL  
DECONGESTANT AND ANTITUSSIVE  
DYE FREE. PRESERVATIVE FREE.  
SODIUM FREE. SUGAR FREE.

**Each tablet contains:**  
Chlorpheniramine maleate 3.5 mg  
Dextromethorphan HBr 29 mg  
Phenylephrine HCl 10 mg

**100 Tablets**  
MANUFACTURED FOR:  
GIL PHARMACEUTICAL CORP.  
PONCE, PUERTO RICO 00716

**Drug Facts**

<b>Drug Facts</b> (continued)
<b>Ask a doctor before use if you have</b> • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
<b>Ask a doctor or pharmacist before use if you are</b> • taking sedatives or tranquilizers • taking any other oral nasal decongestant or stimulant
<b>When using this product</b> • <b>do not use more than directed</b> • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or

## Drug Facts

### Active ingredients .....Purposes

(in each tablet)

Chlorpheniramine maleate 3.5 mg .....Antihistamine  
Dextromethorphan HBr 29 mg .... Cough suppressant  
Phenylephrine HCl 10 mg ..... Nasal decongestant

### Uses

Temporarily relieves:

- cough due to minor throat and bronchial irritation associated with a cold or inhaled irritants
- nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - temporarily restores free breathing through the nose

### Warnings

#### Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

• Be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children

#### Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever • cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- **Adults and Children 12 years and over:** Take one tablet every 6-8 hours. Do not take more than 4 tablets in a 24 hour period.
- **Children 6 to 12 years of age:** Take 1/2 tablet every 6-8 hours. Do not take more than 2 tablets in a 24 hour period.
- **Children under 6 years:** Consult a doctor.

### Other information

- store at room temperature
- do not use if safety seal is broken or missing

### Inactive ingredients

Hydroxypropyl Methylcellulose, Lactose Monohydrate, Magnesium Stearate, Microcrystalline Cellulose and Polyethylene Glycol

### Questions or comments?

Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 04/22

PEEL HERE FOR MORE INFORMATION

## PHENAGIL

chlorpheniramine maleate, dextromethorphan hbr and phenylephrine hcl tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58552-337
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	3.5 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	29 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	29 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	3mm
<b>Flavor</b>		<b>Imprint Code</b>	323;Gil
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58552-337-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/26/2022	
2	NDC:58552-337-02	2 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/26/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/26/2022	

**Labeler** - Gil Pharmaceutical Corp. (176826592)

**Registrant** - Syntho Pharmaceuticals, Inc. (113616187)

### Establishment

Name	Address	ID/FEI	Business Operations
Syntho Pharmaceuticals, Inc.		088797407	manufacture(58552-337)

Revised: 10/2025

Gil Pharmaceutical Corp.