

**NIGHTTIME COLD AND FLU RELIEF- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid
Safeway, Inc.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours,
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center

(1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **When using other Day Time or Night Time products, carefully read each label to insure correct dosing**

Other information

- **each 30 mL contains;** potassium 5 mg
- **each 30 mL contains:** sodium 24 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acesulfame potassium, alcohol, citric acid, FD&C blue #1, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments

Call **1-888-723-3929 Monday-Friday 7AM-6PM PST**

Principal Display Panel

Compare to Vicks® NyQuil® Cold & Flu

Nighttime Cold & Flu Relief

Acetaminophen 650 mg-Pain Reliever/Fever Reducer

Dextromethorphan HBr 30 mg- Cough Suppressant

Doxylamine succinate 12.5 mg-Antihistamine

CHERRY FLAVOR

- Relieves headache, fever, sore throat, minor aches & pains
- Sneezing, runny nose, cough
- For ages 12 years & over
- Nighttime relief
- Alcohol 10%

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY

BETTER LIVING BRANDS LLC

P.O. BOX 99, PLEASANTON, CA 94566-0009

www.betterlivingbrandsLLC.com

Product Label

Drug Facts

Active ingredients (in each 30 mL) **Purposes**

Acetaminophen 650 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 30 mg.....Cough suppressant
 Doxylamine succinate 12.5 mg.....Antihistamine

Uses ■ temporarily relieves these common cold/flu symptoms
 ■ minor aches and pains ■ headache ■ sore throat
 ■ runny nose and sneezing ■ fever
 ■ cough due to minor throat and bronchial irritation

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks

← PEEL HERE **Drug Facts** (continued under label)



Compare to
 Vicks® NyQuil®
 Cold & Flu
 active ingredients*

NDC 21130-343-08

Nighttime Cold & Flu Relief

ACETAMINOPHEN 650 mg
 - Pain Reliever/Fever Reducer
DEXTROMETHORPHAN HBr 30 mg
 - Cough Suppressant
DOXYLAMINE SUCCINATE 12.5 mg
 - Antihistamine

CHERRY FLAVOR

- Relieves headaches, fever, sore throat, minor aches & pains
- Sneezing, runny nose, cough
- For ages 12 years & over
- Nighttime relief
- Alcohol 10%

8 FL OZ (237 mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.



DISTRIBUTED BY
 BETTER LIVING BRANDS LLC
 P.O. BOX 99, PLEASANTON, CA
 94566-0009
 1-888-723-3929
 www.betterlivingbrandsLLC.com



Scan here for more information

OUR PROMISE
 QUALITY & SATISFACTION
100% GUARANTEED
 OR YOUR MONEY BACK.



Drug Facts (continued)

after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
 ■ liver disease ■ glaucoma
 ■ cough that occurs with too much phlegm (mucus)
 ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
 ■ trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are
 ■ taking sedatives or tranquilizers
 ■ taking the blood thinning drug warfarin

When using this product
 ■ excitability may occur, especially in children
 ■ marked drowsiness may occur ■ avoid alcoholic drinks
 ■ be careful when driving a motor vehicle or operating machinery
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
 ■ pain or cough gets worse or lasts more than 7 days
 ■ new symptoms occur
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.

Drug Facts (continued)

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ do not take more than directed (see Overdose warning)
 ■ do not take more than 4 doses in any 24-hour period
 ■ measure only with dosing cup provided. Do not use any other dosing device. ■ mL = milliliter
 ■ keep dosing cup with product
 ■ adults and children 12 years and over: 30 mL every 6 hours
 ■ children under 12 years of age: do not use
 ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information
 ■ each 30 mL contains: potassium 5 mg
 ■ each 30 mL contains: sodium 24 mg
 ■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acesulfame potassium, alcohol, citric acid, FD&C blue #1, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?
 Call 1-888-723-3929 Monday-Friday 7AM-6PM PST

SIGNATURE CARE Nighttime Cold & Flu Relief

NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-343
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-343-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/30/2014	
2	NDC:21130-343-24	2 in 1 PACKAGE	06/30/2014	
2		355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:21130-343-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/30/2014	

Labeler - Safeway, Inc. (009137209)