

4X MEDICATED TOOTHACHE AND GUM CREAM- benzocaine cream Walgreens

5820627 WLG 4x Med Cream (80-15601)

Benzalkonium chloride 0.13% . . . Oral antiseptic

Benzocaine 20% . . . Oral pain reliever

Menthol 0.26% . . . Oral pain reliever

Zinc chloride 0.15% . . . Oral astringent

Use

for the temporary relief of pain due to toothaches * to help protect against infection of minor oral irritation

Methemoglobinemia warning: use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develop: * pale, gray, or blue colored skin (cyanosis) * headache * rapid heart rate * shortness of breath * dizziness or lightheadedness * fatigue or lack of energy

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

Do not use * more than directed * for more than 7 days unless told to do so by a dentist or doctor * for teething * in children under 2 years of age

Stop use and ask a doctor if * swelling, rash or fever develops * irritation, pain or redness persists or worsens * symptoms do not improve in 7 days * allergic reaction occurs

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

cut open tip of tube on score mark * adults and children 2 years of age and older: apply a small amount of product to the cavity and around the gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or doctor * children under 12 years of age should be supervised in the use of this product * children under 2 years of age: do not use

do not use if tip is cut prior to opening * this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist can be consulted * do not use continuously * this formula will stay in place for extended duration of relief * avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying

ammonium glycyrrhizate, flavor, PEG-8, PEG-75, sodium saccharin, sorbic acid, titanium dioxide

NDC 0363-0607-19

Walgreens



Compare to the active ingredients in Orajel™ 4X Medicated for Toothache & Gum Cream*

4x Medicated

Cream

- 20% benzocaine to relieve oral pain
- Antiseptic to kill harmful bacteria
- Astringent to help prevent infection
- Menthol to soothe

Long-lasting oral pain relief

Instant Toothache & Gum Relief

ORAL PAIN RELIEVER / ASTRINGENT / ANTISEPTIC

Wintergreen
flavor

NET WT 0.5 OZ (14 g)

TAMPER-EVIDENT: DO NOT USE IF TUBE TIP IS CUT PRIOR TO USE. RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION.

¹Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
²This product is not manufactured or distributed by Church & Dwight Co., Inc., owner of the trademark Orajel™.
 DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
 walgreens.com 1-800-925-4733 ©2023 Walgreen Co.

ITEM 683212 W00000-0000-0



2321627R2
5820627

W3REVI 124-F

Drug Facts

Active ingredients
 Benzocaine 20%
 Menthol 0.26%
 Zinc chloride 0.15%
 Oral pain reliever
 Oral pain reliever
 Oral antiseptic

Purpose
 • first aid to help prevent infection in minor oral irritation

Uses • for the temporary relief of pain due to minor irritation or injury of the mouth and gums

Warnings
Methemoglobinemia warning: use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

Do not use • more than directed • for more than 7 days unless told to do so by a dentist or doctor • for teething • in children under 2 years of age

Stop use and ask a doctor if • symptoms do not improve in 7 days • swelling, rash or fever develops • irritation, pain or redness persists or worsens

Keep out of reach of children. In case of overdose or allergic reaction, get medical help or contact a Poison Control Center right away.

Directions • cut open tip of tube on score mark • adults and children 2 years of age and older: apply to affected area up to 4 times daily or as directed by a dentist or doctor • children under 2 years of age: do not use

Other information • do not use if tip is cut prior to opening • avoid using toothpaste or drinking soft drinks or fruit juices for at least one hour after applying

Inactive ingredients ammonium glycyrrhizate, flavor, PEG-8, PEG-75, PEG-8, PEG-75, sodium saccharin, sorbic acid, titanium dioxide



pe code,
pe prior
move

4X MEDICATED TOOTHACHE AND GUM CREAM

benzocaine cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0607
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 g in 100 g
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
ZINC CHLORIDE (UNII: 86Q357L16B) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	0.15 g in 100 g
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A) (MENTHOL, UNSPECIFIED FORM - UNII:L7T10EIP3A)	MENTHOL, UNSPECIFIED FORM	0.26 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
SORBIC ACID (UNII: X045WJ989B)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white (Cloudy)	Score	
Shape		Size	
Flavor	WINTERGREEN	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0607-79	1 in 1 CARTON	06/02/2020	
1		9.4 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:0363-0607-19	1 in 1 CARTON	02/16/2024	
2		14 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

OTC Monograph Drug	M017	06/02/2020	
--------------------	------	------------	--

Labeler - Walgreens (008965063)

Registrant - Lornamead Products (119559733)

Establishment

Name	Address	ID/FEI	Business Operations
Lornamead Products		119559733	manufacture(0363-0607) , pack(0363-0607)

Revised: 11/2025

Walgreens