3X MEDICATED MOUTH SORE GEL- benzocaine gel
HEB

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

5820619 3x Med Gel

Active ingredients Purpose

Benzocaine 20% . . . . . .Oral pain reliever

Menthol 0.1% . . . . Oral pain reliever

Zinc chloride 0.15% . . . .Oral astringent

Uses

temporarily relieves pain caused by * canker sores * cold sores * fever blisters * minor irritation or injury of the mouth and gums

Methemoglobinemia warning: use of this product may cause methoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: * pale, gray, or blue colored skin (cyanosis) * headache * rapid heart rate * shortness of breath * dizziness or lightheadedness * fatigue or lack of energy

Allergy alert

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics

Do not use * more than directed * for more than 7 days unless told to do so by a dentist or doctor * for teething * in children under 2 years of age

Stop use and ask a doctor if * swelling, rash or fever develops * irritation, pain or redness persists or worsens * symptoms do not improve in 7 days * allergic reaction occurs

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Directions * cut open tip of tube on score mark * do not use if tip is cut prior to opening * adults and children 2 years of age and older: apply to affected area up to 4 times daily or as directed by a dentist or doctor. Children under 12 years of age should be supervised in the use of this product * children under 2 years of age: do not use

allantoin, carbomer, disodium EDTA, flavor, polyethylene glycol, polysorbate 60, propylene glycol, pvp, sodium saccharin, sorbic acid, stearyl alcohol, water

4.313

0.875

1.813

Compare to the active ingredients
in Orajel™ 3X Medicated
For All Mouth Sores Gel*

• 20% Benzocaine to
  relieve oral pain

• Menthol to soothe gums

• Astringent to help protect
  sores from irritation

H·E·B **3X** MEDICATED
MOUTH SORE RELIEF

FAST-ACTING GEL

ORAL PAIN RELIEVER
ASTRINGENT

H·E·B **3X** MEDICATED
MOUTH SORE RELIEF

Immediate pain relief of canker sores, cold sores,
cheek bites & pain from dentures or braces

NET WT. 0.42 OZ (11.9g)

0.469

FAST-ACTING GEL

ORAL PAIN RELIEVER
ASTRINGENT

H·E·B **3X** MEDICATED
MOUTH SORE RELIEF

Immediate pain relief of canker sores, cold sores,
cheek bites & pain from dentures or braces

NET WT. 0.42 OZ (11.9g)

100%
GUARANTEE
promise

H·E·B
If you aren't completely pleased
with this product, we'll be happy to
replace it or refund your money.
You have our word on it.

* This product is not manufactured or distributed by Church & Dwight Co., Inc.,
owner of the registered trademark Orajel™.

MADE WITH PRIDE AND CARE FOR H-E-B®
SAN ANTONIO, TX 78204

0 41220 33151 2

7.594

0.937

Retain carton for complete product information

Questions or Comments? Call 1-877-777-0743

Inactive ingredients: alcohol, carbomer, disodium EDTA, flavor, polyethylene glycol, polysorbate 60, propylene glycol, PVP, sodium saccharin, sorbic acid, stearyl alcohol, water.

Directions: • cut open tip of tube on score mark • do not use if top is cut prior to opening • **adults and children 2 years of age and older:** apply to affected area up to 4 times daily or as directed by a dentist or doctor. Children under 12 years of age should be supervised in the use of this product • **children under 2 years of age:** do not use

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Stop use and ask a doctor if • swelling, rash or fever develops • irritation, pain or redness persists or worsens • symptoms do not improve in 7 days • allergic reaction occurs

Do not use • more than directed • for more than 7 days unless told to do so by a dentist or doctor • for teething • in children under 2 years of age

Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics • fatigue or lack of energy

Methemoglobinemia warning: use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops • pale, gray or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness

Warnings

Uses temporarily relieves pain caused by • canker sores • cold sores • fever blisters • minor irritation or injury of the mouth and gums

Zinc chloride 0.18%..Oral astringent

Menthol 0.1%...

Benzocaine 20%..Oral pain reliever

Active Ingredients   **Purpose**

Drug Facts

4.25

2.781

6.187

3X MEDICATED MOUTH SORE GEL
benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:37808-691
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC CHLORIDE (UNII: 86Q357L16B) (ZINC CATION - UNII:13S1S8SF37)	ZINC CATION	0.15 g in 100 g
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A) (MENTHOL, UNSPECIFIED FORM - UNII:L7T10EIP3A)	MENTHOL, UNSPECIFIED FORM	0.1 g in 100 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
CARBOMER HOMOPOLYMER TYPE B (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: HHT01ZNK31)	
SORBIC ACID (UNII: X045WJ989B)	
ALLANTOIN (UNII: 344S277G0Z)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
METHYL SALICYLATE (UNII: LAV5U5022Y)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	yellow (Clear to yellow tint)	**Score**	
Shape		**Size**	
Flavor	WINTERGREEN	**Imprint Code**	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-691-19	1 in 1 CARTON	06/02/2020	
1		9.4 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	06/02/2020	

Labeler - HEB (007924756)

Registrant - Lornamead (126440440)

Establishment

Name	Address	ID/FEI	Business Operations
Lornamead		080046418	manufacture(37808-691)

Revised: 1/2023 HEB