

IBUPROFEN- ibuprofen tablet, coated
RITE AID CORPORATION

601R RITE AID 11822 7892 IBUPROFEN 200 MG TABLETS

DRUG FACTS

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonstreoidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pains due to :

- backache
- headache
- menstrual cramps
- minor pain of arthritis
- muscular aches
- the common cold
- toothache

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin

under a doctor's care for any serious condition

taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problem or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of the body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

**do not take more than directed
the smallest effective dose should be used**

adults and children 12 years and over:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor.

children under 12 years:

- ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? Call 1-877-290-4008

NDC 11822-7892-7

PAIN RELIEF IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)



200
TABLETS

Read all product information before using. Keep the box for important information.

Uses ■ temporarily relieves minor aches and pains due to: backache, headache, menstrual cramps, minor pain of arthritis, muscular aches, the common cold, toothache ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: asthma (wheezing), hives, rash, skin redness, and swelling. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used

adults and children 12 years and over: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years: ■ ask a doctor

Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY: RITE AID SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money. 601R 08211

www.riteaid.com Code No. G00RUGS515 JUL462

CAMP HILL, PA 17011

MADE IN INDIA

WARNING: IBUPROFEN MAY CAUSE SEVERE STOMACH BLEEDING. THE CHANCE IS HIGHER IF YOU ARE AGE 60 OR OLDER, HAVE HAD STOMACH ULCERS OR BLEEDING PROBLEMS, TAKE A BLOOD-THINNING (ANTICOAGULANT) OR STEROID DRUG, TAKE OTHER DRUGS CONTAINING PRESCRIPTION OR NONPRESCRIPTION NSAIDS (ASPIRIN, IBUPROFEN, NAPROXEN, OR OTHERS), HAVE 3 OR MORE ALCOHOLIC DRINKS EVERY DAY WHILE USING THIS PRODUCT, TAKE MORE OR FOR A LONGER TIME THAN DIRECTED.

Drug Facts (continued)
directions before use ■ store at 20-25°C (68-77°F)
Inactive ingredients: croscarmellose sodium, iron oxide red, hydroxypropyl methylcellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide
Questions or comments? Call 1-877-290-4008

Drug Facts (continued)
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery
Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug
When using this product ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

Drug Facts (continued)
■ you have symptoms of heart problem or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of the body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).
Directions ■ do not take more than directed ■ the smallest effective dose should be used
adults and children 12 years and over: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years: ■ ask a doctor
Other information ■ read all warnings and

Compare to the active ingredients in Advil® Tablets*

PAIN RELIEF
IBUPROFEN
IBUPROFEN TABLETS USP, 200 mg
PAIN RELIEVER/FEVER REDUCER (NSAID)

ACTUAL SIZE
50 TABLETS

NDC 11822-7892-2

Coating/Print Omit Area
Exp Date: _____
Lot No.: _____
4 11822 7892 0

THIS PRODUCT IS NOT MANUFACTURED OR DISTRIBUTED BY RITE AID. IT IS THE PROPERTY OF THE REGISTERED PROPRIETOR. PLEASE CONSUME HEALTHY AND REGISTERED PRODUCTS WITH CARE. © 2011 RITE AID. ALL RIGHTS RESERVED. PLEASE RECYCLE YOUR MONEY.

Coating Omit

E826R1 Coating/Print Omit Area

Coating Omit

NDC 11822-7892-5

PAIN RELIEF IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)



100
TABLETS

Read all product information before using. Keep the box for important information.

Uses ■ temporarily relieves minor aches and pains due to: ■ headache, ■ backache, ■ menstrual cramps, ■ minor pain of arthritis, ■ muscular aches, ■ the common cold, ■ toothache ■ temporarily reduces fever

Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ asthma (wheezing) ■ hives ■ rash ■ shock ■ skin redness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F) **Questions or comments?** Call 1-877-290-4008

DISTRIBUTED BY: RTE AD, SATISFACTION GUARANTEE, 30 HUNTER LANE, CAMP HILL, PA 17011 If you're not satisfied, we'll happily refund your money. 601R 0821
www.rtead.com MADE IN INDIA

Code No. - GO/DRUGS/515 HULL468

Lot No.:

Exp. Date:

READ AND KEEP CAREFUL FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each tablet): ibuprofen (NSAID) ■ pain reliever/fever reducer ■ nonsteroidal anti-inflammatory drug

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ backache ■ menstrual cramps ■ minor pain of arthritis ■ the common cold ■ temporary reduces fever

Warnings ■ Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ asthma (wheezing) ■ hives ■ rash ■ shock ■ skin redness. If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug ■ taking with food or milk if stomach upset occurs

Drug Facts (continued)

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ you have blood or black stools ■ you have symptoms of heart problem or stroke ■ chest pain ■ dizziness ■ fainting ■ lightheadedness ■ ringing in the ears ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ any new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Other information ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, zinc stearate, and others

Questions or comments? Call 1-877-290-4008

This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Tablets. 601R 0821
DISTRIBUTED BY: RTE AD, SATISFACTION GUARANTEE, 30 HUNTER LANE, CAMP HILL, PA 17011
www.rtead.com
MADE IN INDIA

NDC 11822-7892-7

PAIN RELIEF

IBUPROFEN

PAIN RELIEVER/FEVER REDUCER
(NSAID)



ACTUAL SIZE
200
TABLETS

Compare to the active ingredient in Advil® Tablets*

Varnish/Print Omit Area



Lot No.:
Exp. Date:

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Code: GO/DRUGS/515

18C2207

Varnish Omit



Varnish/Print Omit Area

E6011AR5

Coating/Print
Omit Area



This product is not manufactured or distributed by
Pfizer Consumer Healthcare, owner of the registered
trademark Advil® Tablets, 601R 0021
DISTRIBUTED BY: RITE RID, 30 HUNTER LANE,
GMP HILL, PA 17011 www.riteaid.com MADE IN INDIA
SATISFACTION GUARANTEE:
If you're not satisfied,
happily refund your money.

Code: 601R0021S15

Read all product information before using.
Keep the box for important information.

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen
200 mg (NSAID)* Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- Temporarily relieves minor aches and pains due to:
 - backache
 - headache
 - menstrual cramps
 - minor pain of arthritis
 - muscular aches
 - the common cold
 - toothache
- Temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- hives
- rash
- shock
- skin reddening
- facial swelling
- skin redness

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing

Drug Facts (continued)

- weakness in one part or side of the body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and over:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Drug Facts (continued)
Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide
Questions or comments?
Call 1-817-250-4000
DANGER: FINGER DO NOT USE THIS PRODUCT IF THE IMPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.



Coating
Omit

Compare to the active ingredient in Advil® Tablets*

PAIN RELIEF
IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg
PAIN RELIEVER/FEVER REDUCER
(NSAID)

ACTUAL SIZE
100
TABLETS



Coating/Print Omit Area

NDC 11822-7892-5



Coating
Omit

Compare to the active ingredient in Advil® Tablets†

NDC 11822-7892-9

PAIN RELIEF
IBUPROFEN
IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)

ACTUAL SIZE

500
TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE CUT, TORN, BROKEN OR MISSING IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS

DRUG FACTS (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use: If you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer. Right before or after heart surgery.

Ask a doctor before use if: You have problems or serious side effects from taking pain relievers or fever reducers. You have a history of stomach bleeding, vomiting, or diarrhea. You have high blood pressure, heart disease, new or worse, kidney disease, asthma, or had a stroke. You are taking a diuretic.

Ask a doctor or pharmacist before use if you are: Taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin. Under a doctor's care for any serious condition. Taking any other drug.

When using this product: Take with food or milk if stomach upset occurs.

Stop use and ask a doctor if: You experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the genital area
- any new symptoms appear

If pregnant or breast-feeding: Ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions: Do not take more than directed. The smallest effective dose should be used.

adults and children 12 years and older:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information: Read all warnings and directions before use. Store at 20-25°C (68-77°F).

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? Call 1-877-290-4008

NDC 11822-7892-2

PAIN RELIEF
IBUPROFEN
IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)

50
TABLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses: temporarily relieves minor aches and pains due to: backache, headache, menstrual cramps, minor pain of arthritis, muscular aches, the common cold, toothache

Warnings: temporarily reduces fever

Warnings Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- hives
- rash
- shock
- skin redness

Directions:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information:

- **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**
- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY: RTE AD
30 HUNTER LANE, PA 17011

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Code No.: GODRUC5615 HJL461

Lot No.: 601R 092T

Exp. Date:

DRUG FACTS (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding:** Ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions: Do not take more than directed. The smallest effective dose should be used.

- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information:

- **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**
- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY: RTE AD
30 HUNTER LANE, PA 17011

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Code No.: GODRUC5615 HJL461

Lot No.: 601R 092T

Exp. Date:

DRUG FACTS (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use: If you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer. Right before or after heart surgery.

Ask a doctor before use if: You have problems or serious side effects from taking pain relievers or fever reducers. You have a history of stomach bleeding, vomiting, or diarrhea. You have high blood pressure, heart disease, new or worse, kidney disease, asthma, or had a stroke. You are taking a diuretic.

Ask a doctor or pharmacist before use if you are: Taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin. Under a doctor's care for any serious condition. Taking any other drug.

When using this product: Take with food or milk if stomach upset occurs.

Stop use and ask a doctor if: You experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problem or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the genital area
- any new symptoms appear

If pregnant or breast-feeding: Ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions: Do not take more than directed. The smallest effective dose should be used.

adults and children 12 years and older:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information: Read all warnings and directions before use. Store at 20-25°C (68-77°F).

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? Call 1-877-290-4008

DRUG FACTS (continued under label)

This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Tablets. 601R 0821

DISTRIBUTED BY: RTE AD
30 HUNTER LANE, PA 17011

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Code No.: GODRUC5615 HJL461

Lot No.:

Exp. Date:

PREPARE HERE

NDC 11822-7892-2

PAIN RELIEF
IBUPROFEN
IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)

50
TABLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses: temporarily relieves minor aches and pains due to: backache, headache, menstrual cramps, minor pain of arthritis, muscular aches, the common cold, toothache

Warnings: temporarily reduces fever

Warnings Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- hives
- rash
- shock
- skin redness

Directions:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information:

- **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**
- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY: RTE AD
30 HUNTER LANE, PA 17011

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Code No.: GODRUC5615 HJL461

Lot No.: 601R 092T

Exp. Date:

DRUG FACTS (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding:** Ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions: Do not take more than directed. The smallest effective dose should be used.

- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information:

- **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**
- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY: RTE AD
30 HUNTER LANE, PA 17011

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Code No.: GODRUC5615 HJL461

Lot No.: 601R 092T

Exp. Date:

DRUG FACTS (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding:** Ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy, unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions: Do not take more than directed. The smallest effective dose should be used.

- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information:

- **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**
- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY: RTE AD
30 HUNTER LANE, PA 17011

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Code No.: GODRUC5615 HJL461

Lot No.: 601R 092T

Exp. Date:

DRUG FACTS (continued under label)

This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil® Tablets. 601R 0821

DISTRIBUTED BY: RTE AD
30 HUNTER LANE, PA 17011

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

Code No.: GODRUC5615 HJL461

Lot No.:

Exp. Date:

PREPARE HERE

Adhesive Area

NDC 11822-7892-0

PAIN RELIEF IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)



24
TABLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: backache, headache, menstrual cramps, minor pain of arthritis, muscular aches, the common cold, toothache ■ temporarily reduces fever

Warnings **Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ asthma (wheezing) ■ hives ■ facial swelling ■ hives ■ rash ■ skin redness. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and older: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor

Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008
DISTRIBUTED BY: FORTA INC., 30 HUNTER LANE, CAMP HILL, PA 17011
www.fortainc.com MADE IN INDIA
Code No.: G0DRUGS515 HJL460 601R 0821

Lot No.:
Exp. Date:

Code: G0/DRUGS515

Important: Read all product information before using. Keep this box for important information.

Compare to the actual ingredients in **Real Tablets**

**PAIN RELIEF
IBUPROFEN**
IBUPROFEN TABLETS USP, 200 mg
PAIN RELIEVER/FEVER REDUCER
(NSAID)

24
TABLETS

NDC 11822-7892-0

Lot No.:
Exp. Date:

0 11822 78912 7

MADE IN INDIA
CAMP HILL, PA 17011 www.fortainc.com
DISTRIBUTED BY: FORTA INC., 30 HUNTER LANE, CAMP HILL, PA 17011
www.fortainc.com MADE IN INDIA
Code No.: G0DRUGS515 HJL460 601R 0821

This product is not manufactured or distributed by Forta Consumer Healthcare, owner of the registered trademark name, Real Tablets.

HJCN2205

Drug Facts

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better

Drug Facts (continued)

Directions

- do not take more than directed ■ the smallest effective dose should be used

adults and children 12 years and older:

- take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information ■ read all warnings and

Drug Facts (continued)

Directions before use

- store at 20-25°C (68-77°F)

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? Call 1-877-290-4008

WARNING: Do not use this product if the seal over the mouth of the bottle is cut, torn, broken or missing.

IBUPROFEN
ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-7892
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	114
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-7892-5	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	
2	NDC:11822-7892-0	24 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	
3	NDC:11822-7892-9	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	
4	NDC:11822-7892-2	50 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	
5	NDC:11822-7892-7	200 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

ANDA	ANDA091239	06/01/2020	
------	------------	------------	--

Labeler - RITE AID CORPORATION (014578892)

Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LTD		925822975	manufacture(11822-7892)

Revised: 10/2021

RITE AID CORPORATION