

IBUPROFEN- ibuprofen tablet, coated
RITE AID

612R RITE AID 1822 7893 IBUPROFEN CAPLETS 200 MG ORANGE

DRUG FACTS

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pains due to:

- backache
- headache
- menstrual cramps
- minor pain of arthritis
- muscular aches
- the common cold
- toothache

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's, except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problem or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of the body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- **adults and children 12 years and over:**

take 1 caplet every 4 to 6 hours while symptoms persist

if pain or fever does not respond to 1 caplet, 2 caplets may be used

do not exceed 6 caplets in 24 hours, unless directed by a doctor.

- **children under 12 years:**

ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD & C yellow # 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments? call **1-877-290-4008**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ temporarily relieves minor aches and pains due to:
● backache ● headache ● menstrual cramps ● minor pain of arthritis ● muscular aches ● the common cold ● toothache ■ temporarily reduces fever

Warnings ■ **Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ rash ■ shock ■ skin redness, if facial swelling ■ throat ■ rash ■ skin redness, if an allergic reaction occurs. Stop use and seek medical help right away. ■ **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed ■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. ■ **Pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. ■ **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ **adults and children 12 years and over:** ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ **do not exceed 6 caplets in 24 hours, unless directed by a doctor** ■ **children under 12 years:** ■ ask a doctor

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

DISTRIBUTED BY: RITE AID **SATISFACTION GUARANTEE:**
30 HUNTER LANE, CAMP HILL, PA 17011. If you're not satisfied, we'll happily refund your money. 6/2R 0821

www.riteaid.com **MADE IN INDIA**
HGL472 Code No.: GODRUGS615
Lot No.:

Exp. Date:

NDC 11822-7893-5

PAIN RELIEF IBUPROFEN

IBUPROFEN CAPLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)



100
CAPLETS**
(*CAPSULE-SHAPED
TABLETS)

Read all product information before using. Keep the box for important information.

Uses ■ temporarily relieves minor aches and pains due to: backache, headache, menstrual cramps, minor pain of arthritis, muscular aches, the common cold, toothache

Warnings ■ temporarily reduces fever

Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ asthma (wheezing) ■ blisters ■ facial swelling ■ hives ■ rash ■ shock ■ skin reddening ■ skin itching ■ stomach bleeding. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problem or stroke: ■ chest pain ■ trouble breathing

Directions ■ do not take more than directed ■ the smallest effective dose should be used

adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years: ■ ask a doctor

Other information ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Other information ■ TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ■ ask a doctor ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

Coating/Print Omit Area

Lot No.:
Exp. Date:
3 11822 7893 5 0

MADE IN INDIA
COATED WITH
IF YOU'RE NOT SURE, CALL
DISTRIBUTOR: RITE AID
SANTERSON GURANTER
OWNER OF THE REGISTERED TRADEMARK: MORTON
JOHNSON CONSUMER INC., MEDICAL CONSUMER HEALTHCARE DIVISION
812R 0821

Code: G0DRUGS515

Read all product information before using.
Keep the box for important information.

Drug Facts

Active ingredient (in each caplet) Purpose
Ibuprofen
200 mg (NSAID) *.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:
● backache ● headache ● menstrual cramps
● minor pain of arthritis ● muscular aches
● the common cold ● toothache
temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:
■ asthma (wheezing) ■ blisters ■ facial swelling
■ hives ■ rash ■ shock ■ skin reddening
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you: ■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood-thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if

■ you have problems or serious side effects from taking pain relievers or fever reducers
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are

■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problem or stroke:
■ chest pain ■ trouble breathing

Drug Facts (continued)

● weakness in one part or side of the body
● slurred speech ● leg swelling
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ do not take more than directed
■ the smallest effective dose should be used

adults and children 12 years and over: ■ take 1 caplet every 4 to 6 hours while symptoms persist

■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years: ■ ask a doctor

Other information

■ read all warnings and directions before use
■ store at 20-25°C (68-77°F)

Compare to the active ingredient of Motrin® 100 Tablets

PAIN RELIEF

IBUPROFEN

IBUPROFEN CAPLETS USP, 200 mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

ACTUAL SIZE

100 CAPLETS**

(*CAPSULE-SHAPED TABLETS)



NDC 11822-7893-5

Coating/Print Omit Area

3 11822 7893 5 0

Drug Facts (continued)
Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, FD&C Yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

E827R1

Coating/Print Omit Area

Lot No.:

Exp. Date:

SATISFACTION GUARANTEE:
If you're not satisfied, we'll happily refund your money.
30 HUNTER LANE
CAMP HILL, PA 17011
www.riteaid.com MADE IN INDIA
Code No.: G0DRUGS515 H6L467
612R 0821

IBUPROFEN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-7893
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE ((CAPLET))	Size	10mm
Flavor		Imprint Code	120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-7893-5	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	
2	NDC:11822-7893-2	50 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	06/01/2020	

Labeler - RITE AID (014578892)

Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LTD		925822975	manufacture(11822-7893)

Revised: 1/2022

RITE AID