

IBUPROFEN- ibuprofen tablet, coated
RITE AID

611R RITE AID 11822 8925 IBUPROFEN TABLETS 200 MG ORANGE

DRUG FACTS

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pains due to:

- backache
- headache
- menstrual cramps
- minor pain of arthritis
- muscular aches
- the common cold
- toothache

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- asthma (wheezing)
- blisters
- facial swelling
- hives
- rash
- shock
- skin reddening

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAID's, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problem or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of the body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- **adults and children 12 years and over:**

take 1 tablet every 4 to 6 hours while symptoms persist
if pain or fever does not respond to 1 tablet, 2 tablets may be used
do not exceed 6 tablets in 24 hours, unless directed by a doctor.

- **children under 12 years:**

ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

colloidal silicon dioxide, croscarmellose sodium, FD & C yellow # 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments? Call 1-877-290-4008

NDC 11822-8925-5
**PAIN RELIEF
IBUPROFEN**
IBUPROFEN TABLETS USP, 200 mg
PAIN RELIEVER
FEVER REDUCER
(NSAID)
100
TABLETS

Read all product information before using. Keep the box for important information.

Uses ■ temporarily relieves minor aches and pains due to: backache, headache, menstrual cramps, minor pain of arthritis, muscular aches, the common cold, toothache
■ temporarily reduces fever

Warnings **Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ asthma (wheezing) ■ blisters ■ facial swelling ■ hives ■ rash ■ shock ■ skin reddening. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ **do not take more than directed** ■ **the smallest effective dose should be used** ■ **adults and children 12 years and over:** ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ **children under 12 years:** ■ ask a doctor

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Questions or comments? Call 1-877-290-4008

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money.

DISTRIBUTED BY: RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011
www.riteaid.com **MADE IN INDIA**
HILL471 Code No.: GO/DRUGS/515 611R 0821

Lot No.:
Exp. Date:

Coating/Print
Omit Area



This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., Medical Consumer Healthcare Division, owner of the registered trademark Motrin® Tablets, 6118 (02) DISTRIBUTED BY: RTE MID, 30 HUNTER LANE, CAMP HILL, PA 17011
www.rteusa.com MADE IN INDIA
SAMS PRACTICE DISTRIBUTOR: If you're not satisfied, we'll happily refund your money.

COLE-CONDUCTS15

HICH2220

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen
200 mg (NSAID)* Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - headache
 - menstrual cramps
 - minor pain of arthritis
 - muscular aches
 - the common cold
 - toothache
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- hives
- rash
- shock
- skin reddening
- facial swelling
- skin redness

If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
 - right before or after heart surgery
- Ask a doctor before use if**
- you have problems or serious side effects from taking pain relievers or fever reducers
 - the stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 - you are taking a diuretic.

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- Stop use and ask a doctor if**
- you experience any of the following signs of stomach bleeding:
 - have bloody or black stools
 - feel faint
 - vomit blood
 - have stomach pain that does not get better
 - have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing

Drug Facts (continued)

- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

if pregnant or breast-feeding, ask a health professional before use. It is especially important to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and older:

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)



Coating
Omit

Compare to the active ingredient in Motrin® IB Tablets*

ACTUAL SIZE

100 TABLETS

PAIN RELIEF
IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER/FEVER REDUCER
(NSAID)



NDC 11822-8925-5

E827R1

Coating/Print Omit Area



Coating
Omit

Drug Facts (continued)
Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, FD&C yellow #3 aluminum lake, hydroxypropyl methylcellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, Talc, titanium dioxide
Questions or comments? call 1-877-298-4008
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if: ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are: ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if: ■ you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools

■ you have stomach pain that does not get better

- chest pain
- trouble breathing
- weakness in one part or side of the body
- slurred speech
- leg swelling

■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-2222).

Directions ■ do not take more than directed

■ the smallest effective dose should be used

adults and children 12 years ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor

children under 12 years: ■ ask a doctor

Other information ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call 1-877-298-4008

Compare to the active ingredient in Motrin® IB Tablets!

NDC 11822-8925-9

PAIN RELIEF

IBUPROFEN

IBUPROFEN TABLETS USP, 200 mg

PAIN RELIEVER
FEVER REDUCER
(NSAID)

ACTUAL SIZE

500 TABLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Ibuprofen 200 mg (NSAID)*. Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscle aches
 - minor pain of arthritis
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- facial swelling
- asthma (wheezing)
- hives
- rash
- shock
- skin redness

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued under label)

†This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Motrin® IB Tablets. 611R 0821

DISTRIBUTED BY: RATE AID, 30 HUNTER LANE, CAMP HILL, PA 17011

MADE IN INDIA

SATISFACTION GUARANTEE: If you're not satisfied, we'll happily refund your money. Code No. 600005856 | 1118.09

Lot No.: **FREE HERE**

Exp. Date:

0 11822-8925-9 6

IBUPROFEN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-8925
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M280L1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6130)	

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	115
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-8925-5	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	
2	NDC:11822-8925-9	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	06/01/2020	

Labeler - RITE AID (014578892)

Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LTD		925822975	manufacture(11822-8925)

Revised: 1/2022

RITE AID