ARTH RX TOPICAL ANALGESIC- methyl nicotinate, caps aicin lotion HCD Sales

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Arth RX Topical Analgesic

Active Ingredients Purpose

Methyl nicotinate 0.5% External Analgesic Capsaicin 0.025 % External Analgesic

Uses: Temporarily relieves minor aches and pains of muscles and joints due to:

- Arthritis - Strains - Sprains

WARNINGS

FOR EXTERNAL USE ONLY

When Using This Product

- Avoid contact with eyes
- Do not apply to wounds or damaged skin
- Do not bandage tightly

Stop use and ask doctor if

- Condition worsens
- Excessive irritation of the skin develops
- Pain persists for more than 7 days
- Symptoms clear up and occur again within a few days

Keep out of reach of children. If swallowed get medical help or contact a poison control center immediately.

Directions

- Adults apply to affected area no more than 3-4 times daily
- Children under 18 consult a physician
- Apply to small area of the skin as test for sensitivity at least 60 minutes prior to applying to wide area.

Inactive Ingredients

Isopropanol, water, extracts of arnica and rue, polyacrylamide, C 13-14 isoparaffin. tween 80, laureth-7, boswellia, extract of chamomile, aloe, fragrance.

Questions or comments

Toll free 1-800-729-8446

Products containing capsaicin may cause a burning, stinging or tingling sensation where applied. This is a natural reaction and varies with each individual usually decreasing or disappearing after continued use. Please read package insert prior to use. may be removed from the test area by wiping or olive oil.

Distributed by: HCD sales, Inc.

Tampa, FL 33549

www.hcdsales.com

arth.Rx TOPICAL ANALGESIC ARTHRITIS PAIN RELIEF PENETRATING PAIN RELIEF, DOUBLE ACTION FORMULA WITH DOCTOR RECOMMENDED CAPSAICIN NEW WT 3 OZ



Active Ingredients Methyl nicotinate 0.5% External Analgesia Capsaicin 0.025%

Purpose External Analgesic

Uses: Temporarily relieves minor aches and pains of muscles and joints due to: Arthritis • Strains • Sprains

FOR EXTERNAL USE ONLY

When Using This Product

- Avoid contact with eyes
- Do not apply to wounds or damaged skin
- Do not bandage tightly

Stop use and ask doctor if

- Condition worsens
- Excessive irritation of the skin develops
- Pain persists for more than 7 days
- Symptoms clear up and occur again within a few days

Keep out of reach of children. If swallowed get medical help or contact a poison control center immediately.

Directions

- Adults apply to affected area no more than 3-4 times daily
- Children under 18 consult a physician
- Apply to small area of the skin as test for sensitivity at least 60 minutes prior to applying to wide area.



WITH DOCTOR RECOMMENDED CAPSAICIN

NET WT 3 0Z (85 g)

Drug Facts (continued)

Inactive Ingredients

Isopropanol, water, extracts of arnica and rue, polyacrylamide, C 13-14 isoparaffin, tween 80, laureth-7, boswellia, extract of chamomile, aloe, fragrance.

Questions or comments Toll free 1-800-729-8446

Products containing capsaicin may cause a burning, stinging or tingling sensation where applied. This is a natural reaction and varies with each individual usually decreasing or disappearing after continued use. Please read package insert prior to use. may be removed from the test area by wiping with rubbing alcohol or olive oil.

Distributed by: HCD Sales, Inc. Tampa, FL 33549 www.hcdsales.com

0.025 g in 100 g

ARTH RX TOPICAL ANALGESIC

methyl nicotinate, capsaicin lotion

Product Information

HUMAN OTC DRUG NDC:76069-100 Product Type Item Code (Source)

Route of Administration TOPICAL

CAPSAICIN (UNII: S07O44R1ZM) (CAPSAICIN - UNII:S07O44R1ZM)

Active Ingredient/Active Moiety Basis of Strength Ingredient Name Strength METHYL NICOTINATE (UNII: 7B1AVU9DJN) (NIACIN - UNII:2679MF687A) METHYL NICOTINATE 0.5 g in 100 g

CAPSAICIN

Inactive Ingredients			
Ingredient Name	Strength		
ISOPROPYL ALCOHOL (UNII: ND2M416302)			
WATER (UNII: 059QF0KO0R)			
POLYQUATERNIUM-7 (70/30 ACRYLAMIDE/DADMAC; 1600 KD) (UNII: 0L414VCS5Y)			
C13-14 ISOPARAFFIN (UNII: E4F12ROE70)			
POLYSORBATE 80 (UNII: 6 OZP39 ZG8 H)			
LAURETH-7 (UNII: Z95S6G8201)			

ALOE (UNII: V5VD430 Y	7W9)			
Packaging		M. I. d. and a D. a	W 1 c . E 15 .	
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:/6069-100-03	85 g in 1 BOTTLE; Type 0: Not a Combination Product	02/05/2017		
Date of the Table of the Company of				
Marketing Information				
Marketing Categor	y Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not fina	al part348	02/12/2011		

Labeler - HCD Sales (137317314)

Revised: 2/2017 HCD Sales