

**EFAVIRENZ, EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE -
efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet, film coated
Cipla USA Inc.**

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use EFAVIRENZ, EMTRICITABINE, AND TENOFOVIR DISOPROXIL FUMARATE TABLET safely and effectively. See full prescribing information for EFAVIRENZ, EMTRICITABINE, AND TENOFOVIR DISOPROXIL FUMARATE TABLET.

**EFAVIRENZ, EMTRICITABINE, AND TENOFOVIR DISOPROXIL FUMARATE tablets, for oral use
Initial U.S. Approval: 2006**

WARNING: POSTTREATMENT ACUTE EXACERBATION OF HEPATITIS B

See full prescribing information for complete boxed warning.

Severe acute exacerbations of hepatitis B virus (HBV) have been reported in patients coinfecting with HBV and HIV-1 who have discontinued products containing emtricitabine (FTC) and/or tenofovir disoproxil fumarate (TDF), and may occur with discontinuation of Efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet. Closely monitor hepatic function with both clinical and laboratory follow-up for at least several months in patients who are coinfecting with HIV-1 and HBV and discontinue Efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet. If appropriate, initiation of anti-hepatitis B therapy may be warranted. (5.1)

RECENT MAJOR CHANGES

Warnings and Precautions

Nervous System Symptoms (5.6) 10/2019
Immune Reconstitution Syndrome (5.12) 10/2019

INDICATIONS AND USAGE

Efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet is a three-drug combination of efavirenz (EFV), a non-nucleoside reverse transcriptase inhibitor, and emtricitabine (FTC) and tenofovir disoproxil fumarate (TDF), both HIV-1 nucleoside analog reverse transcriptase inhibitors, and is indicated as a complete regimen or in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and pediatric patients weighing at least 40 kg. (1)

DOSAGE AND ADMINISTRATION

- Testing: Consult Full Prescribing Information for important testing recommendations prior to initiation and during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. (2.1)
- Recommended dosage in adults and pediatric patients weighing at least 40 kg: One tablet once daily taken orally on an empty stomach, preferably at bedtime. (2.2)
- Renal impairment: Not recommended in patients with estimated creatinine clearance below 50 mL/min. (2.3)
- Hepatic impairment: Not recommended in patients with moderate to severe hepatic impairment. (2.4)
- Dosage adjustment with rifampin coadministration: An additional 200 mg/day of efavirenz is recommended for patients weighing 50 kg or more. (2.5)

DOSAGE FORMS AND STRENGTHS

Tablets: 600 mg of efavirenz, 200 mg of emtricitabine, and 300 mg of tenofovir disoproxil fumarate. (3)

CONTRAINDICATIONS

- Previously demonstrated hypersensitivity (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to efavirenz, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. (4)
- Coadministration with voriconazole. (4)
- Coadministration with elbasvir/grazoprevir. (4)

WARNINGS AND PRECAUTIONS

- Rash: Discontinue if severe rash develops. (5.2,6.1)
- Hepatotoxicity: Monitor liver function tests before and during treatment in patients with underlying

hepatic disease, including hepatitis B or C coinfection, marked transaminase elevations, or who are taking medications associated with liver toxicity. Among reported cases of hepatic failure, a few occurred in patients with no pre-existing hepatic disease. (5.3, 6.2,8.7)

- Risk of adverse reactions or loss of virologic response due to drug interactions: Consult full prescribing information prior to and during treatment for important potential drug interactions. Consider alternatives to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablets in patients taking other medications with a known risk of Torsade de Pointes or in patients at higher risk of Torsade de Pointes. (5.4)
- Serious psychiatric symptoms: Immediate medical evaluation is recommended. (5.5, 6.1)
- Nervous system symptoms (NSS): NSS are frequent, usually begin 1–2 days after initiating therapy, and resolve in 2–4 weeks. Dosing at bedtime may improve tolerability. NSS are not predictive of onset of psychiatric symptoms. (2.2, 5.6)
- New onset or worsening renal impairment: Can include acute renal failure and Fanconi syndrome. Prior to initiation and during use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, assess serum creatinine, estimated creatinine clearance, urine glucose, and urine protein in all patients. In patients with chronic kidney disease, also assess serum phosphorus. Avoid administering Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with concurrent or recent use of nephrotoxic drugs. (5.7)
- Embryo fetal toxicity: Fetal harm may occur when administered to a pregnant woman during the first trimester. Avoid pregnancy while receiving Efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet and for 12 weeks after discontinuation. (5.8, 8.1)
- Decreases in bone mineral density (BMD): Consider assessment of BMD in patients with a history of pathological fracture or other risk factors for osteoporosis or bone loss. (5.9)
- Convulsions: Use caution in patients with a history of seizures. (5.10)
- Lactic acidosis/severe hepatomegaly with steatosis: Discontinue treatment in patients who develop symptoms or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity. (5.11)
- Immune reconstitution syndrome: May necessitate further evaluation and treatment. (5.12)
- Redistribution/accumulation of body fat: Observed in patients receiving antiretroviral therapy. (5.13)

----- **ADVERSE REACTIONS** -----

Most common adverse reactions (incidence greater than or equal to 10%) observed in an active-controlled clinical trial of EFV, FTC, and TDF are diarrhea, nausea, fatigue, headache, dizziness, depression, insomnia, abnormal dreams, and rash. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Cipla Ltd. at 1-866-604-3268 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **DRUG INTERACTIONS** -----

- Consult Full Prescribing Information prior to and during treatment for important potential drug interactions. (4, 5.4,7)
- HIV-1 protease inhibitors: Coadministration of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with either lopinavir/ritonavir or darunavir and ritonavir increases tenofovir concentrations. Monitor for evidence of tenofovir toxicity. Coadministration of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with either atazanavir or atazanavir and ritonavir is not recommended. (7.3)

----- **USE IN SPECIFIC POPULATIONS** -----

- Pregnancy: Avoid pregnancy while receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate and for 12 weeks after discontinuation. (5.8, 8.3)
- Lactation: Breastfeeding is not recommended. (8.2)
- Females and Males of Reproductive Potential: Pregnancy testing and contraception are recommended. (8.3)
- Pediatrics: The incidence of rash was higher than in adults. (5.2, 6.1)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 3/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: POSTTREATMENT ACUTE EXACERBATION OF HEPATITIS B

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Testing Prior to Initiation and During Treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet

2.2 Recommended Dosage for Adults and Pediatric Patients Weighing at Least 40 kg

2.3 Not Recommended in Patients with Moderate or Severe Renal Impairment

2.4 Not Recommended in Patients with Moderate to Severe Hepatic Impairment

2.5 Dosage Adjustment with Rifampin

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Severe Acute Exacerbation of Hepatitis B in Patients Coinfected with HIV-1 and HBV

5.2 Rash

5.3 Hepatotoxicity

5.4 Risk of Adverse Reactions or Loss of Virologic Response Due to Drug Interactions

5.5 Psychiatric Symptoms

5.6 Nervous System Symptoms

5.7 New Onset or Worsening Renal Impairment

5.8 Embryo-Fetal Toxicity

5.9 Bone Loss and Mineralization Defects

5.10 Convulsions

5.11 Lactic Acidosis/Severe Hepatomegaly with Steatosis

5.12 Immune Reconstitution Syndrome

5.13 Fat Redistribution

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Efavirenz

7.2 Drugs Affecting Renal Function

7.3 Established and Potentially Significant Interactions

7.4 Efavirenz Assay Interference

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

8.7 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: POSTTREATMENT ACUTE EXACERBATION OF HEPATITIS B

Severe acute exacerbations of hepatitis B virus (HBV) have been reported in patients who are coinfectd with HIV-1 and HBV and have discontinued products containing emtricitabine (FTC) and/or tenofovir disoproxil fumarate (TDF), which are components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.

Closely monitor hepatic function with both clinical and laboratory follow-up for at least several months in patients who are coinfectd with HIV-1 and HBV and discontinue Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. If appropriate, initiation of anti-hepatitis B therapy may be warranted [see Warnings and Precautions (5.1)].

1 INDICATIONS AND USAGE

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is indicated as a complete regimen or in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and pediatric patients weighing at least 40 kg.

2 DOSAGE AND ADMINISTRATION

2.1 Testing Prior to Initiation and During Treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet

Prior to or when initiating Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, test patients for hepatitis B virus infection [see Warnings and Precautions (5.1)].

Prior to initiation and during use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, urine glucose and urine protein in all patients. In patients with chronic kidney disease, also assess serum phosphorus [see Warnings and Precautions (5.7)].

Monitor hepatic function prior to and during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet [see Warnings and Precautions (5.3)].

Perform pregnancy testing before initiation of Efavirenz, Emtricitabine and Tenofovir

disoproxil fumarate tablet in adolescents and adults of childbearing potential [see *Warnings and Precautions (5.8), Use in Specific Populations (8.1, 8.3)*].

2.2 Recommended Dosage for Adults and Pediatric Patients Weighing at Least 40 kg

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is a three-drug fixed-dose combination product containing 600 mg of efavirenz (EFV), 200 mg of emtricitabine (FTC), and 300 mg of tenofovir disoproxil fumarate (TDF). The recommended dosage of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in adults and pediatric patients weighing at least 40 kg is one tablet once daily taken orally on an empty stomach. Dosing at bedtime may improve the tolerability of nervous system symptoms [see *Clinical Pharmacology (12.3)*].

2.3 Not Recommended in Patients with Moderate or Severe Renal Impairment

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended in patients with moderate or severe renal impairment (estimated creatinine clearance below 50 mL/min) [see *Warnings and Precautions (5.7), Use in Specific Populations (8.6)*].

2.4 Not Recommended in Patients with Moderate to Severe Hepatic Impairment

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended in patients with moderate to severe hepatic impairment (Child-Pugh B or C) [see *Warnings and Precautions (5.3) and Use in Specific Populations (8.7)*].

2.5 Dosage Adjustment with Rifampin

If Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is co-administered with rifampin in patients weighing 50 kg or more, take one tablet of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet once daily followed by one additional 200 mg per day of efavirenz [see *Drug Interactions (7.3) and Clinical Pharmacology (12.3)*].

3 DOSAGE FORMS AND STRENGTHS

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate is available as tablets. Each tablet contains 600 mg of efavirenz, 200 mg of emtricitabine and 300 mg of tenofovir disoproxil fumarate (tenofovir DF, which is equivalent to 245 mg of tenofovir disoproxil). The tablets are pink coloured, capsule-shaped, biconvex film-coated tablet, debossed with 'C210' on one side and plain on other side.

4 CONTRAINDICATIONS

- Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is contraindicated in patients with previously demonstrated clinically significant hypersensitivity (e.g., Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to efavirenz, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet [see *Warnings and Precautions (5.2)*].
- Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is contraindicated to

be coadministered with voriconazole or elbasvir/grazoprevir [see *Drug Interactions (7.3) and Clinical Pharmacology (12.3)*].

5 WARNINGS AND PRECAUTIONS

5.1 Severe Acute Exacerbation of Hepatitis B in Patients Coinfected with HIV-1 and HBV

All patients should be tested for the presence of chronic HBV before or when initiating antiretroviral therapy [see *Dosage and Administration (2.1)*]. Severe acute exacerbations of hepatitis B (e.g., liver decompensation and liver failure) have been reported in patients who are coinfecting with HBV and HIV-1 and have discontinued FTC or TDF, two of the components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Patients who are coinfecting with HIV-1 and HBV should be closely monitored, with both clinical and laboratory follow-up for at least several months after stopping treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. If appropriate, initiation of anti-hepatitis B therapy may be warranted, especially in patients with advanced liver disease or cirrhosis, since posttreatment exacerbation of hepatitis may lead to hepatic decompensation and liver failure.

5.2 Rash

In controlled clinical trials, 26% (266/1,008) of adult subjects treated with 600 mg EFV experienced new-onset skin rash compared with 17% (111/635) of those treated in control groups. Rash associated with blistering, moist desquamation, or ulceration occurred in 0.9% (9/1,008) of subjects treated with EFV. The incidence of Grade 4 rash (e.g., erythema multiforme, Stevens-Johnson syndrome) in adult subjects treated with EFV in all trials and expanded access was 0.1%. Rashes are usually mild-to-moderate maculopapular skin eruptions that occur within the first 2 weeks of initiating therapy with EFV (median time to onset of rash in adults was 11 days) and, in most subjects continuing therapy with EFV, rash resolves within 1 month (median duration, 16 days). The discontinuation rate for rash in adult clinical trials was 1.7% (17/1,008). Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet can be reinitiated in patients interrupting therapy because of rash. Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet should be discontinued in patients developing severe rash associated with blistering, desquamation, mucosal involvement, or fever. Appropriate antihistamines and/or corticosteroids may improve the tolerability and hasten the resolution of rash. For patients who have had a life-threatening cutaneous reaction (e.g., Stevens-Johnson syndrome), alternative therapy should be considered [see *Contraindications (4)*].

Experience with EFV in subjects who discontinued other antiretroviral agents of the NNRTI class is limited. Nineteen subjects who discontinued nevirapine because of rash have been treated with EFV. Nine of these subjects developed mild-to-moderate rash while receiving therapy with EFV, and two of these subjects discontinued because of rash.

Rash was reported in 59 of 182 pediatric subjects (32%) treated with EFV [see *Adverse Reactions (6.1)*]. Two pediatric subjects experienced Grade 3 rash (confluent rash with fever, generalized rash), and four subjects had Grade 4 rash (erythema multiforme). The median time to onset of rash in pediatric subjects was 28 days (range 3-1,642 days). Prophylaxis with appropriate antihistamines before initiating therapy with Efavirenz,

Emtricitabine and Tenofovir disoproxil fumarate tablet in pediatric patients should be considered.

5.3 Hepatotoxicity

Postmarketing cases of hepatitis, including fulminant hepatitis progressing to liver failure requiring transplantation or resulting in death, have been reported in patients treated with EFV, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Reports have included patients with underlying hepatic disease, including coinfection with hepatitis B or C, and patients without pre-existing hepatic disease or other identifiable risk factors [see *Warnings and Precautions (5.1)*].

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended for patients with moderate or severe hepatic impairment. Careful monitoring is recommended for patients with mild hepatic impairment receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet [see *Adverse Reactions (6.2)* and *Use in Specific Populations (8.7)*].

Monitoring of liver enzymes before and during treatment is recommended for all patients [see *Dosage and Administration (2.1)*]. Consider discontinuing Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in patients with persistent elevations of serum transaminases to greater than five times the upper limit of the normal range.

Discontinue Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet if elevation of serum transaminases is accompanied by clinical signs or symptoms of hepatitis or hepatic decompensation [see *Adverse Reactions (6.1)*].

5.4 Risk of Adverse Reactions or Loss of Virologic Response Due to Drug Interactions

The concomitant use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and other drugs may result in potentially significant drug interactions [see *Contraindications (4)* and *Drug Interactions (7.3)*], some of which may lead to:

- Loss of therapeutic effect of concomitant drug or Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and possible development of resistance.
- Possible clinically significant adverse reaction from greater exposures of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet or concomitant drug.

QTc prolongation has been observed with the use of EFV [see *Drug Interactions (7.1)* and *Clinical Pharmacology (12.2)*]. Consider alternatives to Efavirenz, Emtricitabine, and Tenofovir disoproxil fumarate tablet when coadministered with a drug with a known risk of Torsade de Pointes or when administered to patients at higher risk of Torsade de Pointes.

See Table 3 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations. Consider the potential for drug interactions prior to and during Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet therapy and review concomitant medications during Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet therapy [see *Dosage and Administration (2.5)*, *Contraindications (4)*, and *Drug Interactions (7)*].

5.5 Psychiatric Symptoms

Serious psychiatric adverse experiences have been reported in patients treated with EFV, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. In controlled trials of 1,008 subjects treated with regimens containing EFV for a mean of 2.1 years and 635 subjects treated with control regimens for a mean of 1.5 years, the frequency (regardless of causality) of specific serious psychiatric events among subjects who received EFV or control regimens, respectively, were: severe depression (2.4%, 0.9%), suicidal ideation (0.7%, 0.3%), nonfatal suicide attempts (0.5%, 0%), aggressive behavior (0.4%, 0.5%), paranoid reactions (0.4%, 0.3%), and manic reactions (0.2%, 0.3%). When psychiatric symptoms similar to those noted above were combined and evaluated as a group in a multifactorial analysis of data from Study AI266006 (006, NCT00002410), a Phase 3 randomized, open-label trial of EFV-containing regimens versus controls in 1,266 subjects (median follow-up 180 weeks, 102 weeks, and 76 weeks for subjects treated with EFV + zidovudine + lamivudine, EFV + indinavir, and indinavir + zidovudine + lamivudine, respectively), treatment with EFV was associated with an increase in the occurrence of these selected psychiatric symptoms. Other factors associated with an increase in the occurrence of these psychiatric symptoms were history of injection drug use, psychiatric history, and receipt of psychiatric medication at trial entry; similar associations were observed in both the EFV and control treatment groups. In Study 006, onset of new serious psychiatric symptoms occurred throughout the trial for both EFV-treated and control-treated subjects. One percent of EFV-treated subjects discontinued or interrupted treatment because of one or more of these selected psychiatric symptoms. There have also been occasional postmarketing reports of death by suicide, delusions, and psychosis-like behavior, although a causal relationship to the use of EFV cannot be determined from these reports. Postmarketing cases of catatonia have also been reported and may be associated with increased EFV exposure. Patients with serious psychiatric adverse experiences should seek immediate medical evaluation to assess the possibility that the symptoms may be related to the use of EFV, and if so, to determine whether the risks of continued therapy outweigh the benefits [see *Adverse Reactions (6)*].

5.6 Nervous System Symptoms

Fifty-three percent (531/1,008) of subjects receiving EFV in controlled trials reported central nervous system symptoms (any grade, regardless of causality) compared to 25% (156/635) of subjects receiving control regimens. These symptoms included dizziness (28.1% of the 1,008 subjects), insomnia (16.3%), impaired concentration (8.3%), somnolence (7.0%), abnormal dreams (6.2%), and hallucinations (1.2%). Other reported symptoms were euphoria, confusion, agitation, amnesia, stupor, abnormal thinking, and depersonalization. The majority of these symptoms were mild to moderate (50.7%); symptoms were severe in 2.0% of subjects. Overall, 2.1% of subjects discontinued therapy as a result. These symptoms usually begin during the first or second day of therapy and generally resolve after the first 2-4 weeks of therapy. After 4 weeks of therapy, the prevalence of nervous system symptoms of at least moderate severity ranged from 5% to 9% in subjects treated with regimens containing EFV and from 3% to 5% in subjects treated with a control regimen. Patients should be informed that these common symptoms were likely to improve with continued therapy and were not predictive of subsequent onset of the less frequent psychiatric symptoms [see *Warnings and Precautions (5.5)*]. Dosing at bedtime may improve the tolerability of these nervous system symptoms [see *Dosage and Administration (2.2)*].

Analysis of long-term data from Study 006 showed that, beyond 24 weeks of therapy,

the incidences of new-onset nervous system symptoms among EFV-treated subjects were generally similar to those in the indinavir-containing control arm.

Late-onset neurotoxicity, including ataxia and encephalopathy (impaired consciousness, confusion, psychomotor slowing, psychosis, delirium), may occur months to years after beginning EFV therapy. Some events of late-onset neurotoxicity have occurred in patients with CYP2B6 genetic polymorphisms which are associated with increased EFV levels despite standard dosing of EFV. Patients presenting with signs and symptoms of serious neurologic adverse experiences should be evaluated promptly to assess the possibility that these events may be related to EFV use, and whether discontinuation of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is warranted.

Patients receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet should be alerted to the potential for additive central nervous system effects when Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is used concomitantly with alcohol or psychoactive drugs.

Patients who experience central nervous system symptoms such as dizziness, impaired concentration, and/or drowsiness should avoid potentially hazardous tasks such as driving or operating machinery.

5.7 New Onset or Worsening Renal Impairment

Emtricitabine and tenofovir are principally eliminated by the kidney; however, EFV is not. Renal impairment, including cases of acute renal failure and Fanconi syndrome (renal tubular injury with severe hypophosphatemia), has been reported with the use of TDF, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate [see *Adverse Reactions (6.2)*].

Prior to initiation and during use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, urine glucose, and urine protein in all patients. In patients with chronic kidney disease, also assess serum phosphorus. Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended in patients with moderate or severe renal impairment (estimated creatinine clearance below 50 mL/min).

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet should be avoided with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple non-steroidal anti-inflammatory drugs [NSAIDs]) [see *Drug Interactions (7.2)*]. Cases of acute renal failure after initiation of high-dose or multiple NSAIDs have been reported in HIV-infected patients with risk factors for renal dysfunction who appeared stable on TDF. Some patients required hospitalization and renal replacement therapy. Alternatives to NSAIDs should be considered, if needed, in patients at risk for renal dysfunction.

Persistent or worsening bone pain, pain in extremities, fractures, and/or muscular pain or weakness may be manifestations of proximal renal tubulopathy and should prompt an evaluation of renal function in patients at risk of renal dysfunction.

Discontinue Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in patients who develop clinically significant decreases in renal function or evidence of Fanconi syndrome.

5.8 Embryo-Fetal Toxicity

Efavirenz may cause fetal harm when administered during the first trimester of pregnancy. Advise adults and adolescents of childbearing potential who are receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet to avoid pregnancy while receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and for 12 weeks after discontinuation [see *Dosage and Administration (2.1)*, *Use in Specific Populations (8.1, 8.3)*].

5.9 Bone Loss and Mineralization Defects

Bone Mineral Density

In clinical trials in HIV-1 infected adults, TDF (a component of Efavirenz, Emtricitabine and tenofovir disoproxil fumarate tablet) was associated with slightly greater decreases in bone mineral density (BMD) and increases in biochemical markers of bone metabolism, suggesting increased bone turnover relative to comparators. Serum parathyroid hormone levels and 1,25 Vitamin D levels were also higher in subjects receiving TDF.

Clinical trials evaluating TDF in pediatric and adolescent subjects were conducted. Under normal circumstances, BMD increases rapidly in pediatric patients. In HIV-1 infected subjects aged 2 years to less than 18 years, bone effects were similar to those observed in adult subjects and suggest increased bone turnover. Total body BMD gain was less in the TDF-treated HIV-1 infected pediatric subjects as compared to the control groups. Similar trends were observed in chronic hepatitis-B infected adolescent subjects aged 12 years to less than 18 years. In all pediatric trials, skeletal growth (height) appeared to be unaffected.

The effects of TDF-associated changes in BMD and biochemical markers on long-term bone health and future fracture risk are unknown. Assessment of BMD should be considered for adult and pediatric patients who have a history of pathologic bone fracture or other risk factors for osteoporosis or bone loss. Although the effect of supplementation with calcium and vitamin D was not studied, such supplementation may be beneficial for all patients. If bone abnormalities are suspected, then appropriate consultation should be obtained.

Mineralization Defects

Cases of osteomalacia associated with proximal renal tubulopathy, manifested as bone pain or pain in extremities and which may contribute to fractures, have been reported in association with TDF use [see *Adverse Reactions (6.2)*]. Arthralgias and muscle pain or weakness have also been reported in cases of proximal renal tubulopathy. Hypophosphatemia and osteomalacia secondary to proximal renal tubulopathy should be considered in patients at risk of renal dysfunction who present with persistent or worsening bone or muscle symptoms while receiving TDF-containing products [see *Warnings and Precautions (5.7)*].

5.10 Convulsions

Convulsions have been observed in adult and pediatric patients receiving EFV, generally in the presence of known medical history of seizures. Caution must be taken in any patient with a history of seizures.

Patients who are receiving concomitant anticonvulsant medications primarily metabolized by the liver, such as phenytoin and phenobarbital, may require periodic monitoring of

plasma levels [see *Drug Interactions (7.3)*].

5.11 Lactic Acidosis/Severe Hepatomegaly with Steatosis

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogs, including TDF and FTC, components of Efavirenz, Emtricitabine and tenofovir disoproxil fumarate tablet, alone or in combination with other antiretrovirals. Treatment with Efavirenz, Emtricitabine and tenofovir disoproxil fumarate tablet should be suspended in any patient who develops clinical or laboratory findings suggestive of lactic acidosis or pronounced hepatotoxicity (which may include hepatomegaly and steatosis even in the absence of marked transaminase elevations).

5.12 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including the components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, Guillain-Barre syndrome, and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.13 Fat Redistribution

Redistribution/accumulation of body fat, including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and "cushingoid appearance," has been observed in patients receiving antiretroviral therapy, including EFV. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in other sections of the labeling:

- Severe Acute Exacerbations of Hepatitis B in Patients Coinfected with HIV-1 and HBV [see *Warnings and Precautions (5.1)*].
- Rash [see *Warnings and Precautions (5.2)*].
- Hepatotoxicity [see *Warnings and Precautions (5.3)*].
- Psychiatric Symptoms [see *Warnings and Precautions (5.5)*].
- Nervous System Symptoms [see *Warnings and Precautions (5.6)*].
- New Onset or Worsening Renal Impairment [see *Warnings and Precautions (5.7)*].
- Embryo-Fetal Toxicity [see *Warnings and Precautions (5.8)*].
- Bone Loss and Mineralization Defects [see *Warnings and Precautions (5.9)*].
- Convulsions [see *Warnings and Precautions (5.10)*].
- Lactic Acidosis/Severe Hepatomegaly with Steatosis [see *Warnings and Precautions (5.11)*].

- Immune Reconstitution Syndrome [see Warnings and Precautions (5.12)].
- Fat Redistribution [see Warnings and Precautions (5.13)].

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Clinical Trials in Adult Subjects

Study 934 was an open-label active-controlled trial in which 511 antiretroviral-naïve subjects received either FTC + TDF administered in combination with EFV (N=257) or zidovudine (AZT)/lamivudine (3TC) administered in combination with EFV (N=254).

The most common adverse reactions (incidence greater than or equal to 10%, any severity) occurring in Study 934 include diarrhea, nausea, fatigue, headache, dizziness, depression, insomnia, abnormal dreams, and rash. Adverse reactions observed in Study 934 were generally consistent with those seen in previous trials of the individual components (Table 1).

Table 1: Selected Adverse Reactions^a (Grades 2-4) Reported in ≥5% in Either Treatment Group in Study 934 (0-144 Weeks)

	FTC+TDF+EFV^b	AZT/3TC+EFV
	N=257	N=254
Fatigue	9%	8%
Depression	9%	7%
Nausea	9%	7%
Diarrhea	9%	5%
Dizziness	8%	7%
Upper respiratory tract infections	8%	5%
Sinusitis	8%	4%
Rash Event ^c	7%	9%
Headache	6%	5%
Insomnia	5%	7%
Anxiety	5%	4%
Nasopharyngitis	5%	3%
Vomiting	2%	5%

^a. Frequencies of adverse reactions are based on all treatment-emergent adverse events, regardless of relationship to study drug.

^b. From Weeks 96 to 144 of the trial, subjects received FTC/TDF administered in combination with EFV in place of FTC+ TDF with EFV.

^c. Rash event includes rash, exfoliative rash, rash generalized, rash macular, rash maculopapular, rash pruritic, and rash vesicular.

In Study 073, subjects with stable, virologic suppression on antiretroviral therapy and no history of virologic failure were randomized to receive Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet or to stay on their baseline regimen. The adverse reactions observed in Study 073 were generally consistent with those seen in Study 934 and those seen with the individual components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet when each was administered in combination with other antiretroviral agents.

Efavirenz, Emtricitabine, or TDF

In addition to the adverse reactions in Study 934 and Study 073, the following adverse reactions were observed in clinical trials of EFV, FTC, or TDF in combination with other antiretroviral agents.

Efavirenz: The most significant adverse reactions observed in subjects treated with EFV were nervous system symptoms [see *Warnings and Precautions (5.6)*], psychiatric symptoms [see *Warnings and Precautions (5.5)*], and rash [see *Warnings and Precautions (5.2)*].

Selected adverse reactions of moderate-to-severe intensity observed in greater than or equal to 2% of EFV-treated subjects in two controlled clinical trials included pain, impaired concentration, abnormal dreams, somnolence, anorexia, dyspepsia, abdominal pain, nervousness, and pruritus.

Pancreatitis has also been reported, although a causal relationship with EFV has not been established. Asymptomatic increases in serum amylase levels were observed in a significantly higher number of subjects treated with EFV 600 mg than in control subjects.

Skin discoloration has been reported with higher frequency among FTC-treated subjects; it was manifested by hyperpigmentation on the palms and/or soles and was generally mild and asymptomatic. The mechanism and clinical significance are unknown.

Clinical Trials in Pediatric Subjects

Efavirenz: Assessment of adverse reactions is based on three pediatric clinical trials in 182 HIV-1 infected pediatric subjects who received EFV in combination with other antiretroviral agents for a median of 123 weeks. The type and frequency of adverse reactions in the three trials were generally similar to that of adult subjects with the exception of a higher incidence of rash, which was reported in 32% (59/182) of pediatric subjects compared to 26% of adults, and a higher frequency of Grade 3 or 4 rash reported in 3% (6/182) of pediatric subjects compared to 0.9% of adults [see *Warnings and Precautions (5.2)*].

Emtricitabine: In addition to the adverse reactions reported in adults, anemia and hyperpigmentation were observed in 7% and 32%, respectively, of pediatric subjects who received treatment with FTC in the larger of two open-label, uncontrolled pediatric trials (N=116).

Tenofovir DF: In a pediatric clinical trial conducted in subjects 12 to less than 18 years of age, the adverse reactions observed in pediatric subjects who received treatment with TDF (N=81) were consistent with those observed in clinical trials of TDF in adults [see *Warnings and Precautions (5.9)*].

Laboratory Abnormalities

Efavirenz, Emtricitabine and Tenofovir DF: Laboratory abnormalities observed in Study 934 were generally consistent with those seen in previous trials (Table 2).

Table 2: Significant Laboratory Abnormalities Reported in $\geq 1\%$ of Subjects in Either Treatment Group in Study 934 (0-144 Weeks)

	FTC + TDF + EFV^a	AZT/3TC + EFV
	N=257	N=254
Any \geq Grade 3 Laboratory Abnormality	30%	26%
Fasting Cholesterol (>240 mg/dL)	22%	24%
Creatine Kinase (M: >990 U/L) (F: >845 U/L)	9%	7%
Serum Amylase (>175 U/L)	8%	4%
Alkaline Phosphatase (>550 U/L)	1%	0%
AST (M: >180 U/L) (F: >170 U/L)	3%	3%
ALT (M: >215 U/L) (F: >170 U/L)	2%	3%
Hemoglobin (<8.0 mg/dL)	0%	4%
Hyperglycemia (>250 mg/dL)	2%	1%
Hematuria (>75 RBC/HPF)	3%	2%
Glycosuria ($\geq 3+$)	$<1\%$	1%
Neutrophils ($<750/\text{mm}^3$)	3%	5%
Fasting Triglycerides (>750 mg/dL)	4%	2%

^a. From Weeks 96 to 144 of the trial, subjects received FTC/TDF administered in combination with EFV in place of FTC + TDF with EFV.

Laboratory abnormalities observed in Study 073 were generally consistent with those in Study 934.

Hepatic Events: In Study 934, 19 subjects treated with EFV, FTC, and TDF and 20 subjects treated with EFV and fixed-dose zidovudine/lamivudine were hepatitis B surface antigen or hepatitis C antibody positive. Among these coinfecting subjects, one subject (1/19) in the EFV, FTC, and TDF arm had elevations in transaminases to greater than five times ULN through 144 weeks. In the fixed-dose zidovudine/lamivudine arm, two subjects (2/20) had elevations in transaminases to greater than five times ULN through 144 weeks. No HBV and/or HCV coinfecting subject discontinued from the trial due to hepatobiliary disorders [see *Warnings and Precautions (5.3)*].

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of EFV, FTC, or TDF. Because postmarketing reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Efavirenz:

Cardiac Disorders

Palpitations

Ear and Labyrinth Disorders

Tinnitus, vertigo

Endocrine Disorders

Gynecomastia

Eye Disorders

Abnormal vision

Gastrointestinal Disorders

Constipation, malabsorption

General Disorders and Administration Site Conditions

Asthenia

Hepatobiliary Disorders

Hepatic enzyme increase, hepatic failure, hepatitis

Immune System Disorders

Allergic reactions

Metabolism and Nutrition Disorders

Redistribution/accumulation of body fat [see Warnings and Precautions (5.13)], hypercholesterolemia, hypertriglyceridemia

Musculoskeletal and Connective Tissue Disorders

Arthralgia, myalgia, myopathy

Nervous System Disorders

Abnormal coordination, ataxia, encephalopathy, cerebellar coordination and balance disturbances, convulsions, hypoesthesia, paresthesia, neuropathy, tremor

Psychiatric Disorders

Aggressive reactions, agitation, delusions, emotional lability, mania, neurosis, paranoia, psychosis, suicide, catatonia

Respiratory, Thoracic and Mediastinal Disorders

Dyspnea

Skin and Subcutaneous Tissue Disorders

Flushing, erythema multiforme, photoallergic dermatitis, Stevens-Johnson syndrome

Emtricitabine: No postmarketing adverse reactions have been identified for inclusion in this section.

Tenofovir DF:

Immune System Disorders

Allergic reaction, including angioedema

Metabolism and Nutrition Disorders

Lactic acidosis, hypokalemia, hypophosphatemia

Respiratory, Thoracic, and Mediastinal Disorders

Dyspnea

Gastrointestinal Disorders

Pancreatitis, increased amylase, abdominal pain

Hepatobiliary Disorders

Hepatic steatosis, hepatitis, increased liver enzymes (most commonly AST, ALT, gamma GT)

Skin and Subcutaneous Tissue Disorders

Rash

Musculoskeletal and Connective Tissue Disorders

Rhabdomyolysis, osteomalacia (manifested as bone pain and which may contribute to fractures), muscular weakness, myopathy

Renal and Urinary Disorders

Acute renal failure, renal failure, acute tubular necrosis, Fanconi syndrome, proximal renal tubulopathy, interstitial nephritis (including acute cases), nephrogenic diabetes insipidus, renal insufficiency, increased creatinine, proteinuria, polyuria

General Disorders and Administration Site Conditions

Asthenia

The following adverse reactions, listed under the body system headings above, may occur as a consequence of proximal renal tubulopathy: rhabdomyolysis, osteomalacia, hypokalemia, muscular weakness, myopathy, hypophosphatemia.

7 DRUG INTERACTIONS

7.1 Efavirenz

Efavirenz has been shown *in vivo* to induce CYP3A and CYP2B6. Other compounds that are substrates of CYP3A or CYP2B6 may have decreased plasma concentrations when coadministered with EFV.

Drugs that induce CYP3A activity (e.g., phenobarbital, rifampin, rifabutin) would be expected to increase the clearance of EFV, resulting in lowered plasma concentrations [see *Dosage and Administration (2.2)*].

There is limited information available on the potential for a pharmacodynamic interaction between EFV and drugs that prolong the QTc interval. QTc prolongation has been observed with the use of EFV [see *Clinical Pharmacology (12.2)*]. Consider alternatives to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet when coadministered with a drug with a known risk of Torsade de Pointes.

7.2 Drugs Affecting Renal Function

FTC and tenofovir are primarily eliminated by the kidneys [see *Clinical Pharmacology (12.3)*]. Coadministration of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with drugs that are eliminated by active tubular secretion may increase concentrations of FTC, tenofovir, and/or the coadministered drug. Some examples include, but are not limited to, acyclovir, adefovir dipivoxil, cidofovir, ganciclovir, valacyclovir, valganciclovir, aminoglycosides (e.g., gentamicin), and high-dose or multiple NSAIDs [see *Warnings and Precautions (5.7)*]. Drugs that decrease renal function may increase concentrations of FTC and/or tenofovir.

7.3 Established and Potentially Significant Interactions

Other important drug interaction information for Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is summarized in Table 3. The drug interactions described are based on trials conducted with either Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, the components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet (EFV, FTC, or TDF) as individual agents, or are potential drug interactions [see *Clinical Pharmacology (12.3)*].

Table 3 Established and Potentially Significant^a Drug Interactions

Concomitant Drug Class: Drug Name	Effect	Clinical Comment
<i>HIV antiviral agents</i>		
Protease inhibitor: atazanavir	↓ atazanavir ↑ tenofovir	Coadministration of atazanavir with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended. The combined effect of EFV plus TDF on atazanavir plasma concentrations is not known. There are insufficient data to support dosing recommendations for atazanavir or atazanavir/ritonavir in combination with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
Protease inhibitor: fosamprenavir calcium	↓ amprenavir	Fosamprenavir (unboosted): Appropriate doses of fosamprenavir and Efavirenz,

		<p>Emtricitabine and Tenofovir disoproxil fumarate tablet with respect to safety and efficacy have not been established.</p> <p>Fosamprenavir/ritonavir: An additional 100 mg/day (300 mg total) of ritonavir is recommended when Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is administered with fosamprenavir/ritonavir once daily. No change in the ritonavir dose is required when Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is administered with fosamprenavir plus ritonavir twice daily.</p>
Protease inhibitor: indinavir	↓ indinavir	The optimal dose of indinavir, when given in combination with EFV, is not known. Increasing the indinavir dose to 1000 mg every 8 hours does not compensate for the increased indinavir metabolism due to EFV.
Protease inhibitor: darunavir/ritonavir lopinavir/ritonavir	↑ tenofovir ↓ lopinavir ↑ tenofovir	<p>Monitor patients receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet concomitantly with ritonavir-boosted darunavir for TDF-associated adverse reactions. Discontinue Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in patients who develop TDF-associated adverse reactions. Do not use once daily administration of lopinavir/ritonavir. Dose increase of lopinavir/ritonavir is recommended for all patients when coadministered with EFV. Refer to the Full Prescribing Information for lopinavir/ritonavir for guidance on coadministration with EFV-or tenofovir-containing regimens, such as Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Patients should be monitored for tenofovir-associated adverse reactions. Discontinue Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate</p>

		tablet in patients who develop TDF-associated adverse reactions.
Protease inhibitor: ritonavir	↑ ritonavir ↑ efavirenz	When ritonavir 500 mg every 12 hours was coadministered with EFV 600 mg once daily, the combination was associated with a higher frequency of adverse clinical experiences (e.g., dizziness, nausea, paresthesia) and laboratory abnormalities (elevated liver enzymes). Monitoring of liver enzymes is recommended when Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is used in combination with ritonavir.
Protease inhibitor: saquinavir	↓ saquinavir	Appropriate doses of the combination of EFV and saquinavir/ritonavir with respect to safety and efficacy have not been established.
CCR5 co-receptor antagonist: maraviroc	↓ maraviroc	Refer to the full prescribing information for maraviroc for guidance on coadministration with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
NRTI: didanosine	↑ didanosine	Patients receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and didanosine should be monitored closely for didanosine-associated adverse reactions. Discontinue didanosine in patients who develop didanosine-associated adverse reactions. Higher didanosine concentrations could potentiate didanosine-associated adverse reactions, including pancreatitis, and neuropathy. Suppression of CD4+ cell counts has been observed in patients receiving TDF with didanosine 400 mg daily. In patients weighing greater than 60 kg, reduce the didanosine dose to 250 mg when it is coadministered with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. In patients weighing less than 60 kg,

		reduce the didanosine dose to 200 mg when it is coadministered with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. When coadministered, Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and Videx EC may be taken under fasted conditions or with a light meal (less than 400 kcal, 20% fat).
NNRTI: Other NNRTIs	↑ or ↓ efavirenz and/or NNRTI	Combining two NNRTIs has not been shown to be beneficial. Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet contains EFV and should not be coadministered with other NNRTIs.
Integrase strand transfer inhibitor: raltegravir	↓ raltegravir	The clinical significance of this interaction has not been directly assessed.
<i>Hepatitis C antiviral agents</i>		
boceprevir	↓ boceprevir	Plasma trough concentrations of boceprevir were decreased when boceprevir was coadministered with EFV, which may result in loss of therapeutic effect. The combination should be avoided.
elbasvir/grazoprevir	↓ elbasvir ↓ grazoprevir	Coadministration of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with elbasvir/grazoprevir is contraindicated [see <i>Contraindications (4)</i>] because it may lead to loss of virologic response to elbasvir/grazoprevir.
glecaprevir/pibrentasvir	↓ glecaprevir ↓ pibrentasvir	Coadministration of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended because it may lead to reduced therapeutic effect of glecaprevir/pibrentasvir.
ledipasvir/sofosbuvir	↑ tenofovir	Patients receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and HARVONI® (ledipasvir/sofosbuvir) concomitantly should be monitored for adverse reactions associated with TDF.
simeprevir	↓ simeprevir ↔ efavirenz	Concomitant administration of simeprevir with EFV is not

		recommended because it may result in loss of therapeutic effect of simeprevir.
sofosbuvir/velpatasvir sofosbuvir/velpatasvir/ voxilaprevir	↑ tenofovir ↓ velpatasvir ↓ voxilaprevir	Coadministration of EFV-containing regimens and EPCLUSA [®] (sofosbuvir/velpatasvir) or VOSEVI [®] (sofosbuvir/velpatasvir/voxilaprevir) is not recommended.
<i>Other agents</i>		
Anticoagulant: warfarin	↑ or ↓ warfarin	Plasma concentrations and effects potentially increased or decreased by EFV.
Anticonvulsants: carbamazepine	↓ carbamazepine ↓ efavirenz	There are insufficient data to make a dose recommendation for Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Alternative anticonvulsant treatment should be used.
phenytoin phenobarbital	↓ anticonvulsant ↓ efavirenz	Potential for reduction in anticonvulsant and/or EFV plasma levels; periodic monitoring of anticonvulsant plasma levels should be conducted.
Antidepressants: bupropion	↓ bupropion	The effect of EFV on bupropion exposure is thought to be due to the induction of bupropion metabolism. Increases in bupropion dosage should be guided by clinical response, but the maximum recommended dose of bupropion should not be exceeded.
sertraline	↓ sertraline	Increases in sertraline dose should be guided by clinical response.
Antifungals: itraconazole	↓ itraconazole ↓ hydroxy- itraconazole	Since no dose recommendation for itraconazole can be made, alternative antifungal treatment should be considered.
ketoconazole	↓ ketoconazole	Drug interaction trials with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and ketoconazole have not been conducted. Efavirenz has the potential to decrease plasma concentrations of ketoconazole.
posaconazole	↓ posaconazole	Avoid concomitant use unless the benefit outweighs the risks.
voriconazole	↓ voriconazole ↑ efavirenz	Coadministration of Efavirenz, Emtricitabine and Tenofovir

		disoproxil fumarate tablet with voriconazole is contraindicated [see <i>Contraindications (4)</i>] because it may lead to reduced therapeutic effect of voriconazole and increased risk of EFV-associated adverse reactions
Anti-infective: clarithromycin	↓ clarithromycin ↑ 14-OH metabolite	Consider alternatives to macrolide antibiotics because of the risk of QT interval prolongation.
Antimycobacterial: rifabutin	↓ rifabutin	Increase daily dose of rifabutin by 50%. Consider doubling the rifabutin dose in regimens where rifabutin is given 2 or 3 times a week.
rifampin	↓ efavirenz	If Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is coadministered with rifampin to patients weighing 50 kg or more, an additional 200 mg/day of EFV is recommended.
Antimalarials: artemether/ lumefantrine atovaquone/proguanil	↓ artemether ↓ dihydroartemisinin ↓ lumefantrine ↓ atovaquone ↓ proguanil	Consider alternatives to artemether/lumefantrine because of the risk of QT interval prolongation [see <i>Warnings and Precautions (5.4)</i>]. Concomitant administration of atovaquone/proguanil with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended.
Calcium channel blockers: diltiazem	↓ diltiazem ↓ desacetyl diltiazem ↓ N-monodes-methyl diltiazem	Diltiazem dose adjustments should be guided by clinical response (refer to the full prescribing information for diltiazem). No dose adjustment of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is necessary when administered with diltiazem.
Others eg, felodipine, nifedipine, verapamil	↓ calcium channel blocker	No data are available on the potential interactions of EFV with other calcium channel blockers that are substrates of CYP3A. The potential exists for reduction in plasma concentrations of the calcium channel blocker. Dose adjustments should be guided by clinical response (refer to the full

		prescribing information for the calcium channel blocker).
HMG-CoA reductase inhibitors: atorvastatin pravastatin simvastatin	↓ atorvastatin ↓ pravastatin ↓ simvastatin	Plasma concentrations of atorvastatin, pravastatin, and simvastatin decreased with EFV. Consult the Full Prescribing Information for the HMG-CoA reductase inhibitor for guidance on individualizing the dose.
Hormonal contraceptives: Oral: ethinyl estradiol/norgestimate Implant: etonogestrel	↓ active metabolites of norgestimate ↓ etonogestrel	A reliable method of barrier contraception must be used in addition to hormonal contraceptives. Efavirenz had no effect on ethinyl estradiol concentrations, but progestin levels (norelgestromin and levonorgestrel) were markedly decreased. No effect of ethinyl estradiol/norgestimate on EFV plasma concentrations was observed. A reliable method of barrier contraception must be used in addition to hormonal contraceptives. Decreased exposure of etonogestrel may be expected. There have been postmarketing reports of contraceptive failure with etonogestrel in EFV-exposed patients.
Immunosuppressants: cyclosporine, tacrolimus, sirolimus, and others metabolized by CYP3A	↓ immuno-suppressant	Decreased exposure of the immunosuppressant may be expected due to CYP3A induction by EFV. These immunosuppressants are not anticipated to affect exposure of EFV. Dose adjustments of the immunosuppressant may be required. Close monitoring of immunosuppressant concentrations for at least 2 weeks (until stable concentrations are reached) is recommended when starting or stopping treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
Narcotic analgesic: methadone	↓ methadone	Coadministration of EFV in HIV-1 infected individuals with a history of

	<p>injection drug use resulted in signs of opiate withdrawal. Methadone dose was increased by a mean of 22% to alleviate withdrawal symptoms. Patients should be monitored for signs of withdrawal and their methadone dose increased as required to alleviate withdrawal symptoms.</p>
--	---

^a. This table is not all inclusive.

7.4 Efavirenz Assay Interference

Cannabinoid Test Interaction: Efavirenz does not bind to cannabinoid receptors. False-positive urine cannabinoid test results have been reported with some screening assays in uninfected and HIV-infected subjects receiving EFV. Confirmation of positive screening tests for cannabinoids by a more specific method is recommended.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Antiretroviral Pregnancy Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in adults and adolescents exposed to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet during pregnancy. Healthcare providers are encouraged to register patients by calling the Antiretroviral Pregnancy Registry (APR) at (800) 258-4263.

Risk Summary

There are retrospective case reports of neural tube defects in infants whose mothers were exposed to EFV-containing regimens in the first trimester of pregnancy. Prospective pregnancy data from the APR are not sufficient to adequately assess this risk. Although a causal relationship has not been established between exposure to EFV in the first trimester and neural tube defects, similar malformations have been observed in studies conducted in monkeys at doses similar to the human dose (*see Data*). In addition, fetal and embryonic toxicities occurred in rats at a dose 10 times less than the human exposure at the recommended clinical human dose (RHD) of EFV. Because of the potential risk of neural tube defects, EFV is not recommended for use in the first trimester of pregnancy. Avoid pregnancy while receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and for 12 weeks after discontinuation. Advise pregnant patients of the potential risk to a fetus.

Available data from the APR show no increase in the overall risk of major birth defects for EFV, FTC, or TDF compared with the background rate for major birth defects of 2.7% in a U.S. reference population of the Metropolitan Atlanta Congenital Defects Program (MACDP) (*see Data*).

The rate of miscarriage is not reported in the APR. The estimated background rate of miscarriage in clinically recognized pregnancies in the U.S. general population is 15- 20%. The background risk of major birth defects and miscarriage for the indicated population

is unknown. The APR uses the MACDP as the U.S. reference population for birth defects in the general population. The MACDP evaluates mothers and infants from a limited geographic area and does not include outcomes for births that occurred at less than 20 weeks' gestation.

In animal reproduction studies, no adverse developmental effects were observed when FTC and TDF were administered separately at doses/exposures ≥ 60 (FTC), ≥ 14 (TDF) and 2.7 (tenofovir) times those at the RHD of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet (*see Data*).

Data

Human Data

Efavirenz: There are retrospective postmarketing reports of findings consistent with neural tube defects, including meningomyelocele, all in infants of mothers exposed to EFV-containing regimens in the first trimester.

Based on prospective reports to the APR of 1,217 exposures to EFV-containing regimens during pregnancy resulting in live births (including over 1,023 live births exposed in the first trimester and 194 exposed in the second/third trimester), there was no increase in overall birth defects with EFV compared with the background birth defect rate of 2.7% in the U.S. reference population of the MACDP. The prevalence of birth defects in live births was 2.3% (95% CI: 1.5% to 3.5%) with first trimester exposure to EFV-containing regimens, and 1.5% (95% CI: 0.3% to 4.5%) with the second/third trimester exposure to EFV-containing regimens. One of these prospectively reported defects with first-trimester exposure was a neural tube defect. A single case of anophthalmia with first-trimester exposure to EFV has also been prospectively reported. This case also included severe oblique facial clefts and amniotic banding, which have a known association with anophthalmia.

Emtricitabine: Based on prospective reports from the APR of 4,005 exposures to FTC-containing regimens during pregnancy resulting in live births (including 2,785 exposed in the first trimester and 1,220 exposed in the second/third trimester), there was no increase in overall major birth defects with FTC compared with the background birth defect rate of 2.7% in the U.S. reference population of the MACDP. The prevalence of birth defects in live births was 2.4% (95% CI: 1.9% to 3.1%) with first trimester exposure to FTC-containing regimens and 2.3% (95% CI: 1.5% to 3.3%) with the second/third trimester exposure to FTC-containing regimens.

Tenofovir DF: Based on prospective reports from the APR of 5,105 exposures to TDF-containing regimens during pregnancy resulting in live births (including 3,535 exposed in the first trimester and 1,570 exposed in the second/third trimester), there was no increase in overall major birth defects with TDF compared with the background birth defect rate of 2.7% in the U.S. reference population of the MACDP. The prevalence of birth defects in live births was 2.3% (95% CI: 1.8% to 2.9%) with first trimester exposure to TDF-containing regimens, and 2.2% (95% CI: 1.6% to 3.1%) with the second/third trimester exposure to TDF-containing regimens.

Animal Data

Efavirenz: Effects of EFV on embryo-fetal development have been studied in three nonclinical species (cynomolgus monkeys, rats, and rabbits). In monkeys, EFV 60 mg/kg/day was administered to pregnant females throughout pregnancy (gestation

Days 20 through 150). The maternal systemic drug exposures (AUC) were 1.3 times the exposures at the RHD, with fetal umbilical venous drug concentrations approximately 0.7 times the maternal values. Three fetuses of 20 fetuses/infants had one or more malformations; there were no malformed fetuses or infants from placebo-treated mothers. The malformations that occurred in these three monkey fetuses included anencephaly and unilateral anophthalmia in one fetus, microphthalmia in a second, and cleft palate in the third. There was no NOAEL (no observable adverse effect level) established for this study because only one dosage was evaluated. In rats, EFV was administered either during organogenesis (gestation Days 7 to 18) or from gestation Day 7 through lactation Day 21 at 50, 100, or 200 mg/kg/day. Administration of 200 mg/kg/day in rats was associated with an increase in the incidence of early resorptions, and doses 100 mg/kg/day and greater were associated with early neonatal mortality. The AUC at the NOAEL (50 mg/kg/day) in this rat study was 0.1 times that in humans at the RHD. Drug concentrations in the milk on lactation Day 10 were approximately 8 times higher than those in maternal plasma. In pregnant rabbits, EFV was neither embryo lethal nor teratogenic when administered at doses of 25, 50, and 75 mg/kg/day over the period of organogenesis (gestation Days 6 through 18). The AUC at the NOAEL (75 mg/kg/day) in rabbits was 0.4 times that in humans at the RHD.

Emtricitabine: FTC was administered orally to pregnant mice (at 0, 250, 500, or 1,000 mg/kg/day), and rabbits (at 0, 100, 300, or 1,000 mg/kg/day) through organogenesis (on gestation days 6 through 15, and 7 through 19, respectively). No significant toxicological effects were observed in embryo-fetal toxicity studies performed with FTC in mice at exposures (AUC) approximately 60 times higher and in rabbits at approximately 120 times higher than human exposures at the RHD. In a pre/postnatal development study in mice, FTC was administered orally at doses up to 1000 mg/kg/day; no significant adverse effects directly related to drug were observed in the offspring exposed daily from before birth (in utero) through sexual maturity at daily exposures (AUC) of approximately 60 times higher than human exposures at the RHD.

Tenofovir DF: TDF was administered orally to pregnant rats (at 0, 50, 150, or 450 mg/kg/day) and rabbits (at 0, 30, 100, or 300 mg/kg/day) through organogenesis (on gestation days 7 through 17, and 6 through 18, respectively). No significant toxicological effects were observed in embryo-fetal toxicity studies performed with TDF in rats at doses up to 14 times the RHD based on body surface area comparisons and in rabbits at doses up to 19 times the RHD based on body surface area comparisons. In a pre/postnatal development study in rats, TDF was administered orally through lactation at doses up to 600 mg/kg/day; no adverse effects were observed in the offspring at tenofovir exposures of approximately 2.7 times higher than human exposures at the RHD.

8.2 Lactation

Risk Summary

The Centers for Disease Control and Prevention recommend that HIV-1 infected mothers not breastfeed their infants to avoid risking postnatal transmission of HIV-1.

Based on limited published data, EFV, FTC, and tenofovir have been shown to be present in human breast milk. It is not known if the components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet affect milk production or have effects on the breastfed child. Because of the potential for: (1) HIV transmission (in HIV-negative

infants); (2) developing viral resistance (in HIV-positive infants); and (3) adverse reactions in a breastfed infant similar to those seen in adults, instruct mothers not to breastfeed if they are receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet .

8.3 Females and Males of Reproductive Potential

Pregnancy Testing

Perform pregnancy testing in adults and adolescents of childbearing potential before initiation of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet because of potential risk of neural tube defects [*see Use in Specific Populations (8.1)*].

Contraception

Advise adults and adolescents of childbearing potential to use effective contraception during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and for 12 weeks after discontinuing Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet due to the long half-life of EFV, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Hormonal methods that contain progesterone may have decreased effectiveness. Always use barrier contraception in combination with other methods of contraception [*see Drug Interactions (7.1, 7.3)*].

8.4 Pediatric Use

The effectiveness and safety of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet as a complete regimen for the treatment of HIV-1 infection was established in pediatric patients with body weight greater than or equal to 40 kg [*see Dosage and Administration (2.2)*]. Use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in this age group is supported by adequate and well-controlled studies of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in adults with HIV-1 infection and data from pediatric studies of the individual components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet (EFV, FTC, and TDF).

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet should only be administered to pediatric patients with a body weight greater than or equal to 40 kg. Because Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is a fixed-dose combination tablet, the dose of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet cannot be adjusted for patients of lower weight [*see Warnings and Precautions (5.2, 5.9), Adverse Reactions (6.1), and Clinical Pharmacology (12.3)*].

8.5 Geriatric Use

Clinical trials of EFV, FTC, or TDF did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. In general, dose selection for elderly patients should be cautious, keeping in mind the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

8.6 Renal Impairment

Because Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is a fixed-dose combination, and cannot be dose adjusted, it is not recommended in patients with moderate or severe renal impairment (estimated creatinine clearance below 50 mL/min)

[see Dosage and Administration (2.3), Warnings and Precautions (5.7)].

8.7 Hepatic Impairment

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is not recommended for patients with moderate or severe hepatic impairment because there are insufficient data to determine an appropriate dose. Patients with mild hepatic impairment may be treated with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet at the approved dose. Because of the extensive cytochrome P450-mediated metabolism of EFV and limited clinical experience in patients with hepatic impairment, caution should be exercised in administering Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet to these patients [see Dosage and Administration (2.4), Warnings and Precautions (5.3), and Clinical Pharmacology (12.3)].

10 OVERDOSAGE

If overdose occurs, the patient should be monitored for evidence of toxicity, and standard supportive treatment applied as necessary. Administration of activated charcoal may be used to aid removal of unabsorbed EFV. Hemodialysis can remove both FTC and TDF (refer to detailed information below) but is unlikely to significantly remove EFV from the blood.

Efavirenz: Some patients accidentally taking 600 mg twice daily have reported increased nervous system symptoms. One patient experienced involuntary muscle contractions.

Emtricitabine: Hemodialysis treatment removes approximately 30% of the FTC dose over a 3-hour dialysis period starting within 1.5 hours of FTC dosing (blood flow rate of 400 mL/min and a dialysate flow rate of 600 mL/min). It is not known whether FTC can be removed by peritoneal dialysis.


Tenofovir DF: Tenofovir is efficiently removed by hemodialysis with an extraction coefficient of approximately 54%. Following a single 300 mg dose of TDF, a 4-hour hemodialysis session removed approximately 10% of the administered tenofovir dose.

11 DESCRIPTION

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is a fixed-dose combination tablet containing EFV, FTC, and TDF. EFV is a non-nucleoside reverse transcriptase inhibitor (NNRTI). FTC is a synthetic nucleoside analog of cytidine. TDF, which is converted *in vivo* to tenofovir, is an acyclic nucleoside phosphonate (nucleotide) analog of adenosine 5'-monophosphate.

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablets are for oral administration. Each tablet contains 600 mg of EFV, 200 mg of FTC, and 300 mg of TDF (equivalent to 245 mg of tenofovir disoproxil) as active ingredients. The tablets include the following inactive ingredients: Croscarmellose sodium, hydroxypropyl cellulose, hypromellose, isopropyl alcohol, microcrystalline cellulose, magnesium stearate, red iron oxide, sodium lauryl sulfate. The tablets are film-coated with a coating material opadry AMB 80W54485 pink containing polyvinyl alcohol-part, hydrolyzed, titanium dioxide, talc, lecithin (soya), xanthan gum, iron oxide yellow, iron oxide red and opadry AMB 80W56843 brown containing polyvinyl alcohol-part, hydrolyzed, titanium dioxide, talc, lecithin (soya), xanthan gum, iron oxide red.


Efavirenz: EFV is chemically described as (S)-6-chloro-4-(cyclopropylethynyl)-1,4-dihydro-4-(trifluoromethyl)-2H-3,1-benzoxazin-2-one. Its molecular formula is $C_{14}H_9ClF_3NO_2$ and its structural formula is:



Efavirenz is a white to slightly pink crystalline powder with a molecular mass of 315.68. It is practically insoluble in water (less than 10 $\mu\text{g}/\text{mL}$).


Emtricitabine: The chemical name of FTC is 5-fluoro-1-(2R,5S)-[2-(hydroxymethyl)-1,3-oxathiolan-5-yl]cytosine. FTC is the (-) enantiomer of a thio analog of cytidine, which differs from other cytidine analogs in that it has a fluorine in the 5-position.

It has a molecular formula of $C_8H_{10}FN_3O_3S$ and a molecular weight of 247.24. It has the following structural formula:



Emtricitabine is a white to off-white crystalline powder with a solubility of approximately 112 mg/mL in water at 25°C.

Tenofovir DF: TDF is a fumaric acid salt of the *bis*-isopropoxycarbonyloxymethyl ester derivative of tenofovir. The chemical name of TDF is 9-[(R)-2[[bis[[[(isopropoxycarbonyl)oxy]-methoxy] phosphinyl]methoxy]propyl]adenine fumarate (1:1). It has a molecular formula of $C_{19}H_{30}N_5O_{10}P \cdot C_4H_4O_4$ and a molecular weight of 635.52. It has the following structural formula:



TDF is a white to off-white crystalline powder with a solubility of 13.4 mg/mL in water at 25°C.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is a fixed-dose combination of antiviral drugs EFV, FTC, and TDF [see *Microbiology (12.4)*].

12.2 Pharmacodynamics

Cardiac Electrophysiology

Efavirenz: The effect of EFV on the QTc interval was evaluated in an open-label, positive and placebo-controlled, fixed single sequence 3-period, 3-treatment crossover QT study in 58 healthy subjects enriched for CYP2B6 polymorphisms. The mean C_{max} of EFV in subjects with CYP2B6 *6/*6 genotype following the administration of 600 mg daily dose for 14 days was 2.25-fold the mean C_{max} observed in subjects with CYP2B6 *1/*1 genotype. A positive relationship between EFV concentration and QTc prolongation was observed. Based on the concentration-QTc relationship, the mean QTc prolongation and its upper bound 90% confidence interval are 8.7 msec and 11.3 msec in subjects with CYP2B6*6/*6 genotype following the administration of 600 mg daily dose for 14 days [see *Warnings and Precautions (5.4)*].

12.3 Pharmacokinetics

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet: One Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is bioequivalent to one Sustiva tablet (600 mg) plus one EMTRIVA[®] capsule (200 mg) plus one VIREAD[®] tablet (300 mg) following single-dose administration to fasting healthy subjects (N=45).

Efavirenz: In HIV-1 infected subjects time-to-peak plasma concentrations were approximately 3-5 hours and steady-state plasma concentrations were reached in 6-10 days. In 35 HIV-1 infected subjects receiving EFV 600 mg once daily, steady-state C_{max} was $12.9 \pm 3.7 \mu\text{M}$ (mean \pm SD), C_{min} was $5.6 \pm 3.2 \mu\text{M}$, and AUC was $184 \pm 73 \mu\text{M}\cdot\text{hr}$. EFV is highly bound (approximately 99.5-99.75%) to human plasma proteins, predominantly albumin. Following administration of ¹⁴C-labeled EFV, 14-34% of the dose was recovered in the urine (mostly as metabolites) and 16-61% was recovered in feces (mostly as parent drug). *In vitro* studies suggest CYP3A and CYP2B6 are the major

isozymes responsible for EFV metabolism. EFV has been shown to induce CYP enzymes, resulting in induction of its own metabolism. EFV has a terminal half-life of 52-76 hours after single doses and 40-55 hours after multiple doses.

Emtricitabine: Following oral administration, FTC is rapidly absorbed, with peak plasma concentrations occurring at 1-2 hours postdose. Following multiple dose oral administration of FTC to 20 HIV-1 infected subjects, the steady-state plasma FTC C_{max} was $1.8 \pm 0.7 \mu\text{g/mL}$ (mean \pm SD) and the AUC over a 24-hour dosing interval was $10.0 \pm 3.1 \mu\text{g}\cdot\text{hr/mL}$. The mean steady-state plasma trough concentration at 24 hours postdose was $0.09 \mu\text{g/mL}$. The mean absolute bioavailability of FTC was 93%. Less than 4% of FTC binds to human plasma proteins *in vitro*, and the binding is independent of concentration over the range of 0.02-200 $\mu\text{g/mL}$. Following administration of radiolabelled FTC, approximately 86% is recovered in the urine and 13% is recovered as metabolites. The metabolites of FTC include 3'-sulfoxide diastereomers and their glucuronic acid conjugate. FTC is eliminated by a combination of glomerular filtration and active tubular secretion with a renal clearance in adults with normal renal function of $213 \pm 89 \text{ mL/min}$ (mean \pm SD). Following a single oral dose, the plasma FTC half-life is approximately 10 hours.

Tenofovir DF: Following oral administration of a single 300 mg dose of TDF to HIV-1 infected subjects in the fasted state, maximum serum concentrations (C_{max}) were achieved in $1.0 \pm 0.4 \text{ hrs}$ (mean \pm SD) and C_{max} and AUC values were $296 \pm 90 \text{ ng/mL}$ and $2287 \pm 685 \text{ ng}\cdot\text{hr/mL}$, respectively. The oral bioavailability of tenofovir from TDF in fasted subjects is approximately 25%. Less than 0.7% of tenofovir binds to human plasma proteins *in vitro*, and the binding is independent of concentration over the range of 0.01-25 $\mu\text{g/mL}$. Approximately 70-80% of the intravenous dose of tenofovir is recovered as unchanged drug in the urine. Tenofovir is eliminated by a combination of glomerular filtration and active tubular secretion, with a renal clearance in adults with normal renal function of $243 \pm 33 \text{ mL/min}$ (mean \pm SD). Following a single oral dose, the terminal elimination half-life of tenofovir is approximately 17 hours.

Effects of Food on Oral Absorption

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet has not been evaluated in the presence of food. Administration of EFV tablets with a high-fat meal increased the mean AUC and C_{max} of EFV by 28% and 79%, respectively, compared to administration in the fasted state. Compared to fasted administration, dosing of TDF and FTC in combination with either a high-fat meal or a light meal increased the mean AUC and C_{max} of tenofovir by 35% and 15%, respectively, without affecting FTC exposures [see *Dosage and Administration (2.2) and Patient Counseling Information (17)*].

Specific Populations

Race

Efavirenz: The pharmacokinetics of EFV in HIV-1 infected subjects appear to be similar among the racial groups studied.

Emtricitabine: No pharmacokinetic differences due to race have been identified following the administration of FTC.

Tenofovir DF: There were insufficient numbers from racial and ethnic groups other than Caucasian to adequately determine potential pharmacokinetic differences among these populations following the administration of TDF.

Gender

Efavirenz, Emtricitabine, and Tenofovir DF: EFV, FTC, and tenofovir pharmacokinetics are similar in male and female subjects.

Pediatric Patients

Efavirenz: In an open-label trial in NRTI-experienced pediatric subjects (mean age 8 years, range 3-16 years), the pharmacokinetics of EFV in pediatric subjects were similar to the pharmacokinetics in adults who received a 600 mg daily dose of EFV. Based on mean steady-state predicted population pharmacokinetic modeling in pediatric subjects weighing >40 kg receiving the 600 mg dose of EFV, C_{\max} was 6.57 $\mu\text{g/mL}$, C_{\min} was 2.82 $\mu\text{g/mL}$, and $\text{AUC}_{(0-24)}$ was 254.78 $\mu\text{M}\cdot\text{hr}$.

Emtricitabine: The pharmacokinetics of FTC at steady state were determined in 27 HIV-1-infected pediatric subjects 13 to 17 years of age receiving a daily dose of 6 mg/kg up to a maximum dose of 240 mg oral solution or a 200-mg capsule; 26 of 27 subjects in this age group received the 200-mg capsule. Mean \pm SD C_{\max} and AUC were 2.7 ± 0.9 $\mu\text{g/mL}$ and 12.6 ± 5.4 $\mu\text{g}\cdot\text{hr/mL}$, respectively. Exposures achieved in pediatric subjects 12 to less than 18 years of age were similar to those achieved in adults receiving a once daily dose of 200 mg.

Tenofovir DF: Steady-state pharmacokinetics of tenofovir were evaluated in 8 HIV-1 infected pediatric subjects (12 to less than 18 years). Mean \pm SD C_{\max} and AUC_{τ} are 0.38 ± 0.13 $\mu\text{g/mL}$ and 3.39 ± 1.22 $\mu\text{g}\cdot\text{hr/mL}$, respectively. Tenofovir exposure achieved in these pediatric subjects receiving oral daily doses of TDF 300 mg was similar to exposures achieved in adults receiving once-daily doses of TDF 300 mg.

Geriatric Patients

Pharmacokinetics of EFV, FTC, and tenofovir have not been fully evaluated in the elderly (65 years of age and older) [see *Use in Specific Populations (8.5)*].

Patients with Impaired Renal Function

Efavirenz: The pharmacokinetics of EFV have not been studied in subjects with renal insufficiency; however, less than 1% of EFV is excreted unchanged in the urine, so the impact of renal impairment on EFV elimination should be minimal.

Emtricitabine and Tenofovir DF: The pharmacokinetics of FTC and TDF are altered in subjects with renal impairment. In subjects with creatinine clearance below 50 mL/min, C_{\max} and $\text{AUC}_{0-\infty}$ of FTC and tenofovir were increased [see *Warnings and Precautions (5.7)*].

Patients with Hepatic Impairment

Efavirenz: A multiple-dose trial showed no significant effect on EFV pharmacokinetics in subjects with mild hepatic impairment (Child-Pugh Class A) compared with controls. There were insufficient data to determine whether moderate or severe hepatic impairment (Child-Pugh Class B or C) affects EFV pharmacokinetics [see *Warnings and Precautions (5.3)* and *Use in Specific Populations (8.7)*].

Emtricitabine: The pharmacokinetics of FTC have not been studied in subjects with hepatic impairment; however, FTC is not significantly metabolized by liver enzymes, so the impact of liver impairment should be limited.

Tenofovir DF: The pharmacokinetics of tenofovir following a 300 mg dose of TDF have

been studied in non-HIV infected subjects with moderate to severe hepatic impairment. There were no substantial alterations in tenofovir pharmacokinetics in subjects with hepatic impairment compared with unimpaired subjects.

Assessment of Drug Interactions

The drug interaction trials described were conducted with either efavirenz, emtricitabine, or tenofovir DF or the components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet (EFV, FTC, or TDF) as individual agents.

Efavirenz: The steady-state pharmacokinetics of EFV and tenofovir were unaffected when EFV and TDF were administered together versus each agent dosed alone. Specific drug interaction trials have not been performed with EFV and NRTIs other than tenofovir, lamivudine, and zidovudine. Clinically significant interactions would not be expected based on NRTIs elimination pathways.

Efavirenz has been shown *in vivo* to cause hepatic enzyme induction, thus increasing the biotransformation of some drugs metabolized by CYP3A and CYP2B6. *In vitro* studies have shown that EFV inhibited CYP isozymes 2C9 and 2C19 with K_i values (8.5-17 μ M) in the range of observed EFV plasma concentrations. In *in vitro* studies, EFV did not inhibit CYP2E1 and inhibited CYP2D6 and CYP1A2 (K_i values 82-160 μ M) only at concentrations well above those achieved clinically. Coadministration of EFV with drugs primarily metabolized by CYP2C9, CYP2C19, CYP3A or CYP2B6 isozymes may result in altered plasma concentrations of the coadministered drug. Drugs which induce CYP3A and CYP2B6 activity would be expected to increase the clearance of EFV resulting in lowered plasma concentrations.

Drug interaction trials were performed with EFV and other drugs likely to be coadministered or drugs commonly used as probes for pharmacokinetic interaction. There was no clinically significant interaction observed between EFV and zidovudine, lamivudine, azithromycin, fluconazole, lorazepam, cetirizine, or paroxetine. Single doses of famotidine or an aluminum and magnesium antacid with simethicone had no effects on EFV exposures. The effects of coadministration of EFV on C_{max} , AUC, and C_{min} are summarized in Table 4 (effect of other drugs on EFV) and Table 5 (effect of EFV on other drugs) see [Drug Interactions (7)].

Table 4: Drug Interactions: Changes in Pharmacokinetic Parameters for EFV in the Presence of the Coadministered Drug

				Mean % Change of EFV Pharmacokinetic Parameters^a (90% CI)		
Coadministered Drug	Dose of Coadministered Drug (mg)	EFV Dose (mg)	N	C_{max}	AUC	C_{min}
Lopinavir/ritonavir	400/100 mg q12h x 9 days	600 mg qd x 9 days	11, 12 ^b	↔	↓ 16 (↓ 38 to ↑ 15)	↓ 16 (↓ 42 to ↑ 20)
Nelfinavir	750 mg q8h x 7 days	600 mg qd x 7 days	10	↓ 12 (↓ 32 to ↑ 13) ^c	↓ 12 (↓ 35 to ↑)	↓ 21 (↓ 53 to ↑)

					18) ^c	33)
Ritonavir	500 mg q 12h x 8 days	600 mg qd x 10 days	9	↑ 14 (↑ 4 to ↑ 26)	↑ 21 (↑ 10 to ↑ 34)	↑ 25 (↑ 7 to ↑ 46) ^c
Boceprevir	800 mg tid x 6 days	600 mg qd x 16 days	NA	↑ 11 (↑ 2 to ↑ 20)	↑ 20 (↑ 15 to ↑ 26)	NA
Rifabutin	300 mg qd x 14 days	600 mg qd x 14 days	11	↔	↔	↓ 12 (↓ 24 to ↑ 1)
Rifampin	600 mg x 7 days	600 mg qd x 7 days	12	↓ 20 (↓ 11 to ↓ 28)	↓ 26 (↓ 15 to ↓ 36)	↓ 32 (↓ 15 to ↓ 46)
Artemether/ lumefantrine	Artemether 20 mg/lumefantrine 120 mg tablets (6 4-tablet doses over 3 days)	600 mg qd x 26 days	12	↔	↓ 17	NA
Simvastatin	40 mg qd x 4 days	600 mg qd x 15 days	14	↓ 12 (↓ 28 to ↑ 8)	↔	↓ 12 (↓ 25 to ↑ 3)
Carbamazepine	200 mg qd x 3 days, 200 mg bid x 3 days, then 400 mg qd x 15 days	600 mg qd x 35 days	14	↓ 21 (↓ 15 to ↓ 26)	↓ 36 (↓ 32 to ↓ 40)	↓ 47 (↓ 41 to ↓ 53)
Diltiazem	240 mg x 14 days	600 mg qd x 28 days	12	↑ 16 (↑ 6 to ↑ 26)	↑ 11 (↑ 5 to ↑ 18)	↑ 13 (↑ 1 to ↑ 26)
Voriconazole	400 mg po q12h x 1 day then 200 mg po q12h x 8 days	400 mg qd x 9 days	NA	↑ 38 ^d	↑ 44 ^d	NA
	300 mg po q12h days 2-7	300 mg qd x 7 days	NA	↓ 14 ^e (↓ 7 to ↓ 21)	↔ ^e	NA
	400 mg po q12h days 2-7	300 mg qd x 7 days	NA	↔ ^e	↑ 17 ^e (↑ 6 to ↑ 29)	NA

NA = not available

a. Increase = ↑; Decrease = ↓; No Effect = ↔

b. Parallel-group design; N for EFV + lopinavir/ritonavir, N for EFV alone.

c. 95% CI

d. 90% CI not available

e. Relative to steady-state administration of EFV (600 mg once daily for 9 days).

No effect on the pharmacokinetic parameters of EFV was observed with the following coadministered drugs: indinavir, saquinavir soft gelatin capsule, simeprevir, ledipasvir/sofosbuvir, sofosbuvir, clarithromycin, itraconazole, atorvastatin, pravastatin, or sertraline.

Table 5 Drug Interactions: Changes in Pharmacokinetic Parameters for Coadministered Drug in the Presence of EFV

				Mean % Change of Coadministered Drug Pharmacokinetic Parameters^a (90% CI)		
Coadministered Drug	Dose of Coadministered Drug (mg)	EFV Dose (mg)	N	C_{max}	AUC	C_{min}
Atazanavir	400 mg qd with a light meal d 1-20	600 mg qd with a light meal d 7-20	27	↓ 59 (↓ 49 to ↓ 67)	↓ 74 (↓ 68 to ↓ 78)	↓ 93 (↓ 90 to ↓ 95)
	400 mg qd d 1-6, then 300 mg qd d 7-20 with ritonavir 100 mg qd and a light meal	600 mg qd 2 h after atazanavir and ritonavir d 7-20	13	↑ 14 ^b (↓ 17 to ↑ 58)	↑ 39 ^b (↑ 2 to ↑ 88)	↑ 48 ^b (↑ 24 to ↑ 76)
	300 mg qd/ritonavir 100 mg qd d 1-10 (pm), then 400 mg qd/ritonavir 100 mg qd d 11-24 (pm) (simultaneous with EFV)	600 mg qd with a light snack d 11-24 (pm)	14	↑ 17 (↑ 8 to ↑ 27)	↔	↓ 42 (↓ 31 to ↓ 51)
Indinavir	1000 mg q8h x 10 days	600 mg qd x 10 days	20			
	After morning dose			↔ ^c	↓ 33 ^c (↓ 26 to ↓ 39)	↓ 39 ^c (↓ 24 to ↓ 51)
	After afternoon dose			↔ ^c	↓ 37 ^c (↓ 26 to ↓ 46)	↓ 52 ^c (↓ 47 to ↓ 57)
	After evening dose				↓ 29 ^c	↓ 57 ^c

				(↓ 11 to ↓ 43)	(↓ 37 to ↓ 54)	(↓ 50 to ↓ 63)
Lopinavir/ritonavir	400/100 mg q12h x 9 days	600 mg qd x 9 days	11, 7 ^d	↔ ^e	↓ 19 ^e (↓ 36 to ↑ 3)	↓ 39 ^e (↓ 3 to ↓ 62)
Nelfinavir Metabolite AG-1402	750 mg q8h x 7 days	600 mg qd x 7 days	10	↑ 21 (↑ 10 to ↑ 33) ↓ 40 (↓ 30 to ↓ 48)	↑ 20 (↑ 8 to ↑ 34) ↓ 37 (↓ 25 to ↓ 48)	↔ ↓ 43 (↓ 21 to ↓ 59)
Ritonavir	500 mg q12h x 8 days	600 mg qd x 10 days	11			
	After AM dose			↑ 24 (↑ 12 to ↑ 38)	↑ 18 (↑ 6 to ↑ 33)	↑ 42 (↑ 9 to ↑ 86) ^f
	After PM dose			↔	↔	↑ 24 (↑ 3 to ↑ 50) ^f
Saquinavir SGC ^g	1200 mg q8h x 10 days	600 mg qd x 10 days	12	↓ 50 (↓ 28 to ↓ 66)	↓ 62 (↓ 45 to ↓ 74)	↓ 56 (↓ 16 to ↓ 77) ^f
Maraviroc	100 mg bid	600 mg qd	12	↓ 51 (↓ 37 to ↓ 62)	↓ 45 (↓ 38 to ↓ 51)	↓ 45 (↓ 28 to ↓ 57)
Raltegravir	400 mg single dose	600 mg qd	9	↓ 36 (↓ 2 to ↓ 59)	↓ 36 (↓ 20 to ↓ 48)	↓ 21 (↓ 51 to ↑ 28)
Boceprevir	800 mg tid x 6 days	600 mg qd x 16 days	NA	↓ 8 (↓ 22 to ↑ 8)	↓ 19 (↓ 11 to ↓ 25)	↓ 44 (↓ 26 to ↓ 58)
Simeprevir	150 mg qd x 14 days	600 mg qd x 14 days	23	↓ 51 (↓ 46 to ↓ 56)	↓ 71 (↓ 67 to ↓ 74)	↓ 91 (↓ 88 to ↓ 92)
Ledipasvir/ Sofosbuvir ^k Ledipasvir Sofosbuvir GS-331007 ^l	90/400 mg qd x 14 days	600 mg qd x 14 days	15	↓ 34 (↓ 25 to ↓ 41) ↔	↓ 34 (↓ 25 to ↓ 41) ↔	↓ 34 (↓ 24 to ↓ 43) NA

				↔	↔	↔
Sofosbuvir ^m GS-331007 ^l	400 mg qd single dose	600 mg qd x14 days	16	↓ 19 (↓ 40 to ↑ 10) ↓ 23 (↓ 16 to ↓ 30)	↔ ↓ 16 (↓ 24 to ↓ 8)	NA NA
Sofosbuvir/Velpatasvir ⁿ Sofosbuvir GS-331007 ^l Velpatasvir	400/100 mg qd × 14 days	600 mg qd × 14 days	14	↑ 38 (↑ 14 to ↑ 67) ↓ 14 (↓ 20 to ↓ 7) ↓ 47 (↓ 57 to ↓ 36)	↔ ↔ ↓ 53 (↓ 61 to ↓ 43)	NA ↔ ↓ 57 (↓ 64 to ↓ 48)
Clarithromycin 14-OH metabolite	500 mg q12h × 7 days	400 mg qd × 7 days	11	↓ 26 (↓ 15 to ↓ 35) ↑ 49 (↑ 32 to ↑ 69)	↓ 39 (↓ 30 to ↓ 46) ↑ 34 (↑ 18 to ↑ 53)	↓ 53 (↓ 42 to ↓ 63) ↑ 26 (↑ 9 to ↑ 45)
Itraconazole Hydroxy-itraconazole	200 mg q 12 h × 28 days	600 mg qd × 14 days	18	↓ 37 (↓ 20 to ↓ 51) ↓ 35 (↓ 12 to ↓ 52)	↓ 39 (↓ 21 to ↓ 53) ↓ 37 (↓ 14 to ↓ 55)	↓ 44 (↓ 27 to ↓ 58) ↓ 43 (↓ 18 to ↓ 60)
Posaconazole	400 mg (oral suspension) bid × 10 and 20 days	400 mg qd × 10 and 20 days	11	↓ 45 (↓ 34 to ↓ 53)	↓ 50 (↓ 40 to ↓ 57)	NA
Rifabutin	300 mg qd × 14 days	600 mg qd × 14 days	9	↓ 32 (↓ 15 to ↓ 46)	↓ 38 (↓ 28 to ↓ 47)	↓ 45 (↓ 31 to ↓ 56)
Artemether/ lumefantrine Artemether dihydroartemisinin lumefantrine	Artemether 20 mg/lumefantrine 120 mg tablets (6 4-tablet doses over 3 days)	600 mg qd × 26 days	12	↓ 21 ↓ 38 ↔	↓ 51 ↓ 46 ↓ 21	NA NA NA

Atorvastatin Total active (including metabolites)	10 mg qd x 4 days	600 mg qd x 15 days	14	↓ 14 (↓ 1 to ↓ 26) ↓ 15 (↓ 2 to ↓ 26)	↓ 43 (↓ 34 to ↓ 50) ↓ 32 (↓ 21 to ↓ 41)	↓ 69 (↓ 49 to ↓ 81) ↓ 48 (↓ 23 to ↓ 64)
Pravastatin	40 mg qd x 4 days	600 mg qd x 15 days	13	↓ 32 (↓ 59 to ↑ 12)	↓ 44 (↓ 26 to ↓ 57)	↓ 19 (↓ 0 to ↓ 35)
Simvastatin Total active (including metabolites)	40 mg qd x 4 days	600 mg qd x 15 days	14	↓ 72 (↓ 63 to ↓ 79) ↓ 68 (↓ 55 to ↓ 78)	↓ 68 (↓ 62 to ↓ 73) ↓ 60 (↓ 52 to ↓ 68)	↓ 45 (↓ 20 to ↓ 62) NA ^h
Carbamazepine Epoxide metabolite	200 mg qd x 3 days, 200 mg bid x 3 days, then 400 mg qd x 29 days	600 mg qd x 14 days	12	↓ 20 (↓ 15 to ↓ 24) ↔	↓ 27 (↓ 20 to ↓ 33) ↔	↓ 35 (↓ 24 to ↓ 44) ↓ 13 (↓ 30 to ↑ 7)
Diltiazem Desacetyl diltiazem N-monodesmethyl diltiazem	240 mg x 21 days	600 mg qd x 14 days	13	↓ 60 (↓ 50 to ↓ 68) ↓ 64 (↓ 57 to ↓ 69) ↓ 28 (↓ 7 to ↓ 44)	↓ 69 (↓ 55 to ↓ 79) ↓ 75 (↓ 59 to ↓ 84) ↓ 37 (↓ 17 to ↓ 52)	↓ 63 (↓ 44 to ↓ 75) ↓ 62 (↓ 44 to ↓ 75) ↓ 37 (↓ 17 to ↓ 52)
Ethinyl estradiol/ Norgestimate Ethinyl estradiol Norelgestromin Levonorgestrel	0.035 mg/0.25 mg x 14 days	600 mg qd x 14 days	21	↔	↔	↔
			21	↓ 46 (↓ 39 to ↓ 52)	↓ 64 (↓ 62 to ↓ 67)	↓ 82 (↓ 79 to ↓ 85)
			6	↓ 80 (↓ 77 to ↓ 83)	↓ 83 (↓ 79 to ↓ 87)	↓ 86 (↓ 80 to ↓ 90)
Methadone	Stable	600 mg	11	↓ 45	↓ 52	NA

	maintenance 35-100 mg daily	qd x 14-21 days		(↓ 25 to ↓ 59)	(↓ 33 to ↓ 66)	
Bupropion Hydroxybupropion	150 mg single dose (sustained- release)	600 mg qd x 14 days	13	↓ 34 (↓ 21 to ↓ 47) ↑ 50 (↑ 20 to ↑ 80)	↓ 55 (↓ 48 to ↓ 62) ↔	NA NA
Sertraline	50 mg qd x 14 days	600 mg qd x 14 days	13	↓ 29 (↓ 15 to ↓ 40)	↓ 39 (↓ 27 to ↓ 50)	↓ 46 (↓ 31 to ↓ 58)
Voriconazole	400 mg po q12h x 1 day then 200 mg po q12h x 8 days	400 mg qd x 9 days	NA	↓ 61 ⁱ	↓ 77 ⁱ	NA
	300 mg po q 12 h days 2-7	300 mg qd x 7 days	NA	↓ 36 ^j (↓ 21 to ↓ 49)	↓ 55 ^j (↓ 45 to ↓ 62)	NA
	400 mg po q12h days 2-7	300 mg qd x 7 days	NA	↑ 23 ^j (↓ 1 to ↑ 53)	↓ 7 ^j (↓ 23 to ↑ 13)	NA

NA= not available

- a. Increase = ↑; Decrease = ↓; No Effect = ↔
- b. Compared with atazanavir 400 mg qd alone.
- c. Comparator dose of indinavir was 800 mg q8h x 10 days.
- d. Parallel-group design; N for EFV + lopinavir/ritonavir, N for lopinavir/ritonavir alone.
- e. Values are for lopinavir. The pharmacokinetics of ritonavir 100 mg q12h are unaffected by concurrent EFV.
- f. 95% CI
- g. Soft Gelatin Capsule
- h. Not available because of insufficient data.
- i. 90% CI not available.
- j. Relative to steady-state administration of voriconazole (400 mg for 1 day, then 200 mg po q12h for 2 days).
- k. Study conducted with efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet coadministered with HARVONI.
- l. The predominant circulating nucleoside metabolite of sofosbuvir.
- m. Study conducted with efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet coadministered with SOVALDI® (sofosbuvir).
- n. Study conducted with efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet coadministered with EPCLUSA.

Emtricitabine and Tenofovir DF: The steady-state pharmacokinetics of FTC and tenofovir were unaffected when FTC and TDF were administered together versus each agent dosed alone.

In vitro and clinical pharmacokinetic drug-drug interaction studies have shown that the potential for CYP mediated interactions involving FTC and tenofovir with other medicinal products is low.

TDF is a substrate of P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP) transporters. When TDF is coadministered with an inhibitor of these transporters, an increase in absorption may be observed.

No clinically significant drug interactions have been observed between FTC and famciclovir, indinavir, sofosbuvir/velpatasvir, stavudine, TDF, and zidovudine. Similarly, no clinically significant drug interactions have been observed between TDF and abacavir, EFV, FTC, entecavir, indinavir, lamivudine, lopinavir/ritonavir, methadone, nelfinavir, oral contraceptives, ribavirin, saquinavir/ritonavir, sofosbuvir, or tacrolimus in trials conducted in healthy volunteers.

Following multiple dosing to HIV-negative subjects receiving either chronic methadone maintenance therapy, oral contraceptives, or single doses of ribavirin, steady-state tenofovir pharmacokinetics were similar to those observed in previous trials, indicating a lack of clinically significant drug interactions between these agents and TDF.

The effects of coadministered drugs on the C_{max} , AUC, and C_{min} of tenofovir are shown in Table 6. The effects of coadministration of TDF on C_{max} , AUC, and C_{min} of coadministered drugs are shown in Table 7.

Table 6: Drug Interactions: Changes in Pharmacokinetic Parameters for Tenofovir in the Presence of the Coadministered Drug^{a,b}

Coadministered Drug	Dose of Coadministered Drug (mg)	N	Mean% Change of Tenofovir Pharmacokinetic Parameters ^c (90% CI)		
			C_{max}	AUC	C_{min}
Atazanavir ^d	400 once daily x 14 days	33	↑ 14 (↑ 8 to ↑ 20)	↑ 24 (↑ 21 to ↑ 28)	↑ 22 (↑ 15 to ↑ 30)
Atazanavir/ritonavir ^d	300/100 once daily	12	↑ 34 (↑ 20 to ↑ 51)	↑ 37 (↑ 30 to ↑ 45)	↑ 29 (↑ 21 to ↑ 36)
Darunavir/ritonavir ^e	300/100 twice daily	12	↑ 24 (↑ 8 to ↑ 42)	↑ 22 (↑ 10 to ↑ 35)	↑ 37 (↑ 19 to ↑ 57)
Didanosine ^f	250 or 400 once daily x 7 days	14	↔	↔	↔
Ledipasvir/sofosbuvir	90/400 once daily	15	↑ 79 (↑ 56 to ↑ 104)	↑ 98 (↑ 77 to ↑ 123)	↑ 163 (↑ 132 to ↑ 197)
Lopinavir/ritonavir	400/100 twice daily x 14 days	24	↔	↑ 32 (↑ 25 to ↑ 38)	↑ 51 (↑ 37 to ↑ 66)
Sofosbuvir	400 once daily	16	↑ 25 (↑ 8 to ↑ 45)	↔	↔
Sofosbuvir/velpatasvir	400/100 once daily	15	↑ 77 (↑ 53 to ↑ 101)	↑ 81 (↑ 68 to ↑ 94)	↑ 121 (↑ 100 to ↑ 142)

			104)	94)	143)
Tipranavir/ ritonavir ^g	500/100 twice daily	22	↓ 23 (↓ 32 to ↓ 13)	↓ 2 (↓ 9 to ↑ 5)	↑ 7 (↓ 2 to ↑ 17)
	750/200 twice daily (23 doses)	20	↓ 38 (↓ 46 to ↓ 29)	↑ 2 (↓ 6 to ↑ 10)	↑ 14 (↑ 1 to ↑ 27)

a. All interaction trials conducted in healthy volunteers.

b. Subjects received TDF 300 mg once daily.

c. Increase = ↑; Decrease = ↓; No Effect = ↔

d. Reyataz Prescribing Information.

e. Prezista Prescribing Information.

f. Subjects received didanosine buffered tablets.

g. Aptivus Prescribing Information.

Table 7: Drug Interactions: Changes in Pharmacokinetic Parameters for Coadministered Drug in the Presence of TDF^{a,b}

Coadministered Drug	Dose of Coadministered Drug (mg)	N	Mean % Change of Co-administered Drug Pharmacokinetic Parameters ^c (90% CI)		
			C _{max}	AUC	C _{min}
Atazanavir ^d	400 once daily x 14 days	34	↓ 21 (↓ 27 to ↓ 14)	↓ 25 (↓ 30 to ↓ 19)	↓ 40 (↓ 48 to ↓ 32)
	Atazanavir/ritonavir 300/100 once daily x 42 days	10	↓ 28 (↓ 50 to ↑ 5)	↓ 25 ^e (↓ 42 to ↓ 3)	↓ 23 ^e (↓ 46 to ↑ 10)
Darunavir ^f	Darunavir/ritonavir 300/100 once daily	12	↑ 16 (↓ 6 to ↑ 42)	↑ 21 (↓ 5 to ↑ 54)	↑ 24 (↓ 10 to ↑ 69)
Didanosine ^g	250 once, simultaneously with TDF and a light meal ^h	33	↓ 20 ⁱ (↓ 32 to ↓ 7)	↔ ⁱ	NA
Lopinavir Ritonavir	Lopinavir/ritonavir 400/100 twice daily x 14 days	24	↔	↔	↔
	Lopinavir/ritonavir 400/100 twice daily x 14 days	24	↔	↔	↔
Tipranavir ^j	Tipranavir/ritonavir 500/100 twice daily	22	↓ 17 (↓ 26 to ↓ 6)	↓ 18 (↓ 25 to ↓ 9)	↓ 21 (↓ 30 to ↓ 10)
	Tipranavir/ritonavir 750/200 twice daily (23 doses)	20	↓ 11 (↓ 16 to ↓ 4)	↓ 9 (↓ 15 to ↓ 3)	↓ 12 (↓ 22 to 0)

a. All interaction trials conducted in healthy volunteers.

b. Subjects received TDF 300 mg once daily.

- c. Increase = ↑; Decrease = ↓; No Effect = ↔
- d. Reyataz Prescribing Information.
- e. In HIV-infected patients, addition of TDF to atazanavir 300 mg plus ritonavir 100 mg, resulted in AUC and C_{\min} values of atazanavir that were 2.3- and 4-fold higher than the respective values observed for atazanavir 400 mg when given alone.
- f. Prezista Prescribing Information.
- g. Videx EC Prescribing Information. Subjects received didanosine enteric-coated capsules.
- h. 373 kcal, 8.2 g fat.
- i. Compared with didanosine (enteric-coated) 400 mg administered alone under fasting conditions.
- j. Aptivus Prescribing Information.

12.4 Microbiology

Mechanism of Action

Efavirenz: EFV is a non-nucleoside reverse transcriptase (RT) inhibitor of HIV-1. Efavirenz activity is mediated predominantly by noncompetitive inhibition of HIV-1 reverse transcriptase. HIV-2 RT and human cellular DNA polymerases α , β , γ , and δ are not inhibited by EFV.

Emtricitabine: Emtricitabine, a synthetic nucleoside analog of cytidine, is phosphorylated by cellular enzymes to form FTC 5'-triphosphate. Emtricitabine 5'-triphosphate inhibits the activity of the HIV-1 RT by competing with the natural substrate deoxycytidine 5'-triphosphate and by being incorporated into nascent viral DNA which results in chain termination. Emtricitabine 5'-triphosphate is a weak inhibitor of mammalian DNA polymerases α , β , ϵ , and mitochondrial DNA polymerase γ .

Tenofovir DF: TDF is an acyclic nucleoside phosphonate diester analog of adenosine monophosphate. TDF requires initial diester hydrolysis for conversion to tenofovir and subsequent phosphorylations by cellular enzymes to form tenofovir diphosphate. Tenofovir diphosphate inhibits the activity of HIV-1 RT by competing with the natural substrate deoxyadenosine 5'-triphosphate and, after incorporation into DNA, by DNA chain termination. Tenofovir diphosphate is a weak inhibitor of mammalian DNA polymerases α , β , and mitochondrial DNA polymerase γ .

Antiviral Activity

Efavirenz, Emtricitabine, and Tenofovir DF: In combination studies evaluating the antiviral activity in cell culture of FTC and EFV together, EFV and tenofovir together, and FTC and tenofovir together, additive to synergistic antiviral effects were observed.

Efavirenz: The concentration of EFV inhibiting replication of wild-type laboratory adapted strains and clinical isolates in cell culture by 90-95% (EC_{90-95}) ranged from 1.7-25 nM in lymphoblastoid cell lines, peripheral blood mononuclear cells, and macrophage/monocyte cultures. Efavirenz demonstrated additive antiviral activity against HIV-1 in cell culture when combined with non-nucleoside reverse transcriptase inhibitors (NNRTIs) (delavirdine and nevirapine), nucleoside reverse transcriptase inhibitors (NRTIs) (abacavir, didanosine, lamivudine, stavudine, zalcitabine, and zidovudine), protease inhibitors (PIs) (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir), and the fusion inhibitor enfuvirtide. Efavirenz demonstrated additive to antagonistic antiviral activity in cell culture with atazanavir. Efavirenz demonstrated antiviral activity against clade B and most non-clade B isolates (subtypes A, AE, AG, C, D, F, G, J, and N), but had reduced antiviral activity against group O viruses. Efavirenz is not active against HIV-2.

Emtricitabine: The antiviral activity in cell culture of FTC against laboratory and clinical isolates of HIV-1 was assessed in lymphoblastoid cell lines, the MAGI-CCR5 cell line, and peripheral blood mononuclear cells. The 50% effective concentration (EC₅₀) values for FTC were in the range of 0.0013-0.64 µM (0.0003-0.158 µg/mL). In drug combination studies of FTC with NRTIs (abacavir, lamivudine, stavudine, zalcitabine, and zidovudine), NNRTIs (delavirdine, EFV, and nevirapine), and PIs (amprenavir, nelfinavir, ritonavir, and saquinavir), additive to synergistic effects were observed. Emtricitabine displayed antiviral activity in cell culture against HIV-1 clades A, B, C, D, E, F, and G (EC₅₀ values ranged from 0.007-0.075 µM) and showed strain-specific activity against HIV-2 (EC₅₀ values ranged from 0.007-1.5 µM).

Tenofovir DF: The antiviral activity in cell culture of tenofovir against laboratory and clinical isolates of HIV-1 was assessed in lymphoblastoid cell lines, primary monocyte/macrophage cells and peripheral blood lymphocytes. The EC₅₀ values for tenofovir were in the range of 0.04-8.5 µM. In drug combination studies of tenofovir with NRTIs (abacavir, didanosine, lamivudine, stavudine, zalcitabine, and zidovudine), NNRTIs (delavirdine, EFV, and nevirapine), and PIs (amprenavir, indinavir, nelfinavir, ritonavir, and saquinavir), additive to synergistic effects were observed. Tenofovir displayed antiviral activity in cell culture against HIV-1 clades A, B, C, D, E, F, G, and O (EC₅₀ values ranged from 0.5-2.2 µM) and showed strain-specific activity against HIV-2 (EC₅₀ values ranged from 1.6 µM-5.5 µM).

Resistance

EFV, FTC, and TDF: HIV-1 isolates with reduced susceptibility to the combination of FTC and tenofovir have been selected in cell culture and in clinical trials. Genotypic analysis of these isolates identified the M184V/I and/or K65R amino acid substitutions in the viral RT. In addition, a K70E substitution in HIV-1 reverse transcriptase has been selected by tenofovir and results in reduced susceptibility to tenofovir.

In a clinical trial of treatment-naïve subjects [*Study 934, see Clinical Studies (14)*] resistance analysis was performed on HIV-1 isolates from all confirmed virologic failure subjects with greater than 400 copies/mL of HIV-1 RNA at Week 144 or early discontinuations. Genotypic resistance to EFV, predominantly the K103N substitution, was the most common form of resistance that developed. Resistance to EFV occurred in 13/19 analyzed subjects in the FTC + TDF group and in 21/29 analyzed subjects in the zidovudine/lamivudine fixed-dose combination group. The M184V amino acid substitution, associated with resistance to FTC and lamivudine, was observed in 2/19 analyzed subject isolates in the FTC + TDF group and in 10/29 analyzed subject isolates in the zidovudine/lamivudine group. Through 144 weeks of Study 934, no subjects developed a detectable K65R substitution in their HIV-1 as analyzed through standard genotypic analysis.

In a clinical trial of treatment-naïve subjects, isolates from 8/47 (17%) analyzed subjects receiving TDF developed the K65R substitution through 144 weeks of therapy; 7 of these occurred in the first 48 weeks of treatment and one at Week 96. In treatment experienced subjects, 14/304 (5%) of TDF treated subjects with virologic failure through Week 96 showed greater than 1.4-fold (median 2.7) reduced susceptibility to tenofovir. Genotypic analysis of the resistant isolates showed a substitution in the HIV-1 RT gene resulting in the K65R amino acid substitution.

Efavirenz: Clinical isolates with reduced susceptibility in cell culture to EFV have been obtained. The most frequently observed amino acid substitution in clinical trials with EFV

is K103N (54%). One or more RT substitutions at amino acid positions 98, 100, 101, 103, 106, 108, 188, 190, 225, 227, and 230 were observed in subjects failing treatment with EFV in combination with other antiretrovirals. Other resistance substitutions observed to emerge commonly included L100I (7%), K101E/Q/R (14%), V108I (11%), G190S/T/A (7%), P225H (18%), and M230I/L (11%).

HIV-1 isolates with reduced susceptibility to EFV (greater than 380-fold increase in EC₉₀ value) emerged rapidly under selection in cell culture. Genotypic characterization of these viruses identified substitutions resulting in single amino acid substitutions L100I or V179D, double substitutions L100I/V108I, and triple substitutions L100I/V179D/Y181C in RT.

Emtricitabine: Emtricitabine-resistant isolates of HIV-1 have been selected in cell culture and in clinical trials. Genotypic analysis of these isolates showed that the reduced susceptibility to FTC was associated with a substitution in the HIV-1 RT gene at codon 184 which resulted in an amino acid substitution of methionine by valine or isoleucine (M184V/I).

Tenofovir DF: HIV-1 isolates with reduced susceptibility to tenofovir have been selected in cell culture. These viruses expressed a K65R substitution in RT and showed a 2- to 4-fold reduction in susceptibility to tenofovir.

Cross Resistance

Efavirenz, Emtricitabine, and Tenofovir DF: Cross resistance has been recognized among NNRTIs. Cross resistance has also been recognized among certain NRTIs. The M184V/I and/or K65R substitutions selected in cell culture by the combination of FTC and tenofovir are also observed in some HIV-1 isolates from subjects failing treatment with tenofovir in combination with either lamivudine or FTC, and either abacavir or didanosine. Therefore, cross resistance among these drugs may occur in patients whose virus harbors either or both of these amino acid substitutions.

Efavirenz: Clinical isolates previously characterized as EFV resistant were also phenotypically resistant in cell culture to delavirdine and nevirapine compared to baseline. Delavirdine- and/or nevirapine-resistant clinical viral isolates with NNRTI resistance-associated substitutions (A98G, L100I, K101E/P, K103N/S, V106A, Y181X, Y188X, G190X, P225H, F227L, or M230L) showed reduced susceptibility to EFV in cell culture. Greater than 90% of NRTI-resistant isolates tested in cell culture retained susceptibility to EFV.

Emtricitabine: Emtricitabine-resistant isolates (M184V/I) were cross resistant to lamivudine but retained susceptibility in cell culture to didanosine, stavudine, tenofovir, zidovudine, and NNRTIs (delavirdine, EFV, and nevirapine). HIV-1 isolates containing the K65R substitution, selected *in vivo* by abacavir, didanosine, and tenofovir, demonstrated reduced susceptibility to inhibition by FTC. Viruses harboring substitutions conferring reduced susceptibility to stavudine and zidovudine (M41L, D67N, K70R, L210W, T215Y/F, and K219Q/E) or didanosine (L74V) remained sensitive to FTC.

Tenofovir DF: Cross resistance has been observed among NRTIs. The K65R substitution in HIV-1 RT selected by tenofovir is also selected in some HIV-1 infected patients treated with abacavir, or didanosine. HIV-1 isolates with the K65R substitution also showed reduced susceptibility to FTC and lamivudine. Therefore, cross resistance among these drugs may occur in patients whose virus harbors the K65R substitution. The K70E substitution selected clinically by TDF results in reduced susceptibility to abacavir,

didanosine, FTC, and lamivudine. HIV-1 isolates from subjects (N=20) whose HIV-1 expressed a mean of 3 zidovudine-associated RT amino acid substitutions (M41L, D67N, K70R, L210W, T215Y/F, or K219Q/E/N) showed a 3.1-fold decrease in the susceptibility to tenofovir. Subjects whose virus expressed an L74V substitution without zidovudine resistance associated substitutions (N=8) had reduced response to TDF. Limited data are available for patients whose virus expressed a Y115F substitution (N=3), Q151M substitution (N=2), or T69 insertion (N=4), all of whom had a reduced response.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Efavirenz: Long-term carcinogenicity studies in mice and rats were carried out with EFV. Mice were dosed with 0, 25, 75, 150, or 300 mg/kg/day for 2 years. Incidences of hepatocellular adenomas and carcinomas and pulmonary alveolar/bronchiolar adenomas were increased above background in females. No increases in tumor incidence above background were seen in males. In studies in which rats were administered EFV at doses of 0, 25, 50, or 100 mg/kg/day for 2 years, no increases in tumor incidence above background were observed. The systemic exposure (based on AUCs) in mice was approximately 1.7-fold that in humans receiving the 600-mg/day dose. The exposure in rats was lower than that in humans. The mechanism of the carcinogenic potential is unknown. However, in genetic toxicology assays, EFV showed no evidence of mutagenic or clastogenic activity in a battery of *in vitro* and *in vivo* studies. These included bacterial mutation assays in *S. typhimurium* and *E. coli*, mammalian mutation assays in Chinese hamster ovary cells, chromosome aberration assays in human peripheral blood lymphocytes or Chinese hamster ovary cells, and an *in vivo* mouse bone marrow micronucleus assay. Given the lack of genotoxic activity of EFV, the relevance to humans of neoplasms in EFV-treated mice is not known.

Efavirenz did not impair mating or fertility of male or female rats, and did not affect sperm of treated male rats. The reproductive performance of offspring born to female rats given EFV was not affected. Because of the rapid clearance of EFV in rats, systemic drug exposures achieved in these studies were equivalent to or below those achieved in humans given therapeutic doses of EFV.

Emtricitabine: In long-term carcinogenicity studies of FTC, no drug-related increases in tumor incidence were found in mice at doses up to 750 mg/kg/day (26 times the human systemic exposure at the therapeutic dose of 200 mg/day) or in rats at doses up to 600 mg/day (31 times the human systemic exposure at the therapeutic dose).

Emtricitabine was not genotoxic in the reverse mutation bacterial test (Ames test), or the mouse lymphoma or mouse micronucleus assays.

Emtricitabine did not affect fertility in male rats at approximately 140-fold or in male and female mice at approximately 60-fold higher exposures (AUC) than in humans given the recommended 200 mg daily dose. Fertility was normal in the offspring of mice exposed daily from before birth (*in utero*) through sexual maturity at daily exposures (AUC) of approximately 60-fold higher than human exposures at the recommended 200 mg daily dose.

Tenofovir DF: Long-term oral carcinogenicity studies of TDF in mice and rats were carried out at exposures up to approximately 16 times (mice) and 5 times (rats) those

observed in humans at the therapeutic dose for HIV-1 infection. At the high dose in female mice, liver adenomas were increased at exposures 16 times that in humans. In rats, the study was negative for carcinogenic findings at exposures up to 5 times that observed in humans at the therapeutic dose.

TDF was mutagenic in the *in vitro* mouse lymphoma assay and negative in an *in vitro* bacterial mutagenicity test (Ames test). In an *in vivo* mouse micronucleus assay, TDF was negative when administered to male mice.

There were no effects on fertility, mating performance, or early embryonic development when TDF was administered to male rats at a dose equivalent to 10 times the human dose based on body surface area comparisons for 28 days prior to mating and to female rats for 15 days prior to mating through Day 7 of gestation. There was, however, an alteration of the estrous cycle in female rats.

13.2 Animal Toxicology and/or Pharmacology

Efavirenz: Nonsustained convulsions were observed in 6 of 20 monkeys receiving EFV at doses yielding plasma AUC values 4- to 13-fold greater than those in humans given the recommended dose.

Tenofovir DF: Tenofovir and TDF administered in toxicology studies to rats, dogs, and monkeys at exposures (based on AUCs) greater than or equal to 6-fold those observed in humans caused bone toxicity. In monkeys the bone toxicity was diagnosed as osteomalacia. Osteomalacia observed in monkeys appeared to be reversible upon dose reduction or discontinuation of tenofovir. In rats and dogs, the bone toxicity manifested as reduced bone mineral density. The mechanism(s) underlying bone toxicity is unknown.

Evidence of renal toxicity was noted in 4 animal species administered tenofovir and TDF. Increases in serum creatinine, BUN, glycosuria, proteinuria, phosphaturia and/or calciuria and decreases in serum phosphate were observed to varying degrees in these animals. These toxicities were noted at exposures (based on AUCs) 2- to 20-times higher than those observed in humans. The relationship of the renal abnormalities, particularly the phosphaturia, to the bone toxicity is not known.

14 CLINICAL STUDIES

Clinical Study 934 (NCT00112047) supports the use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in antiretroviral treatment-naïve HIV-1 infected patients.

Clinical Study 073 (NCT00365612) provides clinical experience in subjects with stable, virologic suppression and no history of virologic failure who switched from their current regimen to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.

In antiretroviral treatment-experienced patients, the use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablets may be considered for patients with HIV-1 strains that are expected to be susceptible to the components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet as assessed by treatment history or by genotypic or phenotypic testing [see *Microbiology (12.4)*].

Study 934: Data through 144 weeks are reported for Study 934, a randomized, open-

label, active-controlled multicenter trial comparing FTC + TDF administered in combination with EFV versus zidovudine/lamivudine fixed-dose combination administered in combination with EFV in 511 antiretroviral-naïve subjects. From Weeks 96 to 144 of the trial, subjects received FTC/TDF fixed-dose combination with EFV in place of FTC + TDF with EFV. Subjects had a mean age of 38 years (range 18-80); 86% were male, 59% were Caucasian, and 23% were Black. The mean baseline CD4⁺ cell count was 245 cells/mm³ (range 2-1191), and median baseline plasma HIV-1 RNA was 5.01 log₁₀ copies/mL (range 3.56-6.54). Subjects were stratified by baseline CD4⁺ cell count (< or ≥200 cells/mm³), and 41% had CD4⁺ cell counts <200 cells/mm³. Fifty-one percent (51%) of subjects had baseline viral loads >100,000 copies/mL. Treatment outcomes through 48 and 144 weeks for those subjects who did not have EFV resistance at baseline (N=487) are presented in Table 8.

Table 8 Virologic Outcomes of Randomized Treatment at Weeks 48 and 144 (Study 934)

Outcomes	At Week 48		At Week 144	
	FTC + TDF +EFV (N=244)	AZT/3TC +EFV (N=243)	FTC + TDF +EFV (N=227) ^a	AZT/3TC +EFV (N=229) ^a
Responder ^b	84%	73%	71%	58%
Virologic failure ^c	2%	4%	3%	6%
Rebound	1%	3%	2%	5%
Never suppressed	0%	0%	0%	0%
Change in antiretroviral regimen	1%	1%	1%	1%
Death	<1%	1%	1%	1%
Discontinued due to adverse event	4%	9%	5%	12%
Discontinued for other reasons ^d	10%	14%	20%	22%

^a. Subjects who were responders at Week 48 or Week 96 (HIV-1 RNA <400 copies/mL) but did not consent to continue trial after Week 48 or Week 96 were excluded from analysis.

^b. Subjects achieved and maintained confirmed HIV-1 RNA <400 copies/mL through Weeks 48 and 144.

^c. Includes confirmed viral rebound and failure to achieve confirmed HIV-1 RNA <400 copies/mL through Weeks 48 and 144.

^d. Includes lost to follow-up, patient withdrawal, noncompliance, protocol violation and other reasons.

Through Week 48, 84% and 73% of subjects in the FTC + TDF group and the zidovudine/lamivudine group, respectively, achieved and maintained HIV-1 RNA <400 copies/mL (71% and 58% through Week 144). The difference in the proportion of subjects who achieved and maintained HIV-1 RNA <400 copies/mL through 48 weeks largely results from the higher number of discontinuations due to adverse events and other reasons in the zidovudine/lamivudine group in this open-label trial. In addition, 80% and 70% of subjects in the FTC + TDF group and the zidovudine/lamivudine group, respectively, achieved and maintained HIV-1 RNA <50 copies/mL through Week 48 (64% and 56% through Week 144). The mean increase from baseline in CD4⁺ cell count was

and 190 cells/mm³ at Week 48 (312 and 271 cells/mm³ at Week 144).

Through 48 weeks, 7 subjects in the FTC + TDF group and 5 subjects in the zidovudine/lamivudine group experienced a new CDC Class C event (10 and 6 subjects through 144 weeks).

Study 073: Study 073 was a 48-week open-label, randomized clinical trial in subjects with stable virologic suppression on combination antiretroviral therapy consisting of at least two NRTIs administered in combination with a protease inhibitor (with or without ritonavir) or a NNRTI.

To be enrolled, subjects were to have HIV-1 RNA <200 copies/mL for at least 12 weeks on their current regimen prior to trial entry with no known HIV-1 substitutions conferring resistance to the components of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and no history of virologic failure.

The trial compared the efficacy of switching to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet or staying on the baseline antiretroviral regimen (SBR). Subjects were randomized in a 2:1 ratio to switch to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet (N=203) or stay on SBR (N=97). Subjects had a mean age of 43 years (range 22-73 years); 88% were male, 68% were white, 29% were Black or African-American, and 3% were of other races. At baseline, median CD4⁺ cell count was 516 cells/mm³, and 96% had HIV-1 RNA <50 copies/mL. The median time since onset of antiretroviral therapy was 3 years, and 88% of subjects were receiving their first antiretroviral regimen at trial enrollment.

At Week 48, 89% and 87% of subjects who switched to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet maintained HIV RNA <200 copies/mL and <50 copies/mL, respectively, compared to 88% and 85% who remained on SBR; this difference was not statistically significant. No changes in CD4⁺ cell counts from baseline to Week 48 were observed in either treatment arm.

16 HOW SUPPLIED/STORAGE AND HANDLING

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablets are pink colored, capsule-shaped, biconvex, film-coated tablets debossed with 'C210' on one side and plain on other side.

Each bottle contains:

30 tablets (NDC 69097-210-02) and silica gel desiccant, and is closed with a child-resistant closure.

Store at 25°C (77°F); excursions permitted to 15°C -30°C (59°F -86°F) [See USP *Controlled Room Temperature*].

- Keep container tightly closed.
- Dispense only in original container.
- Do not use if seal over bottle opening is broken or missing.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Severe Acute Exacerbation of Hepatitis B in Patients Coinfected with HIV-1 and HBV

Inform patients that severe acute exacerbations of hepatitis B have been reported in patients who are coinfecting with HBV and HIV-1 and have discontinued FTC or TDF, and may occur with discontinuation of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Advise patients not to discontinue Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet without first informing their healthcare provider. All patients should be tested for HBV infection before or when starting Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and those who are infected with HBV need close medical follow-up for several months after stopping Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet to monitor for exacerbations of hepatitis [see *Warnings and Precautions (5.1)*].

Rash

Inform patients that a common side effect is rash, and that rashes usually go away without any change in treatment. However, since rash may be serious, advise patients to contact their physician promptly if rash occurs [see *Warnings and Precautions (5.2)*].

Hepatotoxicity

Inform patients to watch for early warning signs of liver inflammation or failure, such as fatigue, weakness, lack of appetite, nausea and vomiting, as well as later signs such as jaundice, confusion, abdominal swelling, and discolored feces, and to consult their health care professional without delay if such symptoms occur [see *Warnings and Precautions (5.3) and Adverse Reactions (6.1)*].

Drug Interactions

Advise patients that Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet may interact with many drugs; therefore, advise patients to report to their healthcare provider the use of any other medication, including other drugs for treatment of hepatitis C virus [see *Warnings and Precautions (5.4) and Drug Interactions (7)*].

Psychiatric Symptoms

- Inform patients that serious psychiatric symptoms including severe depression, suicide attempts, aggressive behavior, delusions, paranoia, psychosis-like symptoms, and catatonia have been reported in patients receiving EFV, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet [see *Warnings and Precautions (5.5)*].
- Advise patients to seek immediate medical evaluation if they experience severe psychiatric adverse experiences.
- Advise patients to inform their physician of any history of mental illness or substance abuse.

Nervous System Symptoms

- Inform patients that central nervous system symptoms (NSS) including dizziness, insomnia, impaired concentration, drowsiness, and abnormal dreams, are commonly reported during the first weeks of therapy with EFV, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Dosing at bedtime may improve the tolerability of these symptoms, which are likely to improve with continued therapy.
- Alert patients to the potential for additive effects when Efavirenz, Emtricitabine and

Tenofovir disoproxil fumarate tablet is used concomitantly with alcohol or psychoactive drugs.

- Instruct patients that if they experience NSS to avoid potentially hazardous tasks such as driving or operating machinery [see *Warnings and Precautions (5.6)* and *Dosage and Administration (2.2)*].
- Inform patients that there is a risk of developing late-onset neurotoxicity, including ataxia and encephalopathy, which may occur months to years after beginning therapy with EFV, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet [see *Warnings and Precautions (5.6)*].

New Onset or Worsening Renal Impairment

Inform patients that renal impairment, including cases of acute renal failure and Fanconi syndrome, has been reported. Advise patients to avoid using Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with concurrent or recent use of a nephrotoxic agent (e.g., high-dose or multiple NSAIDs) [see *Warnings and Precautions (5.7)*].

Embryo-Fetal Toxicity

Apprise patients of the potential harm to the fetus if Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is used during the first trimester of pregnancy, or if the patient becomes pregnant while taking this drug. Instruct adults and adolescents of childbearing potential receiving Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet to avoid pregnancy and to notify their healthcare provider if they become pregnant or plan to become pregnant while taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet [see *Warnings and Precautions (5.8)*]. A reliable form of barrier contraception must always be used in combination with other methods of contraception, including oral or other hormonal contraception. Because of the long half-life of EFV, recommend use of adequate contraceptive measures for 12 weeks after discontinuation of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet [see *Use in Specific Populations (8.1, 8.3)*].

Bone Loss and Mineralization Defects

Inform patients that decreases in bone mineral density have been observed with the use of TDF, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Advise patients that bone mineral density monitoring may be performed in patients who have a history of pathologic bone fracture or other risk factors for osteoporosis or bone loss [see *Warnings and Precautions (5.9)*].

Convulsions

Inform patients that convulsions have been reported with the use of EFV, a component of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Patients who are receiving concomitant anticonvulsant medications primarily metabolized by the liver may require periodic monitoring of plasma levels [see *Warnings and Precautions (5.10)* and *Drug Interactions (7.3)*].

Lactic Acidosis and Severe Hepatomegaly

Inform patients that lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported. Treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet should be suspended in any patient who develops clinical symptoms suggestive of lactic acidosis or pronounced hepatotoxicity [see *Warnings and*

Precautions (5.11)].

Immune Reconstitution Syndrome

Inform patients that in some patients with advanced HIV infection (AIDS), signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is believed that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms. Advise patients to inform their healthcare provider immediately of any symptoms of infection *[see Warnings and Precautions (5.12)].*

Fat Redistribution

Inform patients that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy, including Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and that the cause and long-term health effects of these conditions are not known *[see Warnings and Precautions (5.13)].*

Dosing Instructions

Advise patients to take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet orally on an empty stomach and that it is important to take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet on a regular dosing schedule to avoid missing doses. Advise patients that dosing at bedtime may improve the tolerability of nervous system symptoms *[see Dosage and Administration (2.1)].*

Pregnancy Registry

Advise patients that there is a pregnancy exposure registry that monitors pregnancy outcomes in patients exposed to Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet during pregnancy *[see Use in Specific Populations (8.1)].*

Lactation

Instruct patients not to breastfeed because HIV-1 can be passed to the baby in the breast milk *[see Use in Specific Populations (8.2)].*

Disclaimer: Other brands listed are the registered trademarks of their respective owners and are not trademarks of Cipla Limited.

Manufactured by:

Cipla Ltd. MIDC,
Patalganga
India

Manufactured for:

Cipla USA, Inc.
10 Independence Boulevard, Suite 300
Warren, New Jersey - 07059

Revised: 12/2020

Patient Information

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate

(e FAV e renz, em trye SYE ta been, & ten OF oh vir dye soe PROX il FUE ma rate)

Tablets

What is the most important information I should know about Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet can cause serious side effects, including:

- **Worsening of hepatitis B virus (HBV) infection.** Your healthcare provider will test you for HBV before starting treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. If you have HBV infection and take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, your HBV may get worse (flare-up) if you stop taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. A "flare-up" is when your HBV infection suddenly returns in a worse way than before.
- Do not stop taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet without first talking with your healthcare provider.
- Do not run out of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Refill your prescription or talk to your healthcare provider before your Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is all gone.
- If you stop taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, your healthcare provider will need to check your health often and do blood tests regularly for several months to check your HBV infection, or give you a medication to treat hepatitis B. Tell your healthcare provider about any new or unusual symptoms you may have after you stop taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.

For more information about side effects see the section, "**What are the possible side effects of**

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?"

What is Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is a prescription medicine that contains efavirenz, emtricitabine, and tenofovir disoproxil fumarate combined in 1 tablet. Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is used alone as a complete regimen, or in combination with other anti-HIV-1 medicines to treat people with HIV-1 infection who weigh at least 88 lbs (40 kg). It is not known if Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet is safe and effective for use in children with HIV-1 infection who weigh less than 88 lbs (40 kg).

Who should not take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

Do not take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet if you:

- are allergic to efavirenz
- take the medicine called voriconazole, elbasvir or grazoprevir

Ask your healthcare provider if you are not sure if you take any of these medicines.

Before taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, **tell your healthcare provider about all of your medical conditions, including if you:**

- have liver problems, including hepatitis B or C virus infection
- have heart problems
- have or have had mental problems
- have a history of drug or alcohol abuse
- have nervous system problems
- have kidney problems or receive kidney dialysis treatment
- have bone problems
- have had seizures or take medicines used to treat seizures
- are pregnant or plan to become pregnant. Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet can harm your unborn baby. If you are able to become pregnant, your healthcare provider should do a pregnancy test before you start treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. **You should not become pregnant during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and for 12 weeks after stopping treatment.** Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
- **Females who are able to become pregnant should use 2 effective forms of birth control (contraception) during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and for 12 weeks after stopping treatment.**
- **A barrier form of birth control should always be used along with another type of birth control.** Barrier forms of birth control may include condoms, contraceptive sponges, diaphragm with spermicide, and cervical cap.
- **Birth control methods that contain the hormone progesterone such as birth control pills, injections, vaginal rings, or implants, may not work as well while taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.**
- Talk to your healthcare provider about birth control methods that may be right for you during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
- **Pregnancy Registry:** There is a pregnancy registry for women who take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.
- are breastfeeding or plan to breastfeed. Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet can pass into your breast milk. Do not breastfeed because of the risk of passing HIV-1 to your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and some medicines may interact with each other causing serious side effects.

You can ask your healthcare provider or pharmacist for a list of medicines that interact with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Do not start a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with other medicines.

How should I take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

- Take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet exactly as your healthcare provider tells you to.
- If you take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet with other medicines used to treat HIV-1, your healthcare provider will tell you what medicines to take and how to take them.
- Take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet 1 time each day on an empty stomach. You should take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet at the same time each day.
- Taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet at bedtime may make some side effects less bothersome.
- **Do not miss a dose of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.** Missing a dose lowers the amount of medicine in your blood. Refill your Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet prescription before you run out of medicine.
- Do not change your Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet dose or stop taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet without first talking with your healthcare provider. Stay under a healthcare provider's care during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
- If you take too much Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet, call your healthcare provider or go to the nearest hospital emergency room right away.

What should I avoid while taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

- Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet can cause dizziness, impaired concentration and drowsiness. If you have these symptoms, do not drive a car, use heavy machinery, or do anything that requires you to be alert.

What are the possible side effects of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet may cause serious side effects, including:

- **See "What is the most important information I should know about Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?"**
- **Rash.** Rash is a serious side effect but may also be common. Rashes will usually go away without any change in your treatment. Tell your healthcare provider right away if you develop a rash during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
- **Severe liver problems.** In rare cases, severe liver problems can happen that can lead to death. **Tell your healthcare provider right away if you get these symptoms** : skin or the white part of your eyes turns yellow, dark "tea-colored"

urine, light-colored stools, loss of appetite for several days or longer, nausea, or stomach-area pain.

- **Mental problems.** Serious mental problems including severe depression, suicidal thoughts and actions, aggressive behavior, delusions, catatonia, and paranoid and manic reactions have happened in people who take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. These mental health problems may happen more often in people who have a history of mental problems or drug use, or who take medicines to treat mental problems. Tell your healthcare provider right away if you develop serious mental problems during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
- **Nervous system problems.** Nervous system problems usually begin during the first or second day of treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and usually go away after 2 to 4 weeks of treatment. Some symptoms may occur months to years after beginning Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet therapy. These symptoms may become more severe if you drink alcohol or take mood altering (street) drugs while taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Tell your healthcare provider right away if you develop nervous system problems during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Symptoms of nervous system problems may include:
 - dizziness
 - problems concentrating
 - abnormal dreams
 - unusually happy mood
 - agitation
 - thought problems
 - slow thoughts and physical movement
 - problems sleeping
 - excessive sleepiness or difficulty awakening
 - seeing or hearing things that are not real (hallucinations)
 - confusion
 - memory problems
 - lack of coordination or difficulty with balance

If you have dizziness, trouble concentrating or sleepiness, do not drive a car, use machinery, or do anything that needs you to be alert.

- **New or worse kidney problems, including kidney failure.** Your healthcare provider should do blood and urine tests to check your kidneys before you start and during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Your healthcare provider may tell you to stop taking Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet if you develop new or worse kidney problems during treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
- **Bone problems** can happen in some people who take Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet. Bone problems include bone pain or softening or thinning of bones, which may lead to fractures. Your healthcare provider may need to do tests to check your bones.
- **Seizures** . Your healthcare provider may do blood tests during treatment with

Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet if you take certain medicines used to prevent seizures.

- **Too much lactic acid in your blood (lactic acidosis).** Too much lactic acid is a serious but rare medical emergency that can lead to death. Tell your healthcare provider right away if you develop any of these symptoms:
 - weakness or being more tired than usual
 - being short of breath or fast breathing
 - cold or blue hands and feet
 - fast or abnormal heartbeat
 - unusual muscle pain
 - stomach pain with nausea and vomiting
 - feel dizzy or lightheaded
- **Changes in your immune system (Immune Reconstitution Syndrome) can happen when an HIV-1 infected person starts taking HIV-1 medicines.** Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you develop any new symptoms after starting treatment with Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.
- **Changes in body fat.** Changes in body fat distribution or accumulation have happened in some people taking HIV-1 medicines, including an increased amount of fat in the upper back and neck ("buffalo hump"), in the breasts, and around the trunk. Loss of fat from the legs, arms, and face may also happen. The cause and long-term health effects of these body fat changes are not known.

The most common side effects of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet include:

- diarrhea
- tiredness
- dizziness
- problems sleeping
- rash
- nausea
- headache
- depression
- abnormal dreams

These are not all the possible side effects of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

- Store Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet at room temperature 77°F (25°C).
- Keep Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet in its original container and keep the container tightly closed.

Keep Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet and all other medicines out of reach of children.

General information about the safe and effective use of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet:

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet for a condition for which it was not prescribed. Do not give Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet to other people, even if they have the same symptoms that you have. It may harm them. You can ask your healthcare provider or pharmacist for information about Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet that is written for health professionals.

What are the ingredients of Efavirenz, Emtricitabine and Tenofovir disoproxil fumarate tablet?

Active Ingredients: efavirenz, emtricitabine, and tenofovir disoproxil fumarate

Inactive Ingredients: Croscarmellose sodium, hydroxypropyl cellulose, hypromellose, isopropyl alcohol, microcrystalline cellulose, magnesium stearate, red iron oxide, sodium lauryl sulfate. The tablets are film-coated with a coating material opadry AMB 80W54485 pink containing polyvinyl alcohol-part, hydrolyzed, titanium dioxide, talc, lecithin (soya), xanthan gum, iron oxide yellow, iron oxide red and opadry AMB 80W56843 brown containing polyvinyl alcohol-part, hydrolyzed, titanium dioxide, talc, lecithin (soya), xanthan gum, iron oxide red.

Disclaimer: Other brands listed are the registered trademarks of their respective owners and are not trademarks of Cipla Limited.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Manufactured by:

Cipla Ltd. MIDC,
Patalganga India

Manufactured for:

Cipla USA, Inc.
10 Independence Boulevard, Suite 300
Warren, New Jersey – 07059

Revised: 12/2020

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 69097-210-02 Rx only

Efavirenz, Emtricitabine and Tenofovir Disoproxil

Fumarate Tablets

600 mg/200mg/300 mg

Note to Pharmacist: Do not cover ALERT box with pharmacy label.

ALERT: Find out about medicines that, should Not be taken with Efavirenz, Emtricitabine and Tenofovir Disoproxil Fumarate Tablets 600 mg/200 mg/300 mg.

30 Tablets

Cipla

NDC 69097-210-02 Rx Only

**Efavirenz,
Emtricitabine
and Tenofovir
Disoproxil Fumarate
Tablets**

600 mg/200 mg/300 mg

Note to Pharmacist: Do not cover ALERT box with pharmacy label.

ALERT: Find out about medicines that should NOT be taken with Efavirenz, Emtricitabine and Tenofovir Disoproxil Fumarate Tablets 600 mg/200 mg/300 mg.

30 Tablets

Cipla

Each film-coated tablet contains:
Efavirenz USP 600 mg
Emtricitabine 200 mg
Tenofovir Disoproxil Fumarate 300 mg
(Equivalent to Tenofovir Disoproxil 245 mg)
Usual Dosage: See package insert for dosage and administration.
Store at 25°C (77°F). (see insert)
Keep container tightly closed. Dispense only in original container.
Manufactured by: **Cipla Ltd.**, MIDC Patalganga, India
Manufactured for: **Cipla USA, Inc.**
10 Independence Boulevard, Suite 300, Warren, NJ 07059

208436 Rev. 11/2019

S/N
EXP
LOT

M. L. KD-620 GTIN 00369097210021

69097210021

Area for Batch overprinting
(Serial No., Expiry & Lot will be
overprinted during commercial packing)

NDC 69097-210-02 Rx only

Efavirenz, Emtricitabine and Tenofovir Disoproxil


Fumarate Tablets

600 mg/200mg/300 mg

Note to Pharmacist: Do not cover ALERT box with pharmacy label.

ALERT: Find out about medicines that, should Not be taken with Efavirenz, Emtricitabine and Tenofovir Disoproxil Fumarate Tablets 600 mg/200 mg/300 mg.

30 Tablets



EFAVIRENZ, EMTRICITABINE AND TENOFOVIR DISOPROXIL FUMARATE

efavirenz, emtricitabine and tenofovir disoproxil fumarate tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:69097-210
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EFAVIRENZ (UNII: JE6H2O27P8) (EFAVIRENZ - UNII:JE6H2O27P8)	EFAVIRENZ	600 mg
EMTRICITABINE (UNII: G70B4ETF4S) (EMTRICITABINE - UNII:G70B4ETF4S)	EMTRICITABINE	200 mg
TENOFOVIR DISOPROXIL FUMARATE (UNII: OTT9J7900I) (TENOFOVIR ANHYDROUS - UNII:W4HFE001U5)	TENOFOVIR DISOPROXIL FUMARATE	300 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

HYDROXYPROPYL CELLULOSE (110000 WAMW) (UNII: 5Y0974F5PW)
ISOPROPYL ALCOHOL (UNII: ND2M416302)
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)
MAGNESIUM STEARATE (UNII: 70097M6I30)
FERRIC OXIDE RED (UNII: 1K09F3G675)
SODIUM LAURYL SULFATE (UNII: 368GB5141J)
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TALC (UNII: 7SEV7J4R1U)
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)
XANTHAN GUM (UNII: TTV12P4NEE)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

Product Characteristics

Color	PINK	Score	no score
Shape	CAPSULE (Biconvex)	Size	21mm
Flavor		Imprint Code	Cipla;210
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69097-210-02	30 in 1 BOTTLE; Type 0: Not a Combination Product	03/30/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206894	03/30/2021	

Labeler - Cipla USA Inc. (078719707)

Registrant - Cipla Patalganga (916940208)

Establishment

Name	Address	ID/FEI	Business Operations
Cipla Patalganga		916940208	ANALYSIS(69097-210) , LABEL(69097-210) , MANUFACTURE(69097-210) , PACK(69097-210)

Establishment

Name	Address	ID/FEI	Business Operations
Genvion Corporation 500 Camiel Sys Street, Winnipeg, Manitoba R2J 4K2, Canada		245221226	ANALYSIS(69097-210) , LABEL(69097-210) , PACK(69097-210) , MANUFACTURE(69097-210)

