ALLERGY MULTI SYMPTOM- acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet, coated MARC GLASSMAN, INC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1128B-MAR-2022-1005

Drug Facts

Active ingredients (in each caplet)	Purpose		
Acetaminophen 325 mg	Pain reliever		
Chlorpheniramine maleate 2 mg	Antihistamine		
Phenylephrine HCl 5 mg	Nasal decongestant		

Uses

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 - headache
 - sinus congestion and pressure
 - nasal congestion
 - runny nose and sneezing
 - minor aches and pains
- temporarily relieves these additional symptoms of hay fever:
 - itching of the nose or throat
 - itchy, watery eyes
- helps clear nasal passages
- helps decongest sinus openings and passages

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	 take 2 caplets every 4 hours swallow whole; do not crush, chew, or dissolve do not take more than 10 caplets in 24 hours
children under 12 years	 ask a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, flavor, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

Marc's ®

NDC 68998-228-02

Allergy Multi-Symptom

Acetaminophen, Phenylephrine HCl, Chlorpheniramine Maleate

PAIN RELIEVER, NASAL DECONGESTANT, ANTIHISTAMINE

For Relief of:

- Headache
- Sinus Pressure & Nasal Congestion
- Runny Nose, Sneezing & Itchy, Watery Eyes

May Cause Drowsiness

Actual Size

24 COOL TASTE CAPLETS

	NC		Marcis. Allergy Multi-Symptom	
NC	Orug Facts (continued) Drug Facts (continued) Warnings Wernung free pool of control of the pool of the second hard pool of the	Drug Facts (continued) Distributed by: More Glossman, Inc. Inactive ingredients accutime potassium, coloidal Distributed by: More Glossman, Inc. Ren down of the matching of active potassium, coloidal fav. Distributed by: More Clossman, Inc. More Glossman, Inc. Ren down of the matching of active potassium, coloidal fav. Distributed by: More Clossman, Inc. More Clossman, Inc. Ren down of the matching of active potassim stands Close reliance, Close active Close close active active potassime active potagione gloci, stance acti, tak, titarium dicude Distributed stuck, Mate TORN OR BROKEN	<page-header><section-header><section-header><text><text><text><text><text></text></text></text></text></text></section-header></section-header></page-header>	Drug Facts Drug Facts (continued) Drug Facts Drug Facts (continued) Drug Facts Drug Facts (continued) Detrive inpresentation Enviropease actionaries Detrive inpresentation Enviropease actionaries Distribution Enviropease actionaries Distrese actionaries
	Drug Fact Warnings Warnings Warnings wars Seret wars Seret warn of a sere warning warn	Drug F Inactiv favor to poverheie propylene	Provisiness 24 COOL TASTE CAPLETS F1128802MAR_R1 0 94442 60143 7	Drug Facts

ALLERGY MULTI SYMPTOM

acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet, coated

Product Information						
Product Type	HUMAN OTC DRUG	Item Code (Source)		NDC:6899	NDC:68998-228	
Route of Administration	ORAL					
Active Ingredient/Active Moiety						
Ingredient Name Basis of Strength Strength						
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII:3U6I01965U)			CHLORPHENIRAMI MALEATE	•	2 mg	
PHENYLEPHRINE HYDROCHLOF UNII:1W5297W6MV)	IDE (UNII: 04JA59TNSJ) (PHE	NYLEPHRINE -	PHENYLEPHRINE HYDROCHLORIDE		5 mg	

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CROSPOVIDONE (UNII: 2S7830E561)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: 08232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
Product Characteristics	

Color	white (Off-White)	Score	no score	
Shape	OVAL	Size	17mm	
Flavor	MINT	Imprint Code	AAA;1128	
Contains				

Packaging

ltem Code	Package Description	Marketing Start Date	Marketing End Date	
NDC:68998- 228-02	2 in 1 CARTON	06/22/2020		
	12 in 1 BLISTER PACK; Type 0: Not a Combination Product			
Marketing Information				
	NDC:68998- 228-02	NDC:68998- 228-02 2 in 1 CARTON 12 in 1 BLISTER PACK; Type 0: Not a Combination Product	Item CodePackage DescriptionDateNDC:68998- 228-022 in 1 CARTON06/22/202012 in 1 BLISTER PACK; Type 0: Not a Combination ProductProduct	

Marketing	Application Number or Monograph	Marketing Start	Marketing End
Category	Citation	Date	Date
OTC monograph final	part341	06/22/2020	

Labeler - MARC GLASSMAN, INC. (094487477)

Revised: 10/2022

MARC GLASSMAN, INC.