

IBUPROFEN- ibuprofen capsule, liquid filled

H E B

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

***nonsteroidal anti-inflammatory drug**

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
 - take 1 capsule every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 capsule, 2 capsules may be used
 - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F).
- read all warnings and directions before use. Keep carton.
- swallow whole, do not crush, chew, or dissolve

Inactive ingredients

FD&C blue #1, FD&C yellow#6, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

Questions or comments?

Call toll free **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Compare to Advil® Liqui-Gels® active ingredient**

Solubilized

Ibuprofen Capsules, 200 mg

Pain Reliever/Fever Reducer **(NSAID)**

For Body Aches and Pain

SOFTGELS†

(†Liquid Filled Capsules)

**This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

TAMPER EVIDENT: DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

Product Label

Drug Facts	
Active ingredient (in each capsule) Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* <small>*nonsteroidal anti-inflammatory drug</small>	Purpose Pain reliever/fever reducer
Uses	
<ul style="list-style-type: none"> ■ Temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ headache ■ minor pain of arthritis ■ backache ■ menstrual cramps ■ Temporarily reduces fever 	
Warnings	
<p>Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:</p> <ul style="list-style-type: none"> ■ hives ■ facial swelling ■ shock ■ skin reddening ■ rash ■ blisters <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:</p> <ul style="list-style-type: none"> ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p>	
Do not use	
<ul style="list-style-type: none"> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery 	

Drug Facts (continued)
Ask a doctor before use if
<ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
<ul style="list-style-type: none"> ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug
When using this product
<ul style="list-style-type: none"> ■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
<ul style="list-style-type: none"> ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ chest pain ■ trouble breathing ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use.
<p>It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p>
Keep out of reach of children.
<p>In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>

Drug Facts (continued)	
Directions	
<ul style="list-style-type: none"> ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: <ul style="list-style-type: none"> ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor 	
Other information	
<ul style="list-style-type: none"> ■ each capsule contains: potassium 20 mg ■ store at 20° to 25°C (68° to 77°F) ■ avoid excessive heat above 40°C (104°F) ■ read all warnings and directions before use. Keep carton. ■ swallow whole; do not crush, chew, or dissolve 	
Inactive ingredients	
<p>FD&C blue no. 1, FD&C yellow no. 6, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan</p>	
Questions or comments?	
<p>Call toll free: 1-877-753-3835 Monday-Friday 9AM-5PM EST</p>	
<p><small>*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.</small></p>	
<p>TAMPER EVIDENT: DO NOT USE IF SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.</p> <p>READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION.</p>	

Compare to Advil® Liqui-Gels®
active ingredient**

NDC 37808-750-40



Solubilized Ibuprofen
Capsules, 200 mg

Pain Reliever/Fever Reducer (NSAID)

For Body Aches and Pain



actual size

40 SOFTGELS[†]
(*Liquid Filled Capsules)

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

100% GUARANTEE promise

If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

9407-1804



0 41220 28428 3

FC005009 PLD-CS29A

Lot No.:

Exp. Date:

H-E-B Ibuprofen 200 mg Capsules

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:37808-750

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	AT146
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-750-20	1 in 1 BOX	08/31/2018	08/31/2025
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:37808-750-40	1 in 1 BOX	08/31/2018	08/31/2025
2		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:37808-750-80	80 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2018	08/31/2025
4	NDC:37808-750-30	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/31/2018	08/31/2025

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206999	08/31/2018	08/31/2025

