

**NIGHTTIME COLD AND FLU MULTI SYMPTOM- acetaminophen
dextromethorphan hydrobromide doxylamine succinate capsule, liquid filled
CARDINAL HEALTH 110, LLC. DBA LEADER 70000**

Safeway Nighttime Cold & Flu

Active Ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine Succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use

Other Information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive Ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or Comments?

Call **1-877-753-3929** Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO VICKS® NYQUIL™ COLD & FLU LIQUICAPS™ active ingredients†

Nighttime | Multi-Symptom

Cold & Flu Relief

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

Pain Reliever / Fever Reducer | Cough Suppressant | Antihistamine

Relieves: Fever, Sore Throat, Cough, Runny Nose, Sneezing, Headache, Aches and Pains

Alcohol-Free

SOFTGELS**

**Liquid-Filled Capsules

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil™, and LiquiCaps™ are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY CARDINAL HEALTH

DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

Package Label

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks[®], NyQuil[™], and LiquiCaps[™] are registered trademarks of The Procter & Gamble Company.

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST
Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, purified water, sorbitan, sorbitol, white ink
Other information ■ store between 15-30°C (59-86°F) ■ avoid excessive heat
Directions ■ do not take more than 4 doses in 24 hours ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ swallow whole; do not crush, chew, or dissolve ■ children under 12 years: do not use
Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Keep out of reach of children.
If pregnant or breast-feeding, ask a health professional before use.
Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus)
Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers
When using this product ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ avoid alcoholic drinks ■ marked drowsiness may occur ■ be careful when driving a motor vehicle or operating machinery
Drug Facts (continued)

Drug Facts (continued) ■ runny nose and sneezing ■ cough due to minor throat and bronchial irritation
Warnings ■ liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Drug Facts (continued)

Drug Facts
Active ingredients (in each softgel) Acetaminophen 325 mg Dextromethorphan HBr 15 mg Doxylamine succinate 6.25 mg
Purposes Pain reliever/fever reducer Cough suppressant Antihistamine
Uses ■ temporarily relieves common cold and flu symptoms: ■ sore throat ■ headache ■ minor aches and pains ■ fever

NDC 70000-0516-1

Nighttime | Multi-Symptom

Cold & Flu Relief

Acetaminophen | Dextromethorphan HBr | Doxylamine Succinate
Pain Reliever / Fever Reducer | Cough Suppressant | Antihistamine

Relieves: Fever, Sore Throat,
Cough, Runny Nose, Sneezing,
Headache, Minor Aches & Pains
Alcohol-Free

16 SOFTGELS**
**Liquid-Filled Capsules



Actual Size

COMPARE TO
VICKS[®] NYQUIL[™]
COLD & FLU
LIQUICAPS[™]
active ingredients[†]

100% Money
Back Guarantee

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

LD-C585C
0006966

at No.:
xp. Date:



DISTRIBUTED BY CARDINAL HEALTH
 DUBLIN, OHIO 43017
www.myleader.com 1-800-200-6313
Essential to Care™ since 1979
 Product of China

©2026 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the LEADER LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.



Return to place of purchase.



CIN 5556378 REV. 01/26



0 96295 13833 7

LEADER Nighttime Multi-Symptom Cold & Flu Relief

NIGHTTIME COLD AND FLU MULTI SYMPTOM

acetaminophen dextromethorphan hydrobromide doxylamine succinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0516
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	PC10

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0516-1	16 in 1 CARTON	08/30/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:70000-0516-2	24 in 1 CARTON	08/30/2019	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/30/2019	

Labeler - CARDINAL HEALTH 110, LLC. DBA LEADER 70000 (063997360)

Revised: 1/2026

CARDINAL HEALTH 110, LLC. DBA LEADER 70000