

IBUPROFEN ORAL- ibuprofen oral suspension
A-S Medication Solutions

Ibuprofen Oral Suspension USP, 100 mg/5 mL (OTC)

ACTIVE INGREDIENT(S)

(in each 5 mL)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/fever reducer

USE(S)

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

WARNINGS

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:☐

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- takes more or for a longer time than directed
- **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

DO NOT USE

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

ASK A DOCTOR BEFORE USE IF

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

WHEN USING THIS PRODUCT

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

| Weight (lb) | Age (yr) | Dose (mL)* |
|-------------|---------------|--------------|
| under 24 | under 2 years | ask a doctor |
| 24-35 lbs | 2-3 years | 5 mL |
| 36-47 lbs | 4-5 years | 7.5 mL |
| 48-59 lbs | 6-8 years | 10 mL |
| 60-71 lbs | 9-10 years | 12.5 mL |
| 72-95 lbs | 11 years | 15 mL |

*or as directed by a doctor

Other information

□

- **each 5 mL contains** : sodium 2 mg
- store between 20-25°C (68-77°F)
- **do not use if carton is opened or printed bottle neckband is broken or missing**
- see bottom panel for lot number and expiration date

INACTIVE INGREDIENT SECTION

Berry flavor (with dye)

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Yellow #10, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Berry flavor (dye free)

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Grape flavor

-

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #33, FD&C Blue #1, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

Bubble Gum flavor

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, D&C Red #33, FD&C Red #40, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum.

-

-

QUESTIONS SECTION

Call 1-888-588-1418 from 9 AM to 5 PM EST,
Monday-Friday.



Distributed by:
Camber Consumer Care, Inc.
Piscataway, NJ 08854, USA

Ibuprofen Oral



IBUPROFEN ORAL

ibuprofen oral suspension

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:50090-4129(NDC:69230-308)

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 100 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ACESULFAME POTASSIUM (UNII: 23OV73Q5G9) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| FD&C RED NO. 40 (UNII: WZB9127XOA) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SUCROSE (UNII: C151H8M554) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |

Product Characteristics

| | | | |
|----------|--------|--------------|--|
| Color | orange | Score | |
| Shape | | Size | |
| Flavor | BERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:50090-4129-0 | 1 in 1 CARTON | 02/07/2019 | |
| 1 | | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA210602 | 11/23/2018 | |

Labeler - A-S Medication Solutions (830016429)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--------------------------|---------|-----------|---------------------|
| A-S Medication Solutions | | 830016429 | RELABEL(50090-4129) |

Revised: 12/2023

A-S Medication Solutions