

TOPCARE DAY TIME COLD AND FLU- acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution
Topco Associates LLC

Topco Associates LLC. Day Time Cold & Flu Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- sore throat
- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- fever
- headache
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen

- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- you get nervous, dizzy or sleepless

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions?

1-888-423-0139

Package/Label Principal Display Panel

TopCare® health

COMPARE TO VICKS® DAYQUIL® SEVERE ACTIVE INGREDIENTS

MAXIMUM STRENGTH RELIEF

Day Time Cold & Flu

SEVERE

PAIN RELIEVER – FEVER REDUCER – ACETAMINOPHEN

COUGH SUPPRESSANT – DEXTROMETHORPHAN HBr

EXPECTORANT – GUAIFENESIN

NASAL DECONGESTANT – PHENYLEPHRINE HCl

• Headache, Fever, Sore Throat, Minor Aches & Pains

• Nasal/Sinus Congestion & Sinus Pressure

• Cough

• Chest Congestion

Alcohol Free

Antihistamine Free

Non-Drowsy

12 FL OZ (355 mL)

ORIGINAL FLAVOR

Drug Facts (continued)

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ you get nervous, dizzy or sleepless ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed – see Overdose warning
■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over 30 mL every 4 hrs
children 6 to under 12 yrs 15 mL every 4 hrs
children 4 to under 6 yrs ask a doctor
children under 4 yrs do not use

Other information ■ each 15 mL contains: sodium 6 mg
■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-888-423-0139

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ELK GROVE VILLAGE, IL 60007
©TOPCO PERAD622 QUESTIONS 1-888-423-0139
topcare@topco.com www.topcarebrand.com
Visit here for more information: http://topbrnds.com/49011j

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org


36800 39399 8

TopCare
health

COMPARE TO VICKS® DAYQUIL®
SEVERE ACTIVE INGREDIENTS*

MAXIMUM STRENGTH RELIEF

Day Time
Cold & Flu
SEVERE

PAIN RELIEVER-FEVER REDUCER - **ACETAMINOPHEN**
COUGH SUPPRESSANT - **DEXTROMETHORPHAN HBr**
EXPECTORANT - **GUAIFENESIN**
NASAL DECONGESTANT - **PHENYLEPHRINE HCl**

• Headache, Fever, Sore Throat,
Minor Aches & Pains
• Nasal/Sinus Congestion
& Sinus Pressure
• Cough
• Chest Congestion

Alcohol Free
Antihistamine Free
Non-Drowsy

12 FL OZ (355 mL)

ORIGINAL FLAVOR

Drug Facts **DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: ■ sore throat ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ minor aches and pains ■ fever ■ headache ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ liver disease ■ heart disease ■ diabetes
■ thyroid disease ■ high blood pressure
■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus)

CODE AREA
* no varnish * no color
: 60340 88 F7

TOPCARE DAY TIME COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:36800-603
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)			DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)			GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)			PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL
Inactive Ingredients				
Ingredient Name				Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
GLYCERIN (UNII: PDC6A3C0OX)				
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)				
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SUCROSE (UNII: C151H8M554)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	ORANGE (clear)		Score	
Shape			Size	
Flavor	FRUIT, MENTHOL		Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-603-34	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/22/2014	02/28/2022
2	NDC:36800-603-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/13/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/13/2014	

Labeler - Topco Associates LLC (006935977)

Revised: 10/2024

Topco Associates LLC