

BENZOIN COMPOUND TINCTURE- benzoin resin liquid
Humco Holding Group, Inc.

Humco Benzoin Compound Tincture, USP

Drug Facts

Active Ingredient

Benzoin

Purpose

Oral mucosal protectant

Use

Forms a coating over wound for protecting recurring canker sores

Warnings

For external use only. Do not swallow. Do not exceed recommended dosage.

When using this product

Children under 12 years of age should be supervised in the use of this product.

Do not use for more than 7 days unless directed by a dentist or doctor.

Stop use and consult a dentist or doctor if

sore mouth symptoms do not improve in 7 days. irritation, pain or redness persists or worsens. swelling, rash or fever develops.

Keep out of reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

Adults and children 6 months of age and older: Dry the affected area, with cotton swab, apply undiluted to the affected area not more often than every 2 hours.

Children under 6 months of age: Consult a dentist or doctor.

Other information

Flammable: Keep away from spark, heat or flame.

Inactive Ingredients

Alcohol 77%, Aloe, Storax, Tolu Balsam

Principal Display Panel - 16oz

Drug Facts	
Active ingredients	Purpose
Benzoin.....	Oral mucosal protectant
Indications	
■ Forms a coating over wound for protecting recurring canker sores	
Warnings	
■ Do not exceed recommended dosage	
When using this product	
■ Do not use for more than 7 days unless directed by a dentist or doctor	
Stop use and consult a dentist or doctor promptly if	
■ sore mouth symptoms do not improve in 7 days	
■ irritation, pain or redness persists or worsens	
■ swelling, rash or fever develops	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.	
Directions	
■ Adults and children 6 months of age and older: Dry the affected area. Saturate a cotton applicator with medication. Apply the undiluted medication directly to the affected area. Do not use more often than every 2 hours.	
■ Children under 6 months of age: Consult a dentist or doctor	
Continued ►	

**BENZONIN
COMPOUND
TINCTURE**
ALCOHOL 77%

HUMCOTM
A trusted name
in home health care

*Safety Sealed

**First
Aid**

USP XIX

FIRST AID
ANTISEPTIC

Flammable:
Keep from
spark,
heat or flame.

16 FL oz
(1 pt) 473 mL

***Warning:** Do not use if tamper evident seal is broken or missing. This product is sealed with either a shrink band around cap imprinted "Sealed For Your Protection" or a foil seal under cap.

(Drug Facts continued)

Other information

■ Flammable: Keep away from spark, heat or flame.

Inactive ingredients

■ Alcohol 77%, Aloe, Storax, Tolu Balsam

A43004 10 2021



HUMCO, Texarkana, TX 75501
Questions or Comments? ☎ 1-800-662-3435
"We Help People Feel Better"

✳ Go to: www.simplehomeremedies.com
to find helpful hints, and learn more about
Benzoin Compound.

Drug Facts

Active ingredients	Purpose
Benzoin.....	Oral mucosal protectant

Use

- Forms a coating over wound for protecting recurring canker sores.

Warnings For external use only.

- Do not swallow.
- Do not exceed recommended dosage.

When using this product

- Children under 12 years of age should be supervised in the use of this product.
- Do not use for more than 7 days unless directed by a dentist or doctor.

Stop use and consult a dentist or doctor if

- sore mouth symptoms do not improve in 7 days.
- irritation, pain or redness persists or worsens.
- swelling, rash or fever develops.

Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- Adults and children 6 months of age and older: Dry the affected area, with cotton swab, apply undiluted to the affected area not more often than every 2 hours.
- Children under 6 months of age: Consult a dentist or doctor.

Continued ▶

NDC 0395-0243-16 *Safety Sealed

BENZOIN COMPOUND

TINCTURE USP ALCOHOL 77%

A Trusted Name in Home Health Care



- Oral mucosal protectant
- Wound dressing

16 FL OZ (1 PT) 473 mL

HUMCO™

***Warning:** Do not use if tamper evident seal is broken or missing. This product is sealed with either a shrink band around cap imprinted "Sealed For Your Protection" or a foil seal under cap.

(Drug Facts continued)

Other information

- Flammable: Keep away from spark, heat or flame.

Inactive ingredients

- Alcohol 77%, Aloe, Storax, Tolu Balsam

HUMCO, Texarkana, TX 75501
 Questions or Comments?
 ☎1-800-662-3435
 "We Help People Feel Better"

Go to: www.simplehomeremedies.com to find helpful hints, and learn more about Benzoin Compound.

REV.052009.RLG



3 03950 24316 0

Principal Display Panel - 2 oz

<p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredients</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Benzoin.....</td> <td>Oral mucosal protectant</td> </tr> </tbody> </table> <p>Indications ■ Forms a coating over wound for protecting recurring canker sores</p> <p>Warnings</p> <ul style="list-style-type: none"> Do not exceed recommended dosage <p>When using this product</p> <ul style="list-style-type: none"> Do not use for more than 7 days unless directed by a dentist or doctor <p>Stop use and consult a dentist or doctor promptly if</p> <ul style="list-style-type: none"> sore mouth symptoms do not improve in 7 days irritation, pain or redness persists or worsens swelling, rash or fever develops <p style="text-align: right;"><i>Continued ▶</i></p>	Active ingredients	Purpose	Benzoin.....	Oral mucosal protectant	<p>BENZOIN COMPOUND TINCTURE</p> <p>ALCOHOL 77%</p> <p>HUMCO™</p> <p>A trusted name in home health care</p>	<p>*Safety Sealed</p> <p>First Aid</p> <p>USP XIX</p> <ul style="list-style-type: none"> ORAL MUCOSAL PROTECTANT WOUND DRESSING <p>Flammable: Keep from spark, heat or flame.</p> <p>2 FL oz (59 mL)</p>	<p>*Warning: Do not use if tamper evident seal is broken or missing. This product is sealed with either a shrink band around cap imprinted "Sealed For Your Protection" or a foil seal under cap.</p> <p>Drug Facts (continued)</p> <p>Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.</p> <p>Directions</p> <ul style="list-style-type: none"> Adults and children 6 months of age and older: Dry the affected area. Saturate a cotton applicator with medication. Apply the undiluted medication directly to the affected area. Do not use more often than every 2 hours. Children under 6 months of age: Consult a dentist or doctor <p>Inactive ingredients ■ Alcohol 77%, Aloe, Storax, Tolu Balsam</p> <p>Questions or Comments? ☎1-800-662-3435</p> <p>Packed by: Humco, Texarkana, TX 75501</p>	 <p>3 03950 24392 4</p> <p>R042920</p>
Active ingredients	Purpose							
Benzoin.....	Oral mucosal protectant							

Drug Facts	
Active ingredients	Purpose
Benzoin.....	Oral mucosal protectant
Indications ■ Forms a coating over wound for protecting recurring canker sores	
Warnings ■ Do not exceed recommended dosage	
When using this product ■ Do not use for more than 7 days unless directed by a dentist or doctor	
Stop use and consult a dentist or doctor promptly if ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain or redness persists or worsens ■ swelling, rash or fever develops Continued ▶	

**BENZOIN
COMPOUND
TINCTURE**
ALCOHOL 77%

HUMCO
A trusted name
in home health care

***Safety Sealed
First
Aid**

USP XIX

- ORAL MUCOSAL PROTECTANT
- WOUND DRESSING

Flammable: Keep from spark, heat or flame.

2 FL oz (59 mL)

***Warning:** Do not use if tamper evident seal is broken or missing. This product is sealed with either a shrink band around cap imprinted "Sealed For Your Protection" or a foil seal under cap.

(Drug facts continued)
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.
Directions ■ Adults and children 6 months of age and older: Dry the affected area. Saturate a cotton applicator with medication. Apply the undiluted medication directly to the affected area. Do not use more often than every 2 hours. ■ Children under 6 months of age: Consult a dentist or doctor
Inactive ingredients ■ Alcohol 77%, Aloe, Storax, Tolu Balsam
Questions or Comments? ☎1-800-662-3435



Drug Facts	
Active ingredients	Purpose
Benzoin.....	Oral mucosal protectant
Use ■ Forms a coating over wound for protecting recurring canker sores.	
Warnings For external use only. ■ Do not swallow. ■ Do not exceed recommended dosage.	
When using this product ■ Children under 12 years of age should be supervised in the use of this product. ■ Do not use for more than 7 days unless directed by a dentist or doctor.	
Stop use and consult a dentist or doctor if ■ sore mouth symptoms do not improve in 7 days. ■ irritation, pain or redness persists or worsens. ■ swelling, rash or fever develops. Continued ▶	

NDC 0395-0243-92 ***Safety Sealed**

BENZOIN COMPOUND
TINCTURE USP ALCOHOL 77%

**A Trusted Name
in Home Health Care**

ORAL HEALTH CARE

*Oral mucosal protectant
*Wound dressing

2 FL OZ (59 mL)

Flammable: Keep away from spark, heat or flame.

HUMCO

Distributed by: HUMCO, Texarkana, TX 75501
* Go to: www.simplehomeremedies.com

***Warning:** Do not use if tamper evident seal is broken or missing. This product is sealed with either a shrink band around cap imprinted "Sealed For Your Protection" or a foil seal under cap.

(Drug facts continued)
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.
Directions ■ Adults and children 6 months of age and older: Dry the affected area, with cotton swab, apply undiluted to the affected area not more often than every 2 hours. ■ Children under 6 months of age: Consult a dentist or doctor.
Inactive ingredients ■ Alcohol 77%, Aloe, Storax, Tolu Balsam
Questions or Comments? ☎1-800-662-3435



2025 Label Update

Drug Facts	
Active ingredients	Purpose
Benzoin.....	Oral mucosal protectant
Uses Forms a coating over wound for protecting recurring canker sores	
Warnings For External Use Only. Do not swallow. Do not exceed recommended dosage.	
■ FLAMMABLE: Keep away from spark, heat & flame.	
Stop use and consult a dentist or doctor promptly if ■ the condition persists or gets worse ■ symptoms do not improve within 7 days ■ pain, irritation, swelling, redness, rash or fever develop.	
When using this product do not use in the eyes ■ do not use this product for longer than 1 week unless directed by a dentist or doctor.	
<i>Continued</i> ▶	

BENZOIN COMPOUND TINCTURE
ALCOHOL 77%



*Safety Sealed

• ORAL MUCOSAL PROTECTANT

Flammable: Keep from spark, heat or flame.

2 fl oz (59 mL)

***Warning:** Do not use if tamper evident seal is broken or missing. This product is sealed with either a shrink band around cap imprinted "Sealed For Your Protection" or a foil seal under cap.

Drug Facts (continued)	
Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately (800-222-1222).	
Directions	
■ Adults and children 6 months of age and older: Dry the affected area. Saturate a cotton applicator with medication. Apply the undiluted medication directly to the affected area. Do not use more often than every 2 hours.	
■ Children under 6 months of age: Consult a dentist or doctor	
Inactive ingredients Alcohol 77%, Aloe, Storax, Tolu Balsam	
Questions or Comments? 1-800-662-3435	
Distributed by: Humco, Austin, TX 78701	

REV1124
303950243924

BENZOIN COMPOUND TINCTURE				
benzoin resin liquid				
Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0395-0243	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	BENZOIN RESIN (UNII: GK21SBA74R) (BENZOIN RESIN - UNII:GK21SBA74R)	BENZOIN RESIN	1000 mg in 1 mL	
Inactive Ingredients				
	Ingredient Name		Strength	
	ALCOHOL (UNII: 3K9958V90M)			
	ALOE (UNII: V5VD430YW9)			
	TOLU BALSAM (UNII: TD2LE91MBE)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0395-0243-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/10/2017	
2	NDC:0395-0243-92	59 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/10/2017	
Marketing Information				

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	01/01/2008	

Labeler - Humco Holding Group, Inc. (825672884)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(0395-0243) , analysis(0395-0243) , pack(0395-0243) , label(0395-0243)

Revised: 12/2025

Humco Holding Group, Inc.