NICOTINE TRANSDERMAL SYSTEM- nicotine transdermal system patch, extended release
BluePoint Laboratories

Nicotine Transdermal System- nicotine patch, extended release

Drug Fact

Active ingredient(in each patch)

Step 1 - Nicotine, 21 mg delivered over 24 hours

Step 2 - Nicotine, 14 mg delivered over 24 hours

Step 3 - Nicotine, 7 mg delivered over 24 hours

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

• if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat.
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke **more than 10 cigarettes per day,** use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
Step 1	Step 2	Step 3
Use one 21 mg	Use one 14 mg	Use one 7 mg
patch/day	patch/day	patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- it is important to complete treatment. If you feel you need to use the patch for a

longer period to keep from smoking, talk to your health care provider.

Other information

store at 20 °C - 25 °C (68 °F - 77 °F)

Inactive ingredients

acrylate adhesive, polyester, silicone adhesive

Questions or comments?

1-877-748-1970

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivated to quit.
- 2. Use one patch daily according to directions.
- 3. It is important to complete treatment.
- 4. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
- 5. Use patch with a behavioral support program such as the one described in the enclosed booklet.
- Not for sale to persons under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

See self-help guide inside.

See bottom panel for lot number and expiration date.

For your family's protection, patches are supplied in child-resistant pouches.

Do not use if individual pouch is open or torn.

Manufactured by: AVEVA Drug Delivery Systems A DifGen Company Miramar, FL 33025

For BluePoint Laboratories

Rev. 12/23

^{*}This product is not manufactured or distributed by Dr. Reddy's Laboratories Ltd., distributor of Habitrol.

NDC: 68001-434-88

compare to the active ingredient in Habitrol ®*

Clear Nicotine Transdermal System Patch

Stop Smoking Aid

Includes Behavior Support Program with Self-Help Guide

21 mg

21 mg delivered over 24 hours

If you smoke **MORE**than 10 cigarettes per day: Start with Step 1.

If you smoke 10 or **LESS**cigarettes per day: Start with Step 2.

The full treatment course for STEP 1 is 28 patches (4 weeks); this package contains 7 patches (1 week course) only. Read the enclosed self-help guide for additional information.

STEP 1 7 Clear Patches (1 Week Kit)




NDC: 68001-434-90

STEP 1 14 Clear Patches (2 Week Kit)



NDC: 68001-434-91

STEP 1 28 Clear Patches (4 Week Kit)



PRINCIPAL DISPLAY PANEL - STEP TWO 14 mg CARTON

NDC: 68001-433-88

compare to the active ingredient in Habitrol ®*

Clear Nicotine Transdermal System Patch

Stop Smoking Aid

Includes Behavior Support Program with Self-Help Guide

14 mg

14 mg delivered over 24 hours

If you smoke MOREthan 10 cigarettes per day: Start with Step 1.

If you smoke 10 or **LESS**cigarettes per day: Start with Step 2.

The full treatment course for STEP 2 is 2 or 6 weeks (depending on how many cigarettes you smoke per day; this package contains 7 patches (1 week course) only. Read enclosed self-help guide for additional information.

STEP 2 7 Clear Patches

(1 Week Kit)



NDC: 68001-433-90

STEP 2 14 Clear Patches (2 Week Kit)



PRINCIPAL DISPLAY PANEL - STEP THREE 7 mg CARTON

NDC: 68001-432-88

compare to the active ingredient in Habitrol ®*

Clear Nicotine Transdermal System Patch

Stop Smoking Aid

Includes Behavior Support Program with Self-Help Guide

7 mg

21 mg delivered over 24 hours

For use after completing step 2.


The full treatment course for STEP 3 is 14 patches (2 weeks); this package contains 7 patches (1 week course) only. Read the enclosed self-help guide for additional information.

STEP 3 7 Clear Patches (1 Week Kit)



NDC: 68001-432-90

STEP 3 14 Clear Patches (2 Week Kit)



NICOTINE TRANSDERMAL SYSTEM

nicotine transdermal system patch, extended release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68001-434

Route of Administration TRANSDERMAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R) NICOTINE 21 mg in 24 h

Inactive Ingredients

Ingredient Name Strength

DIMETHICONE (UNII: 92RU3N3Y10)

-	7 al '	_		_
Pa	ч	VО	м	ч
			2	 •

Item Code Package Description Marketing Start Date Date

1	NDC:68001-434- 88	7 in 1 CARTON	06/27/2024	
1		1 h in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:68001-434- 90	14 in 1 CARTON	06/27/2024	
2		1 h in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:68001-434- 91	28 in 1 CARTON	06/27/2024	
3		1 h in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
ANDA	ANDA074612	06/27/2024		

NICOTINE TRANSDERMAL SYSTEM

nicotine transdermal system patch, extended release

Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68001-432		
Route of Administration	TRANS DERMAL				

Active Ingredient/Active Moiety					
Ingredient Name	Basis of Strength	Strength			
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	7 mg in 24 h			

Inactive Ingredients	
Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68001-432- 88	7 in 1 CARTON	06/27/2024	
1		1 h in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:68001-432- 90	14 in 1 CARTON	06/27/2024	
2		1 h in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information				
Marketing Application Number or Monograph Marketing Start Marketing End Category Citation Date Date				
ANDA	ANDA074612	06/27/2024		

NICOTINE TRANSDERMAL SYSTEM

nicotine transdermal system patch, extended release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68001-433

Route of Administration TRANSDERMAL

Active Ingredient/Active Moiety

Ingredient Name

Basis of Strength

NICOTINE (LINIU: 6M3C907Y6P) (NICOTINE LINIU: 6M3C907Y6P)

14 mg in 24 h

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)

NICOTINE

14 mg in 24 h

Inactive Ingredients

Ingredient Name Strength

DIMETHICONE (UNII: 92RU3N3Y10)

Packaging Marketing Start Marketing End Item Code **Package Description Date** Date NDC:68001-433-1 7 in 1 CARTON 06/27/2024 88 1 h in 1 POUCH; Type 0: Not a Combination 1 Product NDC:68001-433-2 14 in 1 CARTON 06/27/2024 1 h in 1 POUCH; Type 0: Not a Combination 2 Product

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA074612	06/27/2024		

Labeler - BluePoint Laboratories (985523874)

Revised: 3/2025 BluePoint Laboratories