UP AND UP 12 HOUR COUGH RELIEF- dextromethorphan polistirex suspension Target Corporation

Target Corporation 12-Hour Cough Relief Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake bottle well before use
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24
	hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24
	hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24
	hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store at 20° to 25° C (68° to 77° F)
- dosing cup provided

Inactive Ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions?

Call 1-888-547-7400

Package/Label Principal Display Panel

Compare to active ingredient in Delsym®

12-hour cough relief

dextromethorphan polistirex extended-release oral suspension cough suppressant contains sodium metabisulfite, a sulfite that may cause allergic-type reactions dosing cup included

day or night

alcohol free

12 HOUR

ORANGE FLAVORED LIQUID

5 FL OZ (148 mL)



Drug Facts Active ingredient (in each 5 mL) Purpose Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide...

Uses temporarily relieves
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains codium metablisulfite, a sulfite that may cause allergic-type reactions.

allergic-type reactions.

Ask a doctor before use if you have

— chronic ough that lasts as occurs with smoking, asthma or emphysema

— cough that occurs with too much phlegm (mucus)

■ cough that occurs with too much prilegin (mucus)

Stop use and ask a doctor if
■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
■ cough lasts more than 7 days, cough comes back, or occurs with fever, main or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdoop, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake bottle well before use
 measure only with dosing cup provided
 do not use dosing cup with other products
 dose as follows or as directed by a doctor

adults and children	10 mL every 12 hours,
12 years of age and over	not to exceed 20 mL in 24 hours
children 6 to under	5 mL every 12 hours,
12 years of age	not to exceed 10 mL in 24 hours
children 4 to under	2.5 mL every 12 hours,
6 years of age	not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

■ each 5 mL contains: sodium 5 mg
■ store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

Inactive ingredients D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose com syrup, methylgaraben, natural and artificial orange laws, polysorbate 80, polyving lacette, povidene, propylgaraben, purified water, sodium metabisulfite, sodium polystyrene sultonate, sucrose, tartaria cald, tagacarthi gum, triacetin, xarihtan gum

Questions? Call 1-888-547-7400



DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided.
Do not use dosing cup with other products

Age (yr)	Dose
12 years	10 mL
to adult	EVERY 12 HOURS
6 to	5 mL
under 12	EVERY 12 HOURS
4 to	2.5 mL
under 6	EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.



Dosing cup included

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Delsym®.

GLUTEN FREE

094 14 0272 R03 ID275438 Distributed by Target Corporation Minneapolis, MN 55403 ©2020 Target Brands, Inc. Shop Target.com





5 FL 0Z (148 mL)



Compare to active ingredient in Delsym®*

12-hour cough relief

extended-release oral suspension cough suppressant

contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

day or night





dextromethorphan polistirex extended-release oral suspension cough suppressant











UP AND UP 12 HOUR COUGH RELIEF

dextromethorphan polistirex suspension

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:11673-114

ORAL **Route of Administration**

Active Ingredient/Active Moiety

Basis of Strength Ingredient Name Strength

Inactive Ingredients	
Ingredient Name	Strength
POLISTIREX (UNII: 5H9W9GTW27)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
METHYLPARABEN (UNII: A218 C7HI9 T)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ACETATE (UNII: 32K497ZK2U)	
PO VIDO NE, UNSPECIFIED (UNII: FZ989 GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)	
SUCROSE (UNII: C151H8 M554)	
TARTARIC ACID (UNII: W4888I119H)	
TRAGACANTH (UNII: 2944357O2O)	
TRIACETIN (UNII: XHX3C3X673)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-114-21	1 in 1 CARTON	10/23/2012	10/01/2016
1		89 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-114-28	1 in 1 CARTON	03/03/2014	
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	10/23/2012	

Revised: 5/2020 Target Corporation