

**IBUPROFEN (NSAID) PAIN RELIEVER/ FEVER REDUCER- ibuprofen tablet**  
**Amneal Pharmaceuticals**

-----  
**Ibuprofen Caplets, USP**  
**Fever reducer/ Pain Reliever (NSAID)**

***Drug Facts***

**ACTIVE INGREDIENT**

***(in each caplet)***

Ibuprofen USP, 200 mg (NSAID)\*\*

\*\*nonsteroidal anti-inflammatory drug

**PURPOSE**

Pain reliever/fever reducer

**INDICATIONS AND USAGE**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

**WARNINGS**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma(wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen,

or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

#### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug
- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug

#### **When using this product**

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

#### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

#### **KEEP OUT OF REACH OF CHILDREN**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### **DOSAGE AND ADMINISTRATION**

- **do not take more than directed**
  - **the smallest effective dose should be used**
-

Adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

#### **OTHER INFORMATION**

- Store between 20 - 25°C (68 - 77°F)
- Read all warnings and directions before use

#### **INACTIVE INGREDIENTS**

Brown Caplets: Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Hypromellose, Iron Oxide Red, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Corn Starch Corn, Titanium Dioxide

Orange Caplets: Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, FD&C Yellow #6, Hypromellose, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Corn Starch, Titanium Dioxide

#### **Questions or Comments**

Call 1-877-835-5472

Monday through Friday 9AM - 5PM EST.

\*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Advil®

Distributed By:

**Ameal Pharmaceuticals**

Glasgow, KY 42141

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

\*Compare to the active ingredient in Advil®

NDC 53746-142-10

SEE NEW WARNINGS INFORMATION

# Ibuprofen Tablets, USP

200 mg

Important: Read all product information before using.

IP-142

Pain reliever / Fever reducer (NSAID) **1000 CAPLETS**



BROWN FILM-COATED (Capsule-shaped Tablets)

**Drug Facts**  
**Active ingredient (in each caplet)** Ibuprofen USP, 200 mg (NSAID)..... Pain reliever/fever reducer  
\*Nonsteroidal anti-inflammatory drug

**Uses**  
• temporarily relieves minor aches and pains due to:  
• headache • muscle aches  
• minor pain of arthritis • toothache • backaches • the common cold  
• menstrual cramps • temporarily reduces fever

**Warnings**  
**Always alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
• hives • facial swelling  
• asthma/wheezing • shock • skin redness • rash • blisters • itchy allergic reactions; stop use and seek medical help right away.

**Stomach bleeding warning:** This product causes a nonsteroidal anti-inflammatory (NSAID) effect, which may cause stomach bleeding. The chances are higher if you:  
• are age 60 or older • have had stomach bleeding or bleeding problems • take a blood thinning (anticoagulant) or aspirin drug • take other drugs containing aspirin or nonsteroidal NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Do not use:** • if you have ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery  
**Ask a doctor before use if:** • you have problems or serious side effects from taking pain relievers or fever reducers • the stomach bleeding warning applies to you • you have a history of stomach problems such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you have asthma • you are taking a diuretic

**Ask a doctor or pharmacist before use if you are:** • taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin • under a doctor's care for any serious condition • using any other drug

Do not use if inner seal imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.  
\*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Advil®.  
Rev. 01-2011  
Distributed by: Amneal Pharmaceuticals  
Glasgow, KY 42141



N 3 53746-142-10 8

Lot No: 3/4"  
Exp. Date: Non-Varmish Area

PEEL BACK FOR ADDITIONAL DRUG FACTS

### Drug Facts (continued)

**When using this product:** • take with food or milk if stomach upset occurs • the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.  
**Stop use and ask a doctor if:** • you experience any of the following signs of stomach bleeding:  
• black stools • dark stool • vomit blood • vomit bloody or black stool  
• take stomach pain medicine not get better  
• you get worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present in the painful area • any new symptoms appear  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important to use ibuprofen during the 3rd trimester of pregnancy because it may affect the fetus. Do not use if you are breastfeeding. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
• do not take more than directed  
• the smallest effective dose should be used

**Adults and Children 12 years and older**  
• take 1 caplet every 4 to 6 hours while symptoms persist  
• if pain or fever does not respond to 1 caplet, 2 caplets may be taken in 24 hours, unless directed by a doctor  
• ask a doctor

**Other information**  
• Store between 20° to 25°C (68° to 77°F) • Read all warnings and directions before use

**Inactive ingredients**  
Anhydrous Lactose, Cantharal Wax, Colloidal Silicon Dioxide, Corn Starch, Hydroxypropyl Methylcellulose, Hydroxypropyl Starch, Magnesium Stearate, Polyethylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Polyethylene Glycol, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Titanium Dioxide

**Questions or Comments?**  
Call 1-877-835-5472  
Monday through Friday, 9AM - 5PM EST

PERMANENT GLUE

\*Compare to the active ingredient in Motrin BP®

NDC 53746-144-10

SEE NEW WARNINGS INFORMATION

# Ibuprofen Tablets, USP

200 mg

Important: Read all product information before using.

IP-144

Pain reliever / Fever reducer (NSAID) **1000 CAPLETS**



ORANGE FILM-COATED (Capsule-shaped Tablets)

**Drug Facts**  
**Active ingredient (in each caplet)** Ibuprofen USP, 200 mg (NSAID)..... Pain reliever/fever reducer  
\*Nonsteroidal anti-inflammatory drug

**Uses**  
• temporarily relieves minor aches and pains due to:  
• headache • muscle aches  
• minor pain of arthritis • toothache • backache • the common cold  
• menstrual cramps • temporarily reduces fever

**Warnings**  
**Always alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
• hives • facial swelling  
• asthma/wheezing • shock • skin redness • rash • blisters • itchy allergic reactions; stop use and seek medical help right away.

**Stomach bleeding warning:** This product causes a nonsteroidal anti-inflammatory (NSAID) effect, which may cause stomach bleeding. The chances are higher if you:  
• are age 60 or older • have had stomach bleeding or bleeding problems • take a blood thinning (anticoagulant) or aspirin drug • take other drugs containing aspirin or nonsteroidal NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Do not use:** • if you have ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery  
**Ask a doctor before use if:** • you have problems or serious side effects from taking pain relievers or fever reducers • the stomach bleeding warning applies to you • you have a history of stomach problems such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you have asthma • you are taking a diuretic

**Ask a doctor or pharmacist before use if you are:** • taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin • under a doctor's care for any serious condition • using any other drug

Do not use if inner seal imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.  
\*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Motrin BP®.  
Rev. 01-2011  
Distributed by: Amneal Pharmaceuticals  
Glasgow, KY 42141



N 3 53746-144-10 2

Lot No: 3/4"  
Exp. Date: Non-Varmish Area

PEEL BACK FOR ADDITIONAL DRUG FACTS

Exp. Date:

Lot No.:



Important: Read all product information before using.

Compare to the active ingredient in Advil®

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

NDC 53746-142-24

SEE NEW WARNINGS INFORMATION

# Ibuprofen Tablets, USP

## 200 mg

PAIN RELIEVER/  
FEVER REDUCER (NSAID)

IP 142

BROWN FILM-COATED  
(Capsule-shaped Tablets)  
**24 CAPLETS**

### Drug Facts (continued)

#### Questions or Comments?

Call 1-877-835-5472

Monday through Friday 9AM - 5PM EST.

Distributed by: **Amneal Pharmaceuticals**  
Glasgow, KY 42141

\*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Advil®.

Rev. 01-2011

**Drug Facts (continued)**

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
- the smallest effective dose should be used

Adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

**Other information**

- Store between 20° to 25°C (68° to 77°F)
- Read all warnings and directions before use

**Inactive ingredients**

Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Corn Starch, Hypromellose, Iron Oxide Red, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Titanium Dioxide

**Drug Facts (continued)**

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear

**Drug Facts**

**Active ingredient (in each caplet)** Purpose  
Ibuprofen USP 200 mg (NSAID)\*\* Pain reliever/  
fever reducer  
\*\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
  - temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Drug Facts (continued)**

**When using this product**

- take 1 caplet every 4 to 6 hours while symptoms persist
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- ask a doctor

**Directions**

- do not take more than directed
- the smallest effective dose should be used

Children under 12 years

- ask a doctor

**Other information**

- Store between 20° to 25°C (68° to 77°F)
- Read all warnings and directions before use

**Inactive ingredients**

Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Corn Starch, EPOC, Yellow 46 Hypromellose, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Titanium Dioxide

**Questions or Comments?**

Call 1-877-835-5472  
Monday through Friday 9AM - 5PM EST.

PERMANENT  
GLUE

Exp. Date:

Lot No.:



NDC 53746-144-24

SEE NEW WARNINGS INFORMATION

# Ibuprofen Tablets, USP

## 200 mg

IP 144

ORANGE FILM-COATED  
(Capsule-shaped Tablets)  
**24 CAPLETS**

**Important:** Read all product information before using.

Compare to the active ingredient in **Motrin IB®**

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

**PAIN RELIEVER/  
FEVER REDUCER (NSAID)**

**Drug Facts** (continued)  
**Questions or Comments?**  
Call 1-877-835-5472  
Monday through Friday 9AM - 5PM EST.

Distributed by: **Amneal Pharmaceuticals**  
Glasgow, KY 42141

\*Amneal Pharmaceuticals is not affiliated with the owner of the trademark Motrin IB®.

Rev. 01-2011

**Drug Facts** (continued)

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- do not take more than directed
- the smallest effective dose should be used

Adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

**Other information**

- Store between 20° to 25°C (68° to 77°F)
- Read all warnings and directions before use

**Inactive ingredients**

Anhydrous Lactose, Carnauba Wax, Colloidal Silicon Dioxide, Corn Starch, FD&C Yellow #6, Hypromellose, Magnesium Stearate, Microcrystalline Cellulose, Polydextrose, Polyethylene Glycol, Povidone, Sodium Lauryl Sulfate, Sodium Starch Glycolate, Titanium Dioxide

**Drug Facts** (continued)

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- if you use more than directed or for longer than directed.

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear

**Drug Facts**

**Active ingredient (in each caplet)** **Purpose**  
Ibuprofen USP, 200 mg (NSAID)\*\* .....: pain reliever/fever reducer  
\*\*nonsteroidal anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - muscular aches
  - toothache
  - the common cold
- minor pain or arthritis
- backache
- menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

## IBUPROFEN (NSAID) PAIN RELIEVER/ FEVER REDUCER

ibuprofen tablet

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:53746-144

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	ORANGE	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	IP;144
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53746-144-24	24 in 1 BOTTLE		
2	NDC:53746-144-01	100 in 1 BOTTLE		
3	NDC:53746-144-10	1000 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072199	12/16/2009	

## IBUPROFEN (NSAID) PAIN RELIEVER/ FEVER REDUCER

ibuprofen tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53746-142
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

**Product Characteristics**

Color	BROWN	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	IP;142
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53746-142-24	24 in 1 BOTTLE		
2	NDC:53746-142-01	100 in 1 BOTTLE		
3	NDC:53746-142-10	1000 in 1 BOTTLE		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072199	12/16/2009	

**Labeler** - Amneal Pharmaceuticals (123797875)**Registrant** - Amneal Pharmaceuticals (123797875)

## Establishment

Name	Address	ID/FEI	Business Operations
Amneal Pharmaceuticals		831227801	ANALYSIS, MANUFACTURE, LABEL, PACK

Revised: 2/2012

Amneal Pharmaceuticals