

NUVICARE HIVES RELIEF- fexofenadine hydrochloride tablet

NUVICARE LLC

Nivicare Hives Relief

Active Ingredient (in each caplet)

Fexofenadine HCl 180 mg

Purpose

Antihistamine

Uses

reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.

Warnings

Severe Allergy Warning:

Get emergency help immediately if you have hives along with any of the following symptoms:

- trouble swallowing
- dizziness or loss of consciousness
- swelling of tongue
- swelling in or around mouth
- trouble speaking
- drooling
- wheezing or problems breathing

These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.

Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for "anaphylaxis or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.

Do not use

- to prevent hives from any known cause such as:
 - foods ■ insect stings ■ medicines ■ latex or rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious.
- If you do not know the cause of your hives, see your doctor for a medical exam. Your

doctor may be able to help you find a cause.

■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ kidney disease. Your doctor should determine if you need a different dose. ■ hives that are an unusual color, look bruised or blistered ■ hives that do not itch

When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve after 3 days of treatment
- the hives have lasted more than 6 weeks

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. in case of overdose, get medical help or contact a poison control center right away.

Directions

adults and children 12 years of age and over: take one 180mg tablet with water once a day; do not take more than 1 tablet in 24 hours

- children under 12 years of age: do not use
- adults 65 years of age and older: ask a doctor
- consumers with kidney disease: ask a doctor

Other information

- Store between 20°C and 25 °C (68 °and 77°F)
- Protect from excessive moisture

Inactive ingredient

Colloidal Silicon Dioxide, Croscarmellose Sodium Hypromellose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol/macrogol, Povidone, Pregelatinized Starch, Red iron oxide, Silica, Titanium dioxide, Yellow iron oxide.

Call 1 (718) 337-8733 or email: support@nuvicare.com

NDC 84324-03 1-01 antihistamine NON-DROWSY Hives Relief Fexofenadine HCl 180 mg ✓ Reduces Hives ✓ Relieves Itching Due to Hives 30 caplets not actual size 24 HOURS * compare to Allegra® Hives 24HR active ingredients. CERTIFIED ORIGINAL PRES	TAMPER EVIDENT Do not use if safety seal under cap is broken or missing.	<table border="1"> <thead> <tr> <th>Drug Facts</th> </tr> </thead> <tbody> <tr> <td> Active Ingredient (in each caplet) Fexofenadine HCl 180 mg (equivalent to Allegra® 180 mg tablets) </td> </tr> <tr> <td> Purpose Antihistamine </td> </tr> <tr> <td> Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring. </td> </tr> <tr> <td> Warnings Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: ■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ drooling ■ wheezing or problems breathing These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later. </td> </tr> </tbody> </table>	Drug Facts	Active Ingredient (in each caplet) Fexofenadine HCl 180 mg (equivalent to Allegra® 180 mg tablets)	Purpose Antihistamine	Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.	Warnings Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: ■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ drooling ■ wheezing or problems breathing These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.	(continued under label) Distributed by: Nuvicare LLC., Brooklyn, NY 11211, U.S.A. www.nuvicare.com PL0281-00 Lot No. Exp. Dt. PEEL HERE
Drug Facts								
Active Ingredient (in each caplet) Fexofenadine HCl 180 mg (equivalent to Allegra® 180 mg tablets)								
Purpose Antihistamine								
Uses reduces hives and relieves itching due to hives (urticaria). This product will not prevent hives or an allergic skin reaction from occurring.								
Warnings Severe Allergy Warning: Get emergency help immediately if you have hives along with any of the following symptoms: ■ trouble swallowing ■ dizziness or loss of consciousness ■ swelling of tongue ■ swelling in or around mouth ■ trouble speaking ■ drooling ■ wheezing or problems breathing These symptoms may be signs of anaphylactic shock. This condition can be life threatening if not treated by a health professional immediately. Symptoms of anaphylactic shock may occur when hives first appear or up to a few hours later.								

Drug Facts (cont)
Not a Substitute for Epinephrine. If your doctor has prescribed an epinephrine injector for anaphylaxis or severe allergy symptoms that could occur with your hives, never use this product as a substitute for the epinephrine injector. If you have been prescribed an epinephrine injector, you should carry it with you at all times.
Do not use ■ to prevent hives from any known cause such as: ■ foods ■ insect stings ■ medicines ■ latex ■ rubber gloves because this product will not stop hives from occurring. Avoiding the cause of your hives is the only way to prevent them. Hives can sometimes be serious.
If you do not know the cause of your hives, see your doctor for a medical exam. Your doctor may be able to help you find a cause. ■ If you have ever had an allergic reaction to this product or any of its ingredients
Ask a doctor before use if you have ■ kidney disease. Your doctor should determine if you need a different dose. ■ Hives that are an unusual color, look bruised or blistered ■ Hives that do not itch
When using this product ■ do not take more than directed ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions)
Stop use and ask a doctor if ■ an allergic reaction to this product occurs. Seek medical help right away. ■ symptoms do not improve after 3 days of treatment ■ the hives have lasted more than 6 weeks
If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a poison control center right away.
Directions: adults and children 12 years of age and over: take one 180mg tablet with water once a day, do not take more than 1 tablet in 24 hours ■ children under 12 years of age: do not use ■ adults 65 years of age and older: ask a doctor ■ consumers with kidney disease: ask a doctor
Other information: ■ Store between 20°C and 25 °C (68 ° and 77°F) ■ Protect from excessive moisture
Inactive ingredient: Colloidal Silicon Dioxide, Crosscarmellose Sodium Hygromellose, Glycol/Macrogel, Microcrystalline Cellulose, Polyethylene Glycol/Macrogel, Povidone, Pregelatinized Starch, Red iron oxide, Silica, Titanium dioxide, Yellow iron oxide.
Questions or comments: Call 1 (718) 337-8733 or email: support@nivicare.com
† This product is not manufactured or distributed by Chatten, Inc., a Sanofi Company, owner of the registered trademark Allegra®.

NUVICARE HIVES RELIEF

fexofenadine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84324-031
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FEXOFENADINE HYDROCHLORIDE (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	180 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	orange	Score	no score
Shape	capsule	Size	17mm
Flavor		Imprint Code	180
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84324-031-01	30 in 1 BOTTLE; Type 0: Not a Combination Product	01/21/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210137	01/21/2026	

Labeler - NUVICARE LLC (119257565)

Revised: 1/2026

NUVICARE LLC