ACETAMINOPHEN- acetaminophen tablet NuCare Pharmaceuticals, Inc.

Major Pharmaceuticals Acetaminophen Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user has ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user

- has liver disease
- is a child with pain of arthritis

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

• do not take more than directed (see overdose warning)

adults and children 12 years and over	 take 2 tablets every 4 to 6 hours while symptoms last do not take more than 10 tablets in 24 hours do not use for more than 10 days unless directed by a doctor
children 6-11 years	 take 1 tablet every 4 to 6 hours while symptoms last do not take more than 5 tablets in 24 hours do not use for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Inactive ingredients

croscarmellose sodium*, povidone, pregelatinized starch, stearic acid
*may contain this ingredient

Questions or comments?

Principal Display Panel



ACETAMINOPHEN

acetaminophen tablet

Droduct	Information
PICKLICI	iniormalkon

Product Type HUMAN OTC DRUG Item Code (Source) NDC:68071-2201(NDC:0904-6719)

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) ACETAMINOPHEN 325 mg

Inactive Ingredients

Ingredient Name Strength

CROSCARMELLOSE SODIUM (UNII: M280L1HH48)

POVIDONE (UNII: FZ 989GH94E)
STEARIC ACID (UNII: 4ELV7Z 65AP)

Produ		L		
Produ	CT	nara	CTEI	ISTICS

Floudet characteristics			
Color	white	Score	no score
Shape	ROUND (beveled edge)	Size	10mm
Flavor		Imprint Code	325MG;L403
Contains			

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:68071- 2201-3	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/26/2020	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/18/2018	

Labeler - NuCare Pharmaceuticals,Inc. (010632300)

Establishment				
Name	Address	ID/FEI	Business Operations	
NuCare Pharmaceuticals, Inc.		010632300	repack(68071-2201)	

Revised: 7/2024 NuCare Pharmaceuticals,Inc.