

**SIGNATURE CARE PAIN RELIEF- naproxen sodium tablet, film coated  
Safeway**

-----  
**Better Living Brands LLC Pain Relief Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - it feels like the pill is stuck in your throat
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **do not take more than directed**
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older:

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years:

- ask a doctor

**Other information**

- **each caplet contains:** sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?**

**1-888-723-3929**

**Principal Display Panel**

Pain Relief

NAPROXEN SODIUM TABLETS 220 mg

Pain Reliever/Fever Reducer (NSAID)

Strength to last 12 hours

Actual Size

COMPARE TO Aleve<sup>®</sup> Caplets active ingredient

50 CAPLETS\*\*

\*\*Capsule-shaped tablets

Actual Size



**OUR PROMISE**  
QUALITY & SATISFACTION  
**100% GUARANTEED**  
OR YOUR MONEY BACK.

**DO NOT USE IF PRINTED FOIL UNDER  
CAP IS BROKEN OR MISSING**

**Pain Relief**  
NAPROXEN SODIUM TABLETS  
220 mg  
Pain Reliever/Fever Reducer  
(NSAID)  
Strength to last 12 hours  
50 CAPLETS\* - Capsule-shaped  
tablets

**DO NOT USE IF PRINTED FOIL UNDER  
CAP IS BROKEN OR MISSING  
SEE CARTON FOR FULL INFORMATION**

NDC 21130-444-71



Quality Guaranteed

# Pain Relief

NAPROXEN SODIUM TABLETS

220 mg  
Pain Reliever/Fever Reducer  
(NSAID)

Strength to last 12 hours





Actual  
Size

**50 CAPLETS\*\***

\*\*Capsule-shaped tablets

**COMPARE TO**  
Aleve® Caplets  
active ingredient\*\*\*



EXP. DATE :

EXP.

LOT NO.



**Drug Facts**

**Active ingredient (in each caplet)** Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)\*  
**Purposes** Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ minor pain of arthritis ■ muscular aches ■ backache  
■ menstrual cramps ■ headache ■ toothache  
■ the common cold ■ temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you: ■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition  
■ taking any other drug

**Drug Facts (continued)**

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding: ■ feel faint

- vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

**Adults and children 12 years and older:**

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

**Children under 12 years:** ■ ask a doctor

**Other information**

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Drug Facts** (continued)

**Inactive ingredients:** F D&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?** 1-888-723-3929

**KEEP CARTON FOR REFERENCE**

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve® Capslets.  
**DISTRIBUTED BY:**  
**BETTER LIVING BRANDS LLC**  
 P.O. BOX 99, PLEASANTON, CA 94566-0009  
 1-888-723-3929  
 www.betterlivingbrandsllc.com

## SIGNATURE CARE PAIN RELIEF

naproxen sodium tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-444
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	BLUE (Light Blue)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (oval)	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	L368
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-444-71	1 in 1 CARTON	10/02/2015	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-444-78	1 in 1 CARTON	09/28/2015	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:21130-444-79	400 in 1 BOTTLE; Type 0: Not a Combination Product	09/04/2015	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	09/04/2015	

**Labeler** - Safeway (009137209)

Revised: 12/2019

Safeway