

**POLYMYXIN B - polymyxin b injection, powder, lyophilized, for solution**  
**Eugia US LLC**

-----

**Polymyxin B for Injection, USP**

**500,000 Units per Vial**

**Rx only**

To reduce the development of drug-resistant bacteria and maintain the effectiveness of polymyxin B and other antibacterial drugs, polymyxin B should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

**WARNING**

**CAUTION:** WHEN THIS DRUG IS GIVEN INTRAMUSCULARLY, INTRAVENOUSLY AND/OR INTRATHECALLY, IT SHOULD BE GIVEN ONLY TO HOSPITALIZED PATIENTS, SO AS TO PROVIDE CONSTANT SUPERVISION BY A PHYSICIAN.

RENAL FUNCTION SHOULD BE CAREFULLY DETERMINED AND PATIENTS WITH RENAL DAMAGE AND NITROGEN RETENTION SHOULD HAVE REDUCED DOSAGE. PATIENTS WITH NEPHROTOXICITY DUE TO POLYMYXIN B SULFATE USUALLY SHOW ALBUMINURIA, CELLULAR CASTS, AND AZOTEMIA. DIMINISHING URINE OUTPUT AND A RISING BUN ARE INDICATIONS FOR DISCONTINUING THERAPY WITH THIS DRUG.

NEUROTOXIC REACTIONS MAY BE MANIFESTED BY IRRITABILITY, WEAKNESS, DROWSINESS, ATAXIA, PERIORAL PARESTHESIA, NUMBNESS OF THE EXTREMITIES, AND BLURRING OF VISION. THESE ARE USUALLY ASSOCIATED WITH HIGH SERUM LEVELS FOUND IN PATIENTS WITH IMPAIRED RENAL FUNCTION AND/OR NEPHROTOXICITY.


THE CONCURRENT OR SEQUENTIAL USE OF OTHER NEUROTOXIC AND/OR NEPHROTOXIC DRUGS WITH POLYMYXIN B SULFATE, PARTICULARLY BACITRACIN, STREPTOMYCIN, NEOMYCIN, KANAMYCIN, GENTAMICIN, TOBRAMYCIN, AMIKACIN, CEPHALORIDINE, PAROMOMYCIN, VIOMYCIN, AND COLISTIN SHOULD BE AVOIDED.

THE NEUROTOXICITY OF POLYMYXIN B SULFATE CAN RESULT IN RESPIRATORY PARALYSIS FROM NEUROMUSCULAR BLOCKADE, ESPECIALLY WHEN THE DRUG IS GIVEN SOON AFTER ANESTHESIA AND/OR MUSCLE RELAXANTS.

USAGE IN PREGNANCY: THE SAFETY OF THIS DRUG IN HUMAN PREGNANCY HAS NOT BEEN ESTABLISHED.

**DESCRIPTION**

Polymyxin B for Injection, USP is one of a group of basic polypeptide antibiotics derived from *B. polymyxa* (*B. aerospirous*). Polymyxin B sulfate is the sulfate salt of Polymyxins B<sub>1</sub> and B<sub>2</sub>, which are produced by the growth of *Bacillus polymyxa* (Prazmowski) Migula (Fam. Bacillacea). Polymyxin B sulfate, USP is a white or off-white powder. It has a potency of not less than 6,000 polymyxin B units per mg, calculated on the anhydrous basis. The structural formulae are:



Each vial contains 500,000 polymyxin B units for parenteral or ophthalmic administration.

Polymyxin B for Injection, USP is a sterile, white lyophilized cake or powder, suitable for preparation of sterile solutions for intramuscular, intravenous drip, intrathecal, or ophthalmic use.

In the medical literature, dosages have frequently been given in terms of equivalent weights of pure polymyxin B base. Each milligram of pure polymyxin B base is equivalent to 10,000 units of polymyxin B and each microgram of pure polymyxin B base is equivalent to 10 units of polymyxin B.

Aqueous solutions of polymyxin B sulfate may be stored up to 12 months without significant loss of potency if kept under refrigeration. In the interest of safety, solutions for parenteral use should be stored under refrigeration and any unused portion should be discarded after 72 hours. Polymyxin B sulfate should not be stored in alkaline solutions since they are less stable.

## CLINICAL PHARMACOLOGY

Polymyxin B sulfate has a bactericidal action against almost all gram-negative bacilli except the *Proteus* group. Polymyxins increase the permeability of the bacterial cell membrane leading to death of the cell. All gram-positive bacteria, fungi, and the gram-negative cocci are resistant to polymyxin B.

Polymyxin B sulfate is not absorbed from the normal alimentary tract. Since the drug loses 50 percent of its activity in the presence of serum, active blood levels are low. Repeated injections may give a cumulative effect. Levels tend to be higher in infants and children. The drug is excreted slowly by the kidneys. Tissue diffusion is poor and the drug does not pass the blood brain barrier into the cerebrospinal fluid. In therapeutic dosage, polymyxin B sulfate causes some nephrotoxicity with tubule damage to a slight

degree.

## Susceptibility Testing

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

## **INDICATIONS AND USAGE**

*Acute Infections Caused by Susceptible Strains of Pseudomonas aeruginosa.*

Polymyxin B sulfate is a drug of choice in the treatment of infections of the urinary tract, meninges, and bloodstream caused by susceptible strains of *Ps. aeruginosa*. It may also be used topically and subconjunctivally in the treatment of infections of the eye caused by susceptible strains of *Ps. aeruginosa*.

It may be indicated in serious infections caused by susceptible strains of the following organisms, when less potentially toxic drugs are ineffective or contraindicated:

*H influenzae*, specifically meningeal infections.

*Escherichia coli*, specifically urinary tract infections.

*Aerobacter aerogenes*, specifically bacteremia.

*Klebsiella pneumoniae*, specifically bacteremia.

**NOTE: IN MENINGEAL INFECTIONS, POLYMYXIN B SULFATE SHOULD BE ADMINISTERED ONLY BY THE INTRATHECAL ROUTE.**

To reduce the development of drug-resistant bacteria and maintain the effectiveness of polymyxin B and other antibacterial drugs, polymyxin B should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

## **CONTRAINDICATIONS**

This drug is contraindicated in persons with a prior history of hypersensitivity reactions to polymyxins.

## **WARNINGS**

*Clostridium difficile* associated diarrhea (CDAD) has been reported with use of nearly all

antibacterial agents, including Polymyxin B for Injection, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

*C. difficile* produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

## **PRECAUTIONS**

### **General**

Prescribing polymyxin B in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

See **WARNING** box.

Baseline renal function should be done prior to therapy, with frequent monitoring of renal function and blood levels of the drug during parenteral therapy.

Avoid concurrent use of a curariform muscle relaxant and other neurotoxic drugs (ether, tubocurarine, succinylcholine, gallamine, decamethonium and sodium citrate) which may precipitate respiratory depression. If signs of respiratory paralysis appear, respiration should be assisted as required, and the drug discontinued.

As with other antibiotics, use of this drug may result in overgrowth of nonsusceptible organisms, including fungi.

If superinfection occurs, appropriate therapy should be instituted.

### **Information for Patients**

Patients should be counseled that antibacterial drugs including polymyxin B should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When polymyxin B is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by

polymyxin B or other antibacterial drugs in the future.

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

## **ADVERSE REACTIONS**

See **WARNING** box.

**Nephrotoxic reactions:** Albuminuria, cylinduria, azotemia, and rising blood levels without any increase in dosage.

**Neurotoxic reactions:** Facial flushing, dizziness progressing to ataxia, drowsiness, peripheral paresthesias (circumoral and stocking glove), apnea due to concurrent use of curariform muscle relaxants, other neurotoxic drugs or inadvertent overdose, and signs of meningeal irritation with intrathecal administration, e.g., fever, headache, stiff neck and increased cell count and protein cerebrospinal fluid.

**Other reactions occasionally reported:** Drug fever, urticarial rash, pain (severe) at intramuscular injection sites, and thrombophlebitis at intravenous injection sites.

To report SUSPECTED ADVERSE EVENTS, contact FDA at 1-800-FDA-1088 or [www.fda.gov](http://www.fda.gov).

## **DOSAGE AND ADMINISTRATION**

### **PARENTERAL:**

#### **Intravenous**

Dissolve 500,000 polymyxin B units in 300 to 500 mL solutions for parenteral dextrose injection 5% for continuous drip.

#### **Adults and children**

15,000 to 25,000 units/kg body weight/day in individuals with normal kidney function. This amount should be reduced from 15,000 units/kg downward for individuals with kidney impairment. Infusions may be given every 12 hours; however, the total daily dose must not exceed 25,000 units/kg/day.

#### **Infants**

Infants with normal kidney function may receive up to 40,000 units/kg/day without adverse effects.

## **Intramuscular**

Not recommended routinely because of severe pain at injection sites, particularly in infants and children. Dissolve 500,000 polymyxin B units in 2 mL sterile water for injection or sodium chloride injection or procaine hydrochloride injection 1%.

### **Adults and children**

25,000 to 30,000 units/kg/day. This should be reduced in the presence of renal impairment. The dosage may be divided and given at either 4 or 6 hour intervals.

### **Infants**

Infants with normal kidney function may receive up to 40,000 units/kg/day without adverse effects.

**Note:** Doses as high as 45,000 units/kg/day have been used in limited clinical studies in treating prematures and newborn infants for sepsis caused by *Ps aeruginosa*.

## **Intrathecal**

**A treatment of choice for *Ps aeruginosa* meningitis.** Dissolve 500,000 polymyxin B units in 10 mL sodium chloride injection USP for 50,000 units per mL dosage unit.

### **Adults and children over 2 years of age**

Dosage is 50,000 units once daily intrathecally for 3 to 4 days, then 50,000 units once every other day for at least 2 weeks after cultures of the cerebrospinal fluid are negative and sugar content has returned to normal.

### **Children under 2 years of age**

20,000 units once daily, intrathecally for 3 to 4 days or 25,000 units once every other day. Continue with a dose of 25,000 units once every other day for at least 2 weeks after cultures of the cerebrospinal fluid are negative and sugar content has returned to normal.

**IN THE INTEREST OF SAFETY, SOLUTIONS OF PARENTERAL USE SHOULD BE STORED UNDER REFRIGERATION, AND ANY UNUSED PORTIONS SHOULD BE DISCARDED AFTER 72 HOURS.**

## **TOPICAL:**

### **Ophthalmic**

Dissolve 500,000 polymyxin B units in 20 to 50 mL sterile water for injection or sodium chloride injection, USP for a 10,000 to 25,000 units per mL concentration.

For the treatment of *Ps aeruginosa* infections of the eye, a concentration of 0.1 percent

to 0.25 percent (10,000 units to 25,000 units per mL) is administered 1 to 3 drops every hour, increasing the intervals as response indicates.

Subconjunctival injection of up to 100,000 units/day may be used for the treatment of *Ps aeruginosa* infections of the cornea and conjunctiva.

*Note: Avoid total systemic and ophthalmic instillation over 25,000 units/kg/day.*

## **HOW SUPPLIED**

Polymyxin B for Injection, USP is a sterile, white lyophilized cake or powder and is supplied as follows:

### **500,000 Units per Vial**

10 Vials in a Carton

NDC 55150-234-10

### **Storage recommendations:**

**Before reconstitution:** Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

**Protect from light.** Retain in carton until time of use.

**After reconstitution:** Product must be stored under refrigeration, between 2° to 8°C (36° to 46°F) and any unused portion should be discarded after 72 hours.

The vial stoppers are not made with natural rubber latex.

Distributed by:

**Eugia US LLC**

279 Princeton-Hightstown Rd.

E. Windsor, NJ 08520

Manufactured by:

**Eugia Pharma Specialities Limited**

Hyderabad - 500032

India

Revised: June 2023

## **PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 500,000 Units per Vial - Container Label**


**Rx only**                    **NDC 55150-234-10**

**Polymyxin B**

**for Injection USP**

**500,000 Units per Vial**

**For Parenteral or  
Ophthalmic Administration  
Sterile**



Rx only NDC 55150-234-10

**Polymyxin B  
for Injection USP**

**500,000 Units per Vial**

For Parenteral or  
Ophthalmic Administration

Sterile  
**eugia**



**Usual Dosage:** See package insert. Dosages should not exceed 25,000 units per kg per day.

**WARNING:** This drug may be neuro- and nephrotoxic. Administer by the intramuscular, intravenous, and/or intrathecal routes in hospitalized patients only. **FOR HOSPITAL USE ONLY**, if prescribed for intramuscular, intravenous, and/or intrathecal administration.

**STORAGE: Before Reconstitution:** Store at 20° to 25°C (68° to 77°F) [see USP]. **After Reconstitution:** Refrigerate between 2° to 8°C (36° to 46°F). Any unused portion should be discarded after 72 hours. **Protect from light.**

**Discard after** \_\_\_\_\_ / \_\_\_\_\_ / \_\_\_\_\_

Mfd. in India for:  
**Eugia US LLC**  
E. Windsor, NJ 08520

P1432228


Code: TS/DRUGS/13/2010



**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 500,000 Units per Vial -  
Container-Carton (10 Vials)**

Rx only NDC 55150-234-10  
**Polymyxin B  
for Injection USP**  
**500,000 Units per Vial**  
For Parenteral or  
Ophthalmic Administration  
Sterile  
**eugia**

**10 Vials**



**POLYMYXIN B**

polymyxin b injection, powder, lyophilized, for solution



## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:55150-234
<b>Route of Administration</b>	INTRAMUSCULAR, INTRAVENOUS, INTRATHECAL, OPTHALMIC		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	500000 [USP'U]

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:55150-234-10	10 in 1 CARTON	04/04/2016	
1		1 in 1 VIAL; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA206589	04/04/2016	

**Labeler** - Eugia US LLC (968961354)

## Establishment

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Eugia Pharma Specialities Limited		650498244	ANALYSIS(55150-234) , MANUFACTURE(55150-234) , PACK(55150-234)

Revised: 8/2023

Eugia US LLC