

**VALACYCLOVIR HYDROCHLORIDE- valacyclovir hydrochloride tablet, film coated**  
**Wockhardt Limited**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use valacyclovir hydrochloride safely and effectively. See full prescribing information for valacyclovir hydrochloride.

**Valacyclovir hydrochloride tablets**

**Rx only**

**Initial U.S. Approval: 1995**

**RECENT MAJOR CHANGES**

Warnings and Precautions, Central Nervous System Effects (5.3) 3/2010

**INDICATIONS AND USAGE**

Valacyclovir hydrochloride is a nucleoside analogue DNA polymerase inhibitor indicated for:

**Adult Patients (1.1)**

- Cold Sores (Herpes Labialis)
- Genital Herpes
- Treatment in immunocompetent patients (initial or recurrent episode)
- Suppression in immunocompetent or HIV-infected patients
- Reduction of transmission
- Herpes Zoster

**Pediatric Patients (1.2)**

- Cold Sores (Herpes Labialis)
- Chickenpox

**Limitations of Use (1.3)**

- The efficacy and safety of valacyclovir hydrochloride tablets have not been established in immunocompromised patients other than for the suppression of genital herpes in HIV-infected patients.

**DOSAGE AND ADMINISTRATION**

<b>Adult Dosage (2.1)</b>	
Cold Sores	2 grams every 12 hours for 1 day
Genital Herpes	
Initial episode	1 gram twice daily for 10 days
Recurrent episodes	500 mg twice daily for 3 days
Suppressive therapy	
Immunocompetent patients	1 gram once daily
Alternate dose in patients with $\leq 9$ recurrences/yr	500 mg once daily
HIV-infected patients	500 mg twice daily
Reduction of transmission	500 mg once daily
Herpes Zoster	1 gram 3 times daily for 7 days
<b>Pediatric Dosage (2.2)</b>	
Cold Sores ( $\geq 12$ years of age)	2 grams every 12 hours for 1 day
Chickenpox (2 to < 18 years of age)	20 mg/kg 3 times daily for 5 days; not to exceed 1 gram 3 times daily

Valacyclovir oral suspension (25 mg/mL or 50 mg/mL) can be prepared from valacyclovir hydrochloride tablets, 500 mg. (2.3)

**DOSAGE FORMS AND STRENGTHS**

Tablets: 500 mg (unscored), 1 gram (partially bisect) (3)

**CONTRAINDICATIONS**

Hypersensitivity to valacyclovir (e.g., anaphylaxis), acyclovir, or any component of the formulation. (4)

**WARNINGS AND PRECAUTIONS**

- Thrombotic thrombocytopenic purpura/hemolytic uremic syndrome (TTP/HUS): Has occurred in patients with advanced HIV disease and in allogeneic bone marrow transplant and renal transplant patients receiving 8 grams per day of valacyclovir hydrochloride in clinical trials. Discontinue treatment if clinical symptoms and laboratory findings consistent with TTP/HUS occur. (5.1)
- Acute renal failure: May occur in elderly patients (with or without reduced renal function), patients with underlying renal disease who receive higher than recommended doses of valacyclovir hydrochloride for their level of renal function, patients who receive concomitant nephrotoxic drugs, or inadequately hydrated patients. Use with caution in elderly patients and reduce dosage in patients with renal impairment. (2.4, 5.2)
- Central nervous system adverse reactions (e.g., agitation, hallucinations, confusion, and encephalopathy): May occur in both adult and pediatric patients (with or without reduced renal function) and in patients with underlying renal disease who receive higher than recommended doses of valacyclovir hydrochloride for their level of renal function. Elderly patients are more likely to have central nervous system adverse reactions. Use with caution in elderly patients and reduce dosage in patients with renal impairment. (2.4, 5.3)

**ADVERSE REACTIONS**

- The most common adverse reactions reported in at least one indication by >10% of adult patients treated with valacyclovir hydrochloride and more commonly than in patients treated with placebo are headache, nausea, and abdominal pain. (6.1)
- The only adverse reaction occurring in >10% of pediatric patients <18 years of age was headache. (6.2)

To report SUSPECTED ADVERSE REACTIONS, contact Wockhardt USA LLC., at 1-800-346-6854 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 3/2013

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1. INDICATIONS AND USAGE**

- 1.1 Adult Patients
- 1.2 Pediatric Patients
- 1.3 Limitations of Use

**2. DOSAGE AND ADMINISTRATION**

- 2.1 Adult Dosing Recommendations
- 2.2 Pediatric Dosing Recommendations
- 2.3 Extemporaneous Preparation of Oral Suspension
- 2.4 Patients With Renal Impairment

**3. DOSAGE FORMS AND STRENGTHS**

**4. CONTRAINDICATIONS**

**5. WARNINGS AND PRECAUTIONS**

- 5.1 Thrombotic Thrombocytopenic Purpura/Hemolytic Uremic Syndrome (TTP/HUS)
- 5.2 Acute Renal Failure
- 5.3 Central Nervous System Effects

**6. ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience in Adult Patients
- 6.2 Clinical Trials Experience in Pediatric Patients
- 6.3 Postmarketing Experience

**7. DRUG INTERACTIONS**

**8. USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

**10. OVERDOSAGE**

**11. DESCRIPTION**

**12. CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics
- 12.4 Microbiology
- 13. NONCLINICAL TOXICOLOGY**
- 13.1 Carcinogenesis, Mutagenesis, Impairment Of Fertility
- 14. CLINICAL STUDIES**
- 14.1 Cold Sores (Herpes Labialis)
- 14.2 Genital Herpes Infections
- 14.3 Herpes Zoster
- 14.4 Chickenpox
- 16. HOW SUPPLIED/STORAGE AND HANDLING**
- 17. PATIENT COUNSELING INFORMATION**
- 17.1 Importance of Adequate Hydration
- 17.2 Cold Sores (Herpes Labialis)
- 17.3 Genital Herpes
- 17.4 Herpes Zoster
- 17.5 Chickenpox

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### 1. INDICATIONS AND USAGE

#### 1.1 Adult Patients

**Cold Sores (Herpes Labialis):** Valacyclovir hydrochloride tablets are indicated for treatment of cold sores (herpes labialis). The efficacy of valacyclovir hydrochloride tablets initiated after the development of clinical signs of a cold sore (e.g., papule, vesicle, or ulcer) has not been established.

**Genital Herpes: Initial Episode:** Valacyclovir hydrochloride tablets are indicated for treatment of the initial episode of genital herpes in immunocompetent adults. The efficacy of treatment with valacyclovir hydrochloride tablets when initiated more than 72 hours after the onset of signs and symptoms has not been established.

**Recurrent Episodes:** Valacyclovir hydrochloride tablets are indicated for treatment of recurrent episodes of genital herpes in immunocompetent adults. The efficacy of treatment with valacyclovir hydrochloride tablets when initiated more than 24 hours after the onset of signs and symptoms has not been established.

**Suppressive Therapy:** Valacyclovir hydrochloride tablets are indicated for chronic suppressive therapy of recurrent episodes of genital herpes in immunocompetent and in HIV-infected adults. The efficacy and safety of valacyclovir hydrochloride tablets for the suppression of genital herpes beyond 1 year in immunocompetent patients and beyond 6 months in HIV-infected patients have not been established.

**Reduction of Transmission:** Valacyclovir hydrochloride tablets are indicated for the reduction of transmission of genital herpes in immunocompetent adults. The efficacy of valacyclovir hydrochloride tablets for the reduction of transmission of genital herpes beyond 8 months in discordant couples has not been established. The efficacy of valacyclovir hydrochloride tablets for the reduction of transmission of genital herpes in individuals with multiple partners and non-heterosexual couples has not been established. Safer sex practices should be used with suppressive therapy (see current Centers for Disease Control and Prevention [CDC] *Sexually Transmitted Diseases Treatment Guidelines*).

**Herpes Zoster:** Valacyclovir hydrochloride tablets are indicated for the treatment of herpes zoster (shingles) in immunocompetent adults. The efficacy of valacyclovir hydrochloride tablets when initiated more than 72 hours after the onset of rash and the efficacy and safety of valacyclovir hydrochloride tablets for treatment of disseminated herpes zoster have not been established.

#### 1.2 Pediatric Patients

**Cold Sores (Herpes Labialis):** Valacyclovir hydrochloride tablets are indicated for the treatment of cold sores (herpes labialis) in pediatric patients  $\geq 12$  years of age. The efficacy of valacyclovir hydrochloride tablets initiated after the development of clinical signs of a cold sore (e.g., papule, vesicle, or ulcer) has not been established.

**Chickenpox:** Valacyclovir hydrochloride tablets are indicated for the treatment of chickenpox in immunocompetent pediatric patients 2 to < 18 years of age. Based on efficacy data from clinical studies with oral acyclovir, treatment with valacyclovir hydrochloride tablets should be initiated within 24 hours after the onset of rash [see *Clinical Studies (14.4)*].

#### 1.3 Limitations of Use

The efficacy and safety of valacyclovir hydrochloride tablets have not been established in:

- Immunocompromised patients other than for the suppression of genital herpes in HIV-infected patients with a CD4+ cell count  $\geq 100$  cells/mm<sup>3</sup>.
- Patients <12 years of age with cold sores (herpes labialis).
- Patients < 2 years of age or  $\geq 18$  years of age with chickenpox.
- Patients <18 years of age with genital herpes.
- Patients <18 years of age with herpes zoster.
- Neonates and infants as suppressive therapy following neonatal herpes simplex virus (HSV) infection.

### 2. DOSAGE AND ADMINISTRATION

- Valacyclovir hydrochloride tablets may be given without regard to meals.
- Valacyclovir oral suspension (25 mg/mL or 50 mg/mL) may be prepared extemporaneously from valacyclovir hydrochloride tablets, 500 mg for use in pediatric patients for whom a solid dosage form is not appropriate [see *Dosage and Administration (2.3)*].

#### 2.1 Adult Dosing Recommendations

**Cold Sores (Herpes Labialis):** The recommended dosage of valacyclovir hydrochloride tablets for treatment of cold sores is 2 grams twice daily for 1 day taken 12 hours apart. Therapy should be initiated at the earliest symptom of a cold sore (e.g., tingling, itching, or burning).

**Genital Herpes: Initial Episode:** The recommended dosage of valacyclovir hydrochloride tablets for treatment of initial genital herpes is 1 gram twice daily for 10 days. Therapy was most effective when administered within 48 hours of the onset of signs and symptoms.

**Recurrent Episodes:** The recommended dosage of valacyclovir hydrochloride tablets for treatment of recurrent genital herpes is 500 mg twice daily for 3 days. Initiate treatment at the first sign or symptom of an episode.

**Suppressive Therapy:** The recommended dosage of valacyclovir hydrochloride tablets for chronic suppressive therapy of recurrent genital herpes is 1 gram once daily in patients with normal immune function. In patients with a history of 9 or fewer recurrences per year, an alternative dose is 500 mg once daily.

In HIV-infected patients with a CD4+ cell count  $\geq 100$  cells/mm<sup>3</sup>, the recommended dosage of valacyclovir hydrochloride tablets for chronic suppressive therapy of recurrent genital herpes is 500

mg twice daily.

**Reduction of Transmission:** The recommended dosage of valacyclovir hydrochloride tablets for reduction of transmission of genital herpes in patients with a history of 9 or fewer recurrences per year is 500 mg once daily for the source partner.

**Herpes Zoster:** The recommended dosage of valacyclovir hydrochloride tablets for treatment of herpes zoster is 1 gram 3 times daily for 7 days. Therapy should be initiated at the earliest sign or symptom of herpes zoster and is most effective when started within 48 hours of the onset of rash.

## 2.2 Pediatric Dosing Recommendations

**Cold Sores (Herpes Labialis):** The recommended dosage of valacyclovir hydrochloride tablets for the treatment of cold sores in pediatric patients  $\geq 12$  years of age is 2 grams twice daily for 1 day taken 12 hours apart. Therapy should be initiated at the earliest symptom of a cold sore (e.g., tingling, itching, or burning).

**Chickenpox:** The recommended dosage of valacyclovir hydrochloride tablets for treatment of chickenpox in immunocompetent pediatric patients 2 to  $< 18$  years of age is 20 mg/kg administered 3 times daily for 5 days. The total dose should not exceed 1 gram 3 times daily. Therapy should be initiated at the earliest sign or symptom [see *Use in Specific Populations (8.4), Clinical Pharmacology (12.3), Clinical Studies (14.4)*].

## 2.3 Extemporaneous Preparation of Oral Suspension

**Ingredients and Preparation per USP-NF:** Valacyclovir hydrochloride tablets, 500 mg, cherry flavor, and Suspension Structured Vehicle USP-NF (SSV). Valacyclovir oral suspension (25 mg/mL or 50 mg/mL) should be prepared in lots of 100 mL.

**Prepare Suspension at Time of Dispensing as Follows:**

- Prepare SSV according to the USP-NF.
- Using a pestle and mortar, grind the required number of valacyclovir hydrochloride tablets, 500 mg until a fine powder is produced (5 valacyclovir hydrochloride tablets for 25 mg/mL suspension; 10 valacyclovir hydrochloride tablets for 50 mg/mL suspension).
- Gradually add approximately 5 mL aliquots of SSV to the mortar and triturate the powder until a paste has been produced. Ensure that the powder has been adequately wetted.
- Continue to add approximately 5 mL aliquots of SSV to the mortar, mixing thoroughly between additions, until a concentrated suspension is produced, to a minimum total quantity of 20 mL SSV and a maximum total quantity of 40 mL SSV for both the 25 mg/mL and 50 mg/mL suspensions.
- Transfer the mixture to a suitable 100 mL measuring flask.
- Transfer the cherry flavor\* to the mortar and dissolve in approximately 5 mL of SSV. Once dissolved, add to the measuring flask.
- Rinse the mortar at least 3 times with approximately 5 mL aliquots of SSV, transferring the rinsing to the measuring flask between additions.
- Make the suspension to volume (100 mL) with SSV and shake thoroughly to mix.
- Transfer the suspension to an amber glass medicine bottle with a child-resistant closure.
- The prepared suspension should be labeled with the following information "Shake well before using. Store suspension between 2° to 8°C (36° to 46°F) in a refrigerator. Discard after 28 days."

\*The amount of cherry flavor added is as instructed by the suppliers of the cherry flavor.

## 2.4 Patients With Renal Impairment

Dosage recommendations for adult patients with reduced renal function are provided in Table 1 [see *Use in Specific Populations (8.5, 8.6), Clinical Pharmacology (12.3)*]. Data are not available for the use of valacyclovir hydrochloride tablets in pediatric patients with a creatinine clearance  $< 50$  mL/min/1.73 m<sup>2</sup>.

**Table 1. Valacyclovir Hydrochloride Tablets Dosage Recommendations for Adults With Renal Impairment**

Indications	Normal Dosage Regimen (Creatinine Clearance $\geq 50$ mL/min)	Creatinine Clearance (mL/min)		
		30-49	10-29	$< 10$
<b>Cold sores (Herpes labialis)</b>	Two 2 gram doses taken 12 hours apart	Two 1 gram doses taken 12 hours apart	Two 500 mg doses taken 12 hours apart	500 mg single dose
<b>Do not exceed 1 day of treatment.</b>				
<b>Genital herpes:</b>	1 gram every 12 hours	no reduction	1 gram every 24 hours	500 mg every 24 hours
<b>Initial episode</b>				
<b>Genital herpes:</b>	500 mg every 12 hours	no reduction	500 mg every 24 hours	500 mg every 24 hours
<b>Recurrent episode</b>				
<b>Genital herpes:</b>	1 gram every 24 hours	no reduction	500 mg every 24 hours	500 mg every 24 hours
<b>Suppressive therapy</b>				
Immunocompetent patients	500 mg every 24 hours	no reduction	500 mg every 48 hours	500 mg every 48 hours
Alternate dose for immunocompetent patients with $\leq 9$ recurrences/year	500 mg every 24 hours	no reduction	500 mg every 24 hours	500 mg every 24 hours
HIV-infected patients	500 mg every 12 hours	no reduction	500 mg every 24 hours	500 mg every 24 hours
<b>Herpes zoster</b>	1 gram every 8 hours	1 gram every 12 hours	1 gram every 24 hours	500 mg every 24 hours

**Hemodialysis:** Patients requiring hemodialysis should receive the recommended dose of valacyclovir hydrochloride tablets after hemodialysis. During hemodialysis, the half-life of acyclovir after administration of valacyclovir hydrochloride tablets are approximately 4 hours. About one third of acyclovir in the body is removed by dialysis during a 4-hour hemodialysis session.

**Peritoneal Dialysis:** There is no information specific to administration of valacyclovir hydrochloride tablets in patients receiving peritoneal dialysis. The effect of chronic ambulatory peritoneal dialysis (CAPD) and continuous arteriovenous hemofiltration/dialysis (CAVHD) on acyclovir pharmacokinetics has been studied. The removal of acyclovir after CAPD and CAVHD is less pronounced than with hemodialysis, and the pharmacokinetic parameters closely resemble those observed in patients with end-stage renal disease (ESRD) not receiving hemodialysis. Therefore, supplemental doses of valacyclovir hydrochloride tablets should not be required following CAPD or CAVHD.

## 3. DOSAGE FORMS AND STRENGTHS

Tablets:

- 500 mg: blue, film-coated, capsule-shaped tablets debossed with "W152" on one side and plain and other side.
- 1 gram: blue, film-coated, capsule-shaped tablets, debossed with "W153" on one side and partial bisect on both sides.

## 4. CONTRAINDICATIONS

Valacyclovir hydrochloride tablets are contraindicated in patients who have had a demonstrated clinically significant hypersensitivity reaction (e.g., anaphylaxis) to valacyclovir, acyclovir, or any component of the formulation [see *Adverse Reactions (6.3)*].

## 5. WARNINGS AND PRECAUTIONS



Laboratory Abnormality	(n = 967)	(n = 195)	(n = 1,194)	(n = 1,159)	(n = 439)	(n = 269)	(n = 266)	(n = 134)
Hemoglobin (<0.8 x LLN)	0.8%	0%	0.3%	0.2%	0%	0%	0.8%	0.8%
White blood cells (<0.75 x LLN)	1.3%	0.6%	0.7%	0.6%	0.2%	0.7%	0.8%	1.5%
Platelet count (<100,000/mm <sup>3</sup> )	1.0%	1.2%	0.3%	0.1%	0.7%	0.4%	1.1%	1.5%
AST (SGOT) (>2 x ULN)	1.0%	0%	1.0%	<sup>a</sup>	0.5%	4.1%	3.8%	3.0%
Serum creatinine (>1.5 x ULN)	0.2%	0%	0.7%	0%	0%	0%	0%	0%

<sup>a</sup>Data were not collected prospectively.

LLN = Lower limit of normal.

ULN = Upper limit of normal.

## 6.2 Clinical Trials Experience in Pediatric Patients

The safety profile of valacyclovir hydrochloride has been studied in 177 pediatric patients 1 month to <18 years of age. Sixty-five of these pediatric patients, 12 to <18 years of age, received oral caplets for 1 to 2 days for treatment of cold sores. The remaining 112 pediatric patients, 1 month to <12 years of age, participated in 3 pharmacokinetic and safety studies and received valacyclovir oral suspension. Fifty-one of these 112 pediatric patients received oral suspension for 3 to 6 days. The frequency, intensity, and nature of clinical adverse reactions and laboratory abnormalities were similar to those seen in adults.

**Pediatric Patients 12 to <18 Years of Age (Cold Sores):** In clinical studies for the treatment of cold sores, the adverse reactions reported by adolescent patients receiving valacyclovir hydrochloride 2 grams twice daily for 1 day, or valacyclovir hydrochloride 2 grams twice daily for 1 day followed by 1 gram twice daily for 1 day (n = 65, across both dosing groups), or placebo (n = 30), respectively, included headache (17%, 3%) and nausea (8%, 0%).

**Pediatric Patients 1 Month to <12 Years of Age:** Adverse events reported in more than 1 subject across the 3 pharmacokinetic and safety studies in children 1 month to <12 years of age were diarrhea (5%), pyrexia (4%), dehydration (2%), and rhinorrhea (2%). No clinically meaningful changes in laboratory values were observed.

## 6.3 Postmarketing Experience

In addition to adverse events reported from clinical trials, the following events have been identified during postmarketing use of valacyclovir hydrochloride. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to valacyclovir hydrochloride.

**General:** Facial edema, hypertension, tachycardia.

**Allergic:** Acute hypersensitivity reactions including anaphylaxis, angioedema, dyspnea, pruritus, rash, and urticaria [see *Contraindications (4)*].

**CNS Symptoms:** Aggressive behavior; agitation; ataxia; coma; confusion; decreased consciousness; dysarthria; encephalopathy; mania; and psychosis, including auditory and visual hallucinations, seizures, tremors [see *Warnings and Precautions (5.3)*, *Use in Specific Populations (8.5)*, *(8.6)*].

**Eye:** Visual abnormalities.

**Gastrointestinal:** Diarrhea.

**Hepatobiliary Tract and Pancreas:** Liver enzyme abnormalities, hepatitis.

**Renal:** Renal failure, renal pain (may be associated with renal failure) [see *Warnings and Precautions (5.2)*, *Use in Specific Populations (8.5)*, *(8.6)*].

**Hematologic:** Thrombocytopenia, aplastic anemia, leukocytoclastic vasculitis, TTP/HUS [see *Warnings and Precautions (5.1)*].

**Skin:** Erythema multiforme, rashes including photosensitivity, alopecia.

## 7. DRUG INTERACTIONS

No clinically significant drug-drug or drug-food interactions with valacyclovir hydrochloride are known [see *Clinical Pharmacology (12.3)*].

## 8. USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category B. There are no adequate and well-controlled studies of valacyclovir hydrochloride or acyclovir in pregnant women. Based on prospective pregnancy registry data on 749 pregnancies, the overall rate of birth defects in infants exposed to acyclovir in-utero appears similar to the rate for infants in the general population. Valacyclovir hydrochloride should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

A prospective epidemiologic registry of acyclovir use during pregnancy was established in 1984 and completed in April 1999. There were 749 pregnancies followed in women exposed to systemic acyclovir during the first trimester of pregnancy resulting in 756 outcomes. The occurrence rate of birth defects approximates that found in the general population. However, the small size of the registry is insufficient to evaluate the risk for less common defects or to permit reliable or definitive conclusions regarding the safety of acyclovir in pregnant women and their developing fetuses.

Animal reproduction studies performed at oral doses that provided up to 10 and 7 times the human plasma levels during the period of major organogenesis in rats and rabbits, respectively, revealed no evidence of teratogenicity.

### 8.3 Nursing Mothers

Following oral administration of a 500 mg dose of valacyclovir hydrochloride to 5 nursing mothers, peak acyclovir concentrations (C<sub>max</sub>) in breast milk ranged from 0.5 to 2.3 times (median 1.4) the corresponding maternal acyclovir serum concentrations. The acyclovir breast milk AUC ranged from 1.4 to 2.6 times (median 2.2) maternal serum AUC. A 500 mg maternal dosage of valacyclovir hydrochloride twice daily would provide a nursing infant with an oral acyclovir dosage of approximately 0.6 mg/kg/day. This would result in less than 2% of the exposure obtained after administration of a standard neonatal dose of 30 mg/kg/day of intravenous acyclovir to the nursing infant. Unchanged valacyclovir was not detected in maternal serum, breast milk, or infant urine. Caution should be exercised when valacyclovir hydrochloride is administered to a nursing woman.

### 8.4 Pediatric Use

Valacyclovir hydrochloride tablets are indicated for treatment of cold sores in pediatric patients ≥12 years of age [see *Indications and Usage (1.2)*, *Dosage and Administration (2.2)*].

The use of valacyclovir hydrochloride tablets for treatment of cold sores is based on 2 double-blind, placebo-controlled clinical trials in healthy adults and adolescents (≥12 years of age) with a history of recurrent cold sores [see *Clinical Studies (14.1)*].

The use of valacyclovir hydrochloride tablets for treatment of chickenpox in pediatric patients 2 to <18 years of age is based on single-dose pharmacokinetic and multiple-dose safety data from an open-label trial with valacyclovir and supported by efficacy and safety data from 3 randomized, double-blind, placebo-controlled trials evaluating oral acyclovir in pediatric patients with chickenpox [see *Dosage and Administration (2.2)*, *Adverse Reactions (6.2)*, *Clinical Pharmacology (12.3)*, *Clinical Studies (14.4)*].

The efficacy and safety of valacyclovir have not been established in pediatric patients:

- <12 years of age with cold sores
- <18 years of age with genital herpes
- <18 years of age with herpes zoster
- < 2 years of age with chickenpox
- for suppressive therapy following neonatal HSV infection.

The pharmacokinetic profile and safety of valacyclovir oral suspension in children < 12 years of age were studied in 3 open-label studies. No efficacy evaluations were conducted in any of the 3 studies.

Study 1 was a single-dose pharmacokinetic, multiple-dose safety study in 27 pediatric patients 1 to < 12 years of age with clinically suspected varicella-zoster virus (VZV) infection [see *Dosage and Administration* (2.2), *Adverse Reactions* (6.2), *Clinical Pharmacology* (12.3), *Clinical Studies* (14.4)].

Study 2 was a single-dose pharmacokinetic and safety study in pediatric patients 1 month to < 6 years of age who had an active herpes virus infection or who were at risk for herpes virus infection. Fifty-seven subjects were enrolled and received a single dose of 25 mg/kg valacyclovir oral suspension. In infants and children 3 months to < 6 years of age, this dose provided comparable systemic acyclovir exposures to that from a 1 gram dose of valacyclovir in adults (historical data). In infants 1 month to < 3 months of age, mean acyclovir exposures resulting from a 25 mg/kg dose were higher ( $C_{max}$ : ↑30%, AUC: ↑60%) than acyclovir exposures following a 1 gram dose of valacyclovir in adults. Acyclovir is not approved for suppressive therapy in infants and children following neonatal HSV infections; therefore valacyclovir is not recommended for this indication because efficacy cannot be extrapolated from acyclovir.

Study 3 was a single-dose pharmacokinetic, multiple-dose safety study in 28 pediatric patients 1 to < 12 years of age with clinically suspected HSV infection. None of the children enrolled in this study had genital herpes. Each subject was dosed with valacyclovir oral suspension, 10 mg/kg twice daily for 3 to 5 days. Acyclovir systemic exposures in pediatric patients following valacyclovir oral suspension were compared with historical acyclovir systemic exposures in immunocompetent adults receiving the solid oral dosage form of valacyclovir or acyclovir for the treatment of recurrent genital herpes. The mean projected daily acyclovir systemic exposures in pediatric patients across all age-groups (1 to < 12 years of age) were lower ( $C_{max}$ : ↓20%, AUC: ↓33%) compared with the acyclovir systemic exposures in adults receiving valacyclovir 500 mg twice daily, but were higher (daily AUC: ↑16%) than systemic exposures in adults receiving acyclovir 200 mg 5 times daily. Insufficient data are available to support valacyclovir for the treatment of recurrent genital herpes in this age-group because clinical information on recurrent genital herpes in young children is limited; therefore, extrapolating efficacy data from adults to this population is not possible. Moreover, valacyclovir has not been studied in children 1 to < 12 years of age with recurrent genital herpes.

### 8.5 Geriatric Use

Of the total number of subjects in clinical studies of valacyclovir hydrochloride tablets, 906 were 65 and over, and 352 were 75 and over. In a clinical study of herpes zoster, the duration of pain after healing (post-herpetic neuralgia) was longer in patients 65 and older compared with younger adults. Elderly patients are more likely to have reduced renal function and require dose reduction. Elderly patients are also more likely to have renal or CNS adverse events [see *Dosage and Administration* (2.4), *Warnings and Precautions* (5.2, 5.3), *Clinical Pharmacology* (12.3)].

### 8.6 Renal Impairment

Dosage reduction is recommended when administering valacyclovir hydrochloride to patients with renal impairment [see *Dosage and Administration* (2.4), *Warnings and Precautions* (5.2, 5.3)].

## 10. OVERDOSAGE


Caution should be exercised to prevent inadvertent overdose [see *Use in Specific Populations* (8.5), (8.6)]. Precipitation of acyclovir in renal tubules may occur when the solubility (2.5 mg/mL) is exceeded in the intratubular fluid. In the event of acute renal failure and anuria, the patient may benefit from hemodialysis until renal function is restored [see *Dosage and Administration* (2.4)].

## 11. DESCRIPTION

Valacyclovir hydrochloride is the hydrochloride salt of *L*-valyl ester of the antiviral drug acyclovir.

Valacyclovir hydrochloride tablets are for oral administration. Each tablet contains valacyclovir hydrochloride equivalent to 500 mg or 1 gram valacyclovir and the inactive ingredients croscarmellose sodium, FD&C Blue No. 2 aluminium lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc and titanium dioxide.

The chemical name of valacyclovir hydrochloride is *L*-valine, 2-[(2-amino-1,6-dihydro-6-oxo-9H-purin-9-yl)methoxy]ethyl ester, monohydrochloride. It has the following structural formula:



Valacyclovir hydrochloride is a white to off-white powder with the molecular formula  $C_{13}H_{20}N_6O_4 \cdot HCl$  and a molecular weight of 360.80. The maximum solubility in water at 25°C is 174 mg/mL. The  $pK_s$  for valacyclovir hydrochloride are 1.90, 7.47, and 9.43.

## 12. CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Valacyclovir is an antiviral drug [see *Clinical Pharmacology* (12.4)].

### 12.3 Pharmacokinetics

The pharmacokinetics of valacyclovir and acyclovir after oral administration of valacyclovir hydrochloride have been investigated in 14 volunteer studies involving 283 adults and in 3 studies involving 112 pediatric subjects from 1 month to < 12 years of age.

**Pharmacokinetics in Adults: Absorption and Bioavailability:** After oral administration, valacyclovir hydrochloride is rapidly absorbed from the gastrointestinal tract and nearly completely converted to acyclovir and *L*-valine by first-pass intestinal and/or hepatic metabolism.

The absolute bioavailability of acyclovir after administration of valacyclovir hydrochloride is 54.5% ± 9.1% as determined following a 1 gram oral dose of valacyclovir hydrochloride and a 350 mg intravenous acyclovir dose to 12 healthy volunteers. Acyclovir bioavailability from the administration of valacyclovir hydrochloride is not altered by administration with food (30 minutes after an 873 Kcal breakfast, which included 51 grams of fat).

Acyclovir pharmacokinetic parameter estimates following administration of valacyclovir hydrochloride to healthy adult volunteers are presented in Table 3. There was a less than dose-proportional increase in acyclovir maximum concentration ( $C_{max}$ ) and area under the acyclovir concentration-time curve (AUC) after single-dose and multiple-dose administration (4 times daily) of valacyclovir hydrochloride from doses between 250 mg to 1 gram.

There is no accumulation of acyclovir after the administration of valacyclovir at the recommended dosage regimens in adults with normal renal function.

**Table 3. Mean (±SD) Plasma Acyclovir Pharmacokinetic Parameters Following Administration of Valacyclovir Hydrochloride to Healthy Adult Volunteers**

Dose	Single-Dose Administration (N = 8)		Multiple-Dose Administration* (N = 24, 8 per treatment arm)	
	$C_{max}$ (±SD) (mcg/mL)	AUC (±SD) (hr·mcg/mL)	$C_{max}$ (±SD) (mcg/mL)	AUC (±SD) (hr·mcg/mL)
100 mg	0.83 (±0.14)	2.28 (±0.40)	ND	ND
250 mg	2.15 (±0.50)	5.76 (±0.60)	2.11 (±0.33)	5.66 (±1.09)
500 mg	3.28 (±0.83)	11.59 (±1.79)	3.69 (±0.87)	9.88 (±2.01)
750 mg	4.17 (±1.14)	14.11 (±3.54)	ND	ND
1,000 mg	5.65 (±2.37)	19.52 (±6.04)	4.96 (±0.64)	15.70 (±2.27)

\*Administered 4 times daily for 11 days. ND = not done.

**Distribution:** The binding of valacyclovir to human plasma proteins ranges from 13.5% to 17.9%. The binding of acyclovir to human plasma proteins ranges from 9% to 33%.

**Metabolism:** Valacyclovir is converted to acyclovir and L-valine by first-pass intestinal and/or hepatic metabolism. Acyclovir is converted to a small extent to inactive metabolites by aldehyde oxidase and by alcohol and aldehyde dehydrogenase. Neither valacyclovir nor acyclovir is metabolized by cytochrome P450 enzymes. Plasma concentrations of unconverted valacyclovir are low and transient, generally becoming non-quantifiable by 3 hours after administration. Peak plasma valacyclovir concentrations are generally less than 0.5 mcg/mL at all doses. After single-dose administration of 1 gram of valacyclovir hydrochloride, average plasma valacyclovir concentrations observed were 0.5, 0.4, and 0.8 mcg/mL in patients with hepatic dysfunction, renal insufficiency, and in healthy volunteers who received concomitant cimetidine and probenecid, respectively.

**Elimination:** The pharmacokinetic disposition of acyclovir delivered by valacyclovir is consistent with previous experience from intravenous and oral acyclovir. Following the oral administration of a single 1 gram dose of radiolabeled valacyclovir to 4 healthy subjects, 46% and 47% of administered radioactivity was recovered in urine and feces, respectively, over 96 hours. Acyclovir accounted for 89% of the radioactivity excreted in the urine. Renal clearance of acyclovir following the administration of a single 1 gram dose of valacyclovir hydrochloride to 12 healthy volunteers was approximately 255 ± 86 mL/min which represents 42% of total acyclovir apparent plasma clearance.

The plasma elimination half-life of acyclovir typically averaged 2.5 to 3.3 hours in all studies of valacyclovir hydrochloride in volunteers with normal renal function.

**Specific Populations: Renal Impairment:** Reduction in dosage is recommended in patients with renal impairment [see Dosage and Administration (2.4), Use in Specific Populations (8.5), (8.6)].

Following administration of valacyclovir hydrochloride to volunteers with ESRD, the average acyclovir half-life is approximately 14 hours. During hemodialysis, the acyclovir half-life is approximately 4 hours. Approximately one third of acyclovir in the body is removed by dialysis during a 4-hour hemodialysis session. Apparent plasma clearance of acyclovir in dialysis patients was 86.3 ± 21.3 mL/min/1.73 m<sup>2</sup> compared with 679.16 ± 162.76 mL/min/1.73 m<sup>2</sup> in healthy volunteers.

**Hepatic Impairment:** Administration of valacyclovir hydrochloride to patients with moderate (biopsy-proven cirrhosis) or severe (with and without ascites and biopsy-proven cirrhosis) liver disease indicated that the rate but not the extent of conversion of valacyclovir to acyclovir is reduced, and the acyclovir half-life is not affected. Dosage modification is not recommended for patients with cirrhosis.

**HIV Disease:** In 9 patients with HIV disease and CD4+ cell counts <150 cells/mm<sup>3</sup> who received valacyclovir hydrochloride at a dosage of 1 gram 4 times daily for 30 days, the pharmacokinetics of valacyclovir and acyclovir were not different from that observed in healthy volunteers.

**Geriatrics:** After single-dose administration of 1 gram of valacyclovir hydrochloride in healthy geriatric volunteers, the half-life of acyclovir was 3.11 ± 0.51 hours, compared with 2.91 ± 0.63 hours in healthy younger adult volunteers. The pharmacokinetics of acyclovir following single- and multiple-dose oral administration of valacyclovir hydrochloride in geriatric volunteers varied with renal function. Dose reduction may be required in geriatric patients, depending on the underlying renal status of the patient [see Dosage and Administration (2.4), Use in Specific Populations (8.5), (8.6)].

**Pediatrics:** Acyclovir pharmacokinetics have been evaluated in a total of 98 pediatric patients (1 month to <12 years of age) following administration of the first dose of an extemporaneous oral suspension of valacyclovir [see Adverse Reactions (6.2), Use in Specific Populations (8.4)]. Acyclovir pharmacokinetic parameter estimates following a 20 mg/kg dose are provided in Table 4.

**Table 4. Mean (± SD) Plasma Acyclovir Pharmacokinetic Parameter Estimates Following First-Dose Administration of 20 mg/kg Valacyclovir Oral Suspension to Pediatric Patients vs. 1 Gram Single Dose of Valacyclovir Hydrochloride Tablets to Adults**

Parameter	Pediatric Patients (20 mg/kg Oral Suspension)			Adults 1 gram Solid Dose of Valacyclovir Hydrochloride Tablets <sup>a</sup> (N = 15)
	1 to < 2 yr (N = 6)	2 to < 6 yr (N = 12)	6 to < 12 yr (N = 8)	
AUC (mcg·hr/mL)	14.4 (± 6.26)	10.1 (± 3.35)	13.1 (± 3.43)	17.2 (± 3.10)
$C_{max}$ (mcg/mL)	4.03 (± 1.37)	3.75 (± 1.14)	4.71 (± 1.20)	4.72 (± 1.37)

<sup>a</sup> Historical estimates using pediatric pharmacokinetic sampling schedule.

**Drug Interactions:** When valacyclovir hydrochloride is coadministered with antacids, cimetidine and/or probenecid, digoxin, or thiazide diuretics in patients with normal renal function, the effects are not considered to be of clinical significance (see below). Therefore, when valacyclovir hydrochloride is coadministered with these drugs in patients with normal renal function, no dosage adjustment is recommended.

**Antacids:** The pharmacokinetics of acyclovir after a single dose of valacyclovir hydrochloride (1 gram) were unchanged by coadministration of a single dose of antacids (Al<sup>3+</sup> or Mg<sup>++</sup>).

**Cimetidine:** Acyclovir  $C_{max}$  and AUC following a single dose of valacyclovir hydrochloride (1 gram) increased by 8% and 32%, respectively, after a single dose of cimetidine (800 mg).

**Cimetidine Plus Probenecid:** Acyclovir  $C_{max}$  and AUC following a single dose of valacyclovir hydrochloride (1 gram) increased by 30% and 78%, respectively, after a combination of cimetidine and probenecid, primarily due to a reduction in renal clearance of acyclovir.

**Digoxin:** The pharmacokinetics of digoxin were not affected by coadministration of valacyclovir hydrochloride 1 gram 3 times daily, and the pharmacokinetics of acyclovir after a single dose of valacyclovir hydrochloride (1 gram) was unchanged by coadministration of digoxin (2 doses of 0.75 mg).

**Probenecid:** Acyclovir  $C_{max}$  and AUC following a single dose of valacyclovir hydrochloride (1 gram) increased by 22% and 49%, respectively, after probenecid (1 gram).

**Thiazide Diuretics:** The pharmacokinetics of acyclovir after a single dose of valacyclovir hydrochloride (1 gram) were unchanged by coadministration of multiple doses of thiazide diuretics.

## 12.4 Microbiology

**Mechanism of Action:** Valacyclovir is a nucleoside analogue DNA polymerase inhibitor. Valacyclovir hydrochloride is rapidly converted to acyclovir which has demonstrated antiviral activity against HSV types 1 (HSV-1) and 2 (HSV-2) and VZV both in cell culture and *in vivo*.

The inhibitory activity of acyclovir is highly selective due to its affinity for the enzyme thymidine kinase (TK) encoded by HSV and VZV. This viral enzyme converts acyclovir into acyclovir monophosphate, a nucleotide analogue. The monophosphate is further converted into diphosphate by cellular guanylate kinase and into triphosphate by a number of cellular enzymes. In biochemical assays, acyclovir triphosphate inhibits replication of herpes viral DNA. This is accomplished in 3 ways: 1) competitive inhibition of viral DNA polymerase, 2) incorporation and termination of the growing viral DNA chain, and 3) inactivation of the viral DNA polymerase. The greater antiviral activity of acyclovir against HSV compared with VZV is due to its more efficient phosphorylation by the viral TK.

**Antiviral Activities:** The quantitative relationship between the cell culture susceptibility of herpesviruses to antivirals and the clinical response to therapy has not been established in humans, and virus sensitivity testing has not been standardized. Sensitivity testing results, expressed as the concentration of drug required to inhibit by 50% the growth of virus in cell culture ( $EC_{50}$ ), vary greatly depending upon a number of factors. Using plaque-reduction assays, the  $EC_{50}$  values against herpes simplex virus isolates range from 0.09 to 60  $\mu$ M (0.02 to 13.5 mcg/mL) for HSV-1 and from 0.04 to 44  $\mu$ M (0.01 to 9.9 mcg/mL) for HSV-2. The  $EC_{50}$  values for acyclovir against most laboratory strains and clinical isolates of VZV range from 0.53 to 48  $\mu$ M (0.12 to 10.8 mcg/mL). Acyclovir also demonstrates activity against the Oka vaccine strain of VZV with a mean  $EC_{50}$  of 6  $\mu$ M (1.35 mcg/mL).

**Resistance:** Resistance of HSV and VZV to acyclovir can result from qualitative and quantitative changes in the viral TK and/or DNA polymerase. Clinical isolates of VZV with reduced susceptibility to acyclovir have been recovered from patients with AIDS. In these cases, TK-deficient mutants of VZV have been recovered.

Resistance of HSV and VZV to acyclovir occurs by the same mechanisms. While most of the acyclovir-resistant mutants isolated thus far from immunocompromised patients have been found to be TK-deficient mutants, other mutants involving the viral TK gene (TK partial and TK altered) and DNA polymerase have also been isolated. TK-negative mutants may cause severe disease in immunocompromised patients. The possibility of viral resistance to valacyclovir (and therefore, to acyclovir) should be considered in patients who show poor clinical response during therapy.

## 13. NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment Of Fertility

The data presented below include references to the steady-state acyclovir AUC observed in humans treated with 1 gram valacyclovir hydrochloride tablets given orally 3 times a day to treat herpes zoster. Plasma drug concentrations in animal studies are expressed as multiples of human exposure to acyclovir [see *Clinical Pharmacology* (12.3)].

Valacyclovir was noncarcinogenic in lifetime carcinogenicity bioassays at single daily doses (gavage) of valacyclovir giving plasma acyclovir concentrations equivalent to human levels in the mouse bioassay and 1.4 to 2.3 times human levels in the rat bioassay. There was no significant difference in the incidence of tumors between treated and control animals, nor did valacyclovir shorten the latency of tumors.

Valacyclovir was tested in 5 genetic toxicity assays. An Ames assay was negative in the absence or presence of metabolic activation. Also negative were an *in vitro* cytogenetic study with human lymphocytes and a rat cytogenetic study.

In the mouse lymphoma assay, valacyclovir was not mutagenic in the absence of metabolic activation. In the presence of metabolic activation (76% to 88% conversion to acyclovir), valacyclovir was mutagenic.

Valacyclovir was mutagenic in a mouse micronucleus assay.

Valacyclovir did not impair fertility or reproduction in rats at 6 times human plasma levels.

## 14. CLINICAL STUDIES

### 14.1 Cold Sores (Herpes Labialis)

Two double-blind, placebo-controlled clinical trials were conducted in 1,856 healthy adults and adolescents ( $\geq 12$  years old) with a history of recurrent cold sores. Patients self-initiated therapy at the earliest symptoms and prior to any signs of a cold sore. The majority of patients initiated treatment within 2 hours of onset of symptoms. Patients were randomized to valacyclovir hydrochloride 2 grams twice daily on Day 1 followed by placebo on Day 2, valacyclovir hydrochloride 2 grams twice daily on Day 1 followed by 1 gram twice daily on Day 2, or placebo on Days 1 and 2.

The mean duration of cold sore episodes was about 1 day shorter in treated subjects as compared with placebo. The 2 day regimen did not offer additional benefit over the 1-day regimen.

No significant difference was observed between subjects receiving valacyclovir hydrochloride or placebo in the prevention of progression of cold sore lesions beyond the papular stage.

### 14.2 Genital Herpes Infections

**Initial Episode:** Six hundred and forty-three immunocompetent adults with first-episode genital herpes who presented within 72 hours of symptom onset were randomized in a double-blind trial to receive 10 days of valacyclovir hydrochloride 1 gram twice daily ( $n = 323$ ) or oral acyclovir 200 mg 5 times a day ( $n = 320$ ). For both treatment groups: the median time to lesion healing was 9 days, the median time to cessation of pain was 5 days, the median time to cessation of viral shedding was 3 days.

**Recurrent Episodes:** Three double-blind trials (2 of them placebo-controlled) in immunocompetent adults with recurrent genital herpes were conducted. Patients self-initiated therapy within 24 hours of the first sign or symptom of a recurrent genital herpes episode.

In 1 study, patients were randomized to receive 5 days of treatment with either valacyclovir hydrochloride 500 mg twice daily ( $n = 360$ ) or placebo ( $n = 259$ ). The median time to lesion healing was 4 days in the group receiving valacyclovir hydrochloride 500 mg versus 6 days in the placebo group, and the median time to cessation of viral shedding in patients with at least 1 positive culture (42% of the overall study population) was 2 days in the group receiving valacyclovir hydrochloride 500 mg versus 4 days in the placebo group. The median time to cessation of pain was 3 days in the group receiving valacyclovir hydrochloride 500 mg versus 4 days in the placebo group. Results supporting efficacy were replicated in a second trial.

In a third study, patients were randomized to receive valacyclovir hydrochloride 500 mg twice daily for 5 days ( $n = 398$ ) or valacyclovir hydrochloride 500 mg twice daily for 3 days (and matching placebo



twice daily for 2 additional days) (n = 402). The median time to lesion healing was about 4½ days in both treatment groups. The median time to cessation of pain was about 3 days in both treatment groups.

**Suppressive Therapy:** Two clinical studies were conducted, one in immunocompetent adults and one in HIV-infected adults.

A double-blind, 12-month, placebo-and active-controlled study enrolled immunocompetent adults with a history of 6 or more recurrences per year. Outcomes for the overall study population are shown in Table 5.

**Table 5. Recurrence Rates in Immunocompetent Adults at 6 and 12 Months**

Outcome	6 Months			12 Months		
	Valacyclovir Hydrochloride 1 gram once daily (n = 269)	Oral acyclovir 400 mg twice daily (n = 267)	Placebo (n=134)	Valacyclovir Hydrochloride 1 gram once daily (n = 269)	Oral acyclovir 400 mg twice daily (n = 267)	Placebo (n = 134)
Recurrence free	55%	54%	7%	34%	34%	4%
Recurrences	35%	36%	83%	46%	46%	85%
Unknown*	10%	10%	10%	19%	19%	10%

\* Includes lost to follow-up, discontinuations due to adverse events, and consent withdrawn.

Subjects with 9 or fewer recurrences per year showed comparable results with valacyclovir hydrochloride 500 mg once daily.

In a second study, 293 HIV-infected adults on stable antiretroviral therapy with a history of 4 or more recurrences of ano-genital herpes per year were randomized to receive either valacyclovir hydrochloride 500 mg twice daily (n = 194) or matching placebo (n = 99) for 6 months. The median duration of recurrent genital herpes in enrolled subjects was 8 years, and the median number of recurrences in the year prior to enrollment was 5. Overall, the median prestudy HIV-1 RNA was 2.6 log<sub>10</sub> copies/mL. Among patients who received valacyclovir hydrochloride, the prestudy median CD4+ cell count was 336 cells/mm<sup>3</sup>; 11% had <100 cells/mm<sup>3</sup>, 16% had 100 to 199 cells/mm<sup>3</sup>, 42% had 200 to 499 cells/mm<sup>3</sup>, and 31% had ≥500 cells/mm<sup>3</sup>. Outcomes for the overall study population are shown in Table 6.

**Table 6. Recurrence Rates in HIV-Infected Adults at 6 Months**

Outcome	Valacyclovir Hydrochloride Tablets 500 mg twice daily (n = 194)	Placebo (n = 99)
Recurrence free	65%	26%
Recurrences	17%	57%
Unknown*	18%	17%

\* Includes lost to follow-up, discontinuations due to adverse events, and consent withdrawn.

**Reduction of Transmission of Genital Herpes:** A double-blind, placebo-controlled study to assess transmission of genital herpes was conducted in 1,484 monogamous, heterosexual, immunocompetent adult couples. The couples were discordant for HSV-2 infection. The source partner had a history of 9 or fewer genital herpes episodes per year. Both partners were counseled on safer sex practices and were advised to use condoms throughout the study period. Source partners were randomized to treatment with either valacyclovir hydrochloride 500 mg once daily or placebo once daily for 8 months. The primary efficacy endpoint was symptomatic acquisition of HSV-2 in susceptible partners. Overall HSV-2 acquisition was defined as symptomatic HSV-2 acquisition and/or HSV-2 seroconversion in susceptible partners. The efficacy results are summarized in Table 7.

**Table 7. Percentage of Susceptible Partners Who Acquired HSV-2 Defined by the Primary and Selected Secondary Endpoints**

Endpoint	Valacyclovir Hydrochloride* (n = 743)	Placebo (n = 741)
Symptomatic HSV-2 acquisition	4 (0.5%)	16 (2.2%)
HSV-2 seroconversion	12 (1.6%)	24 (3.2%)
Overall HSV-2 acquisition	14 (1.9%)	27 (3.6%)

\* Results show reductions in risk of 75% (symptomatic HSV-2 acquisition), 50% (HSV-2 seroconversion), and 48% (overall HSV-2 acquisition) with valacyclovir hydrochloride versus placebo. Individual results may vary based on consistency of safer sex practices.

### 14.3 Herpes Zoster

Two randomized double-blind clinical trials in immunocompetent adults with localized herpes zoster were conducted. Valacyclovir hydrochloride was compared with placebo in patients less than 50 years of age, and with oral acyclovir in patients greater than 50 years of age. All patients were treated within 72 hours of appearance of zoster rash. In patients less than 50 years of age, the median time to cessation of new lesion formation was 2 days for those treated with valacyclovir hydrochloride compared with 3 days for those treated with placebo. In patients greater than 50 years of age, the median time to cessation of new lesions was 3 days in patients treated with either valacyclovir hydrochloride or oral acyclovir. In patients less than 50 years of age, no difference was found with respect to the duration of pain after healing (post-herpetic neuralgia) between the recipients of valacyclovir hydrochloride and placebo. In patients greater than 50 years of age, among the 83% who reported pain after healing (post-herpetic neuralgia), the median duration of pain after healing [95% confidence interval] in days was: 40 [31, 51], 43 [36, 55], and 59 [41, 77] for 7-day valacyclovir hydrochloride, 14-day valacyclovir hydrochloride, and 7-day oral acyclovir, respectively.

### 14.4 Chickenpox

The use of valacyclovir hydrochloride tablets for treatment of chickenpox in pediatric patients 2 to < 18 years of age is based on single-dose pharmacokinetic and multiple-dose safety data from an open-label trial with valacyclovir and supported by safety and extrapolated efficacy data from 3 randomized, double-blind, placebo-controlled trials evaluating oral acyclovir in pediatric patients.

The single-dose pharmacokinetic and multiple-dose safety study enrolled 27 pediatric patients 1 to < 12 years of age with clinically suspected VZV infection. Each subject was dosed with valacyclovir oral suspension, 20 mg/kg 3 times daily for 5 days. Acyclovir systemic exposures in pediatric patients following valacyclovir oral suspension were compared with historical acyclovir systemic exposures in immunocompetent adults receiving the solid oral dosage form of valacyclovir or acyclovir for the treatment of herpes zoster. The mean projected daily acyclovir exposures in pediatric patients across all age-groups (1 to < 12 years of age) were lower (C<sub>max</sub>: +13%, AUC: +30%) than the mean daily historical exposures in adults receiving valacyclovir 1 gram 3 times daily, but were higher (daily AUC: +50%) than the mean daily historical exposures in adults receiving acyclovir 800 mg 5 times daily. The projected daily exposures in pediatric patients were greater (daily AUC approximately 100% greater) than the exposures seen in immunocompetent pediatric patients receiving acyclovir 20 mg/kg 4 times daily for the treatment of chickenpox. Based on the pharmacokinetic and safety data from this study and the safety and extrapolated efficacy data from the acyclovir studies, oral valacyclovir 20 mg/kg 3 times a day for 5 days (not to exceed 1 gram 3 times daily) is recommended for the treatment of chickenpox in pediatric patients 2 to < 18 years of age. Because the efficacy and safety of acyclovir for the treatment of chickenpox in children < 2 years of age have not been established, efficacy data cannot be extrapolated to support valacyclovir treatment in children < 2 years of age with chickenpox. Valacyclovir is also not recommended for the treatment of herpes zoster in children because safety data up to 7 days' duration are not available [see Use in Specific Populations (8.4)].

## 16. HOW SUPPLIED/STORAGE AND HANDLING

Valacyclovir hydrochloride tablets are blue, film-coated, capsule-shaped tablets containing valacyclovir hydrochloride equivalent to 500 mg valacyclovir debossed with "W152" on one side and plain on other side.

Bottle of 30 (NDC 64679-152-01).

Bottle of 90 (NDC 64679-152-03).

Bottle of 500 (NDC 64679-152-04).

Unit dose pack of 100 (NDC 64679-152-05).

Valacyclovir hydrochloride tablets are blue, film-coated, capsule-shaped tablets containing valacyclovir hydrochloride equivalent to 1 g valacyclovir debossed with "W153" on one side and partial bisect on both sides.

Bottle of 21 (NDC 64679-153-01).

Bottle of 30 (NDC 64679-153-02).

Bottle of 90 (NDC 64679-153-03).

Bottle of 250 (NDC 64679-153-06).

Bottle of 500 (NDC 64679-153-04).

Unit dose pack of 100 (NDC 64679-153-05).

### Storage:

Store at 20° to 25°C (68° to 77°F). [See USP controlled room temperature] Dispense in a well-closed container as defined in the USP.

## 17. PATIENT COUNSELING INFORMATION

See FDA-Approved Patient Labeling.

### 17.1 Importance of Adequate Hydration

Patients should be advised to maintain adequate hydration.

### 17.2 Cold Sores (Herpes Labialis)

Patients should be advised to initiate treatment at the earliest symptom of a cold sore (e.g., tingling, itching, or burning). There are no data on the effectiveness of treatment initiated after the development of clinical signs of a cold sore (e.g., papule, vesicle, or ulcer). Patients should be instructed that treatment for cold sores should not exceed 1 day (2 doses) and that their doses should be taken about 12 hours apart. Patients should be informed that valacyclovir hydrochloride is not a cure for cold sores.

### 17.3 Genital Herpes

Patients should be informed that valacyclovir hydrochloride is not a cure for genital herpes. Because genital herpes is a sexually transmitted disease, patients should avoid contact with lesions or intercourse when lesions and/or symptoms are present to avoid infecting partners. Genital herpes is frequently transmitted in the absence of symptoms through asymptomatic viral shedding. Therefore, patients should be counseled to use safer sex practices in combination with suppressive therapy with valacyclovir hydrochloride. Sex partners of infected persons should be advised that they might be infected even if they have no symptoms. Type-specific serologic testing of asymptomatic partners of persons with genital herpes can determine whether risk for HSV-2 acquisition exists.

Valacyclovir hydrochloride has not been shown to reduce transmission of sexually transmitted infections other than HSV-2.

If medical management of a genital herpes recurrence is indicated, patients should be advised to initiate therapy at the first sign or symptom of an episode.

There are no data on the effectiveness of treatment initiated more than 72 hours after the onset of signs and symptoms of a first episode of genital herpes or more than 24 hours after the onset of signs and symptoms of a recurrent episode.

There are no data on the safety or effectiveness of chronic suppressive therapy of more than 1 year's duration in otherwise healthy patients. There are no data on the safety or effectiveness of chronic suppressive therapy of more than 6 months' duration in HIV-infected patients.

### 17.4 Herpes Zoster

There are no data on treatment initiated more than 72 hours after onset of the zoster rash. Patients should be advised to initiate treatment as soon as possible after a diagnosis of herpes zoster.

### 17.5 Chickenpox

Patients should be advised to initiate treatment at the earliest sign or symptom of chickenpox.

### Manufactured by:

Wockhardt Limited

Mumbai, India.

### Distributed by:

Wockhardt USA LLC.

20 Waterview Blvd.

Parsippany, NJ 07054

USA.

Rev.080213

PHARMACIST: Dispense the patient information leaflet with the drug product

## PATIENT INFORMATION

### Valacyclovir Hydrochloride Tablets

#### Rx only

Read the Patient Information that comes with valacyclovir hydrochloride tablets before you start using it and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment. Ask your healthcare provider or pharmacist if you have questions.

#### What are valacyclovir hydrochloride tablets?

Valacyclovir hydrochloride tablets are prescription antiviral medicine. Valacyclovir hydrochloride tablet lowers the ability of herpes viruses to multiply in your body.

Valacyclovir hydrochloride tablets are used in adults:

- to treat cold sores (also called fever blisters or herpes labialis)
- to treat shingles (also called herpes zoster)
- to treat or control genital herpes outbreaks in adults with normal immune systems

- to control genital herpes outbreaks in adults infected with the human immunodeficiency virus (HIV) with CD4+ cell count greater than 100 cells/mm<sup>3</sup>
- with safer sex practices to lower the chances of spreading genital herpes to others. Even with safer sex practices, it is still possible to spread genital herpes.

Valacyclovir hydrochloride tablets used daily with the following safer sex practices can lower the chances of passing genital herpes to your partner.

- **Do not have sexual contact with your partner when you have any symptom or outbreak of genital herpes.**
- **Use a condom** made of latex or polyurethane whenever you have sexual contact.

Valacyclovir hydrochloride tablets are used in children:

- to treat cold sores (for children ≥12 years of age)
- to treat chickenpox (for children 2 to < 18 years of age).

**Valacyclovir hydrochloride tablet does not cure herpes infections** (cold sores, chickenpox, shingles, or genital herpes).

The efficacy of valacyclovir hydrochloride tablets have not been studied in children who have not reached puberty.

#### **What are cold sores, chickenpox, shingles, and genital herpes?**

**Cold sores** are caused by a herpes virus that may be spread by kissing or other physical contact with the infected area of the skin. They are small, painful ulcers that you get in or around your mouth. It is not known if valacyclovir hydrochloride tablets can stop the spread of cold sores to others.

**Chickenpox** is caused by a herpes virus. It causes an itchy rash of multiple small, red bumps that look like pimples or insect bites usually appearing first on the abdomen or back and face. It can spread to almost everywhere else on the body and may be accompanied by flu-like symptoms.

**Shingles** is caused by the same herpes virus that causes chickenpox. It causes small, painful blisters that happen on your skin. Shingles occurs in people who have already had chickenpox. Shingles can be spread to people who have not had chickenpox or the chickenpox vaccine by contact with the infected areas of the skin. It is not known if valacyclovir hydrochloride tablets can stop the spread of shingles to others.

**Genital herpes** is a sexually transmitted disease. It causes small, painful blisters on your genital area. You can spread genital herpes to others, even when you have no symptoms. If you are sexually active, you can still pass herpes to your partner, even if you are taking valacyclovir hydrochloride tablets. Valacyclovir hydrochloride tablets, taken every day as prescribed and used with the following **safer sex practices**, can lower the chances of passing genital herpes to your partner.

- Do not have sexual contact with your partner when you have any symptom or outbreak of genital herpes.
- Use a condom made of latex or polyurethane whenever you have sexual contact.

Ask your healthcare provider for more information about safer sex practices.

#### **Who should not take valacyclovir hydrochloride tablets?**

**Do not take valacyclovir hydrochloride tablets** if you are allergic to any of its ingredients or to acyclovir. The active ingredient is valacyclovir. See the end of this leaflet for a complete list of ingredients in valacyclovir hydrochloride tablets.

#### **Before taking valacyclovir hydrochloride tablets, tell your healthcare provider:**

**About all your medical conditions**, including:

- **if you have had a bone marrow transplant or kidney transplant, or if you have advanced HIV disease or "AIDS"**. Patients with these conditions may have a higher chance for getting a blood disorder called thrombotic thrombocytopenic purpura/hemolytic uremic syndrome (TTP/HUS). TTP/HUS can result in death.
- **if you have kidney problems**. Patients with kidney problems may have a higher chance for getting side effects or more kidney problems with valacyclovir hydrochloride tablets. Your healthcare provider may give you a lower dose of valacyclovir hydrochloride tablets.
- **if you are 65 years of age or older**. Elderly patients have a higher chance of certain side effects. Also, elderly patients are more likely to have kidney problems. Your healthcare provider may give you a lower dose of valacyclovir hydrochloride tablets.
- **if you are pregnant or planning to become pregnant**. Talk with your healthcare provider about the risks and benefits of taking prescription drugs (including valacyclovir hydrochloride tablets) during pregnancy.
- **if you are breastfeeding**. Valacyclovir hydrochloride tablets may pass into your milk and it may harm your baby. Talk with your healthcare provider about the best way to feed your baby if you are taking valacyclovir hydrochloride tablets.
- **about all the medicines you take**, including prescription and non-prescription medicines, vitamins, and herbal supplements. Valacyclovir hydrochloride tablets may affect other medicines, and other medicines may affect valacyclovir hydrochloride tablets. It is a good idea to keep a complete list of all the medicines you take. Show this list to your healthcare provider and pharmacist any time you get a new medicine.

#### **How should I take valacyclovir hydrochloride tablets?**

Take valacyclovir hydrochloride tablets exactly as prescribed by your healthcare provider. Your dose of valacyclovir hydrochloride tablets and length of treatment will depend on the type of herpes infection that you have and any other medical problems that you have.

- Do not stop valacyclovir hydrochloride tablets or change your treatment without talking to your healthcare provider.
- Valacyclovir hydrochloride tablets can be taken with or without food.
- If you are taking valacyclovir hydrochloride tablets to treat cold sores, chickenpox, shingles, or genital herpes, you should start treatment as soon as possible after your symptoms start. Valacyclovir hydrochloride tablets may not help you if you start treatment too late.
- If you miss a dose of valacyclovir hydrochloride tablets, take it as soon as you remember and then take your next dose at its regular time. However, if it is almost time for your next dose, do not take the missed dose. Wait and take the next dose at the regular time.
- Do not take more than the prescribed number of valacyclovir hydrochloride tablets each day. Call your healthcare provider right away if you take too much valacyclovir hydrochloride tablets.

#### **What are the possible side effects of valacyclovir hydrochloride tablets?**

**Kidney failure and nervous system problems are not common, but can be serious in some patients taking valacyclovir hydrochloride tablets.** Nervous system problems include aggressive behavior, unsteady movement, shaky movements, confusion, speech problems, hallucinations (seeing or hearing things that are really not there), seizures, and coma. Kidney failure and nervous system problems have happened in patients who already have kidney disease and in elderly patients whose kidneys do not work well due to age. **Always tell your healthcare provider if you have kidney problems before taking valacyclovir hydrochloride tablets. Call your doctor right away if you get a nervous system problem while you are taking valacyclovir hydrochloride tablets.**

Common side effects of valacyclovir hydrochloride tablets in adults include headache, nausea, stomach pain, vomiting, and dizziness. Side effects in HIV-infected adults include headache, tiredness, and rash. These side effects usually are mild and do not cause patients to stop taking valacyclovir hydrochloride tablets.

Other less common side effects in adults include painful periods in women, joint pain, depression, low blood cell counts, and changes in tests that measure how well the liver and kidneys work.

The most common side effect seen in children <18 years of age was headache.

**Talk to your healthcare provider if you develop any side effects that concern you.**

These are not all the side effects of valacyclovir hydrochloride. For more information ask your healthcare provider or pharmacist.

**How should I store valacyclovir hydrochloride tablets?**

- Store at 20° to 25°C (68° to 77°F). [See USP controlled room temperature]
- Store valacyclovir hydrochloride suspension between 2° to 8°C (36° to 46°F) in a refrigerator. Discard after 28 days.
- Keep valacyclovir hydrochloride tablets in a tightly closed container.
- Do not keep medicine that is out of date or that you no longer need.
- Keep valacyclovir hydrochloride tablets and all medicines out of the reach of children.

**General information about valacyclovir hydrochloride tablets**

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use valacyclovir hydrochloride tablets for a condition for which it was not prescribed. Do not give valacyclovir hydrochloride tablets to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about valacyclovir hydrochloride tablets. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about valacyclovir hydrochloride tablets that is written for health professionals.

**What are the ingredients in valacyclovir hydrochloride tablets?**

**Active Ingredient:** valacyclovir hydrochloride

**Inactive Ingredients:** croscarmellose sodium, FD&C Blue No. 2 aluminium lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, povidone, talc and titanium dioxide.

**Manufactured by:**

Wockhardt Limited

Mumbai, India.

**Distributed by:**

Wockhardt USA LLC.

20 Waterview Blvd.

Parsippany, NJ 07054

USA.

Rev.080213

NDC 64679-152-01

\*Each film-coated tablet contains valacyclovir 500 mg as valacyclovir HCl.

**Valacyclovir HCl Tablets**

**500 mg\***

Rx only

30 Tablets

WOCKHARDT

Manufactured by: Wockhardt Limited, Mumbai, India.  
Distributed by: Wockhardt USA LLC., 20 Waterview Blvd, Parsippany, NJ 07054, USA.

MH/DRUGS/AD/068

LOT: 206426

EXP: NON-VARNISH AREA

Actual size: 20 mm x 7 mm

NDC 64679-153-04

\*Each film-coated tablet contains valacyclovir 1 gram as valacyclovir HCl.

**Valacyclovir HCl Tablets**

**1 gram\***

Rx only

500 Tablets

WOCKHARDT

Manufactured by: Wockhardt Limited, Mumbai, India.  
Distributed by: Wockhardt USA LLC., 20 Waterview Blvd, Parsippany, NJ 07054, USA.

MH/DRUGS/AD/068

LOT: 206424

EXP: NON-VARNISH AREA

Actual size: 67 mm x 13 mm

NDC 64679-153-06

\*Each film-coated tablet contains valacyclovir 1 gram as valacyclovir HCl.

**Valacyclovir HCl Tablets**

**1 gram\***

Rx only

250 Tablets

WOCKHARDT

Manufactured by: Wockhardt Limited, Mumbai, India.  
Distributed by: Wockhardt USA LLC., 20 Waterview Blvd, Parsippany, NJ 07054, USA.

MH/DRUGS/AD/068

LOT: 216866

EXP: NON-VARNISH AREA

Actual size: 11 mm x 55 mm

**VALACYCLOVIR HYDROCHLORIDE**

valacyclovir hydrochloride tablet, film coated

Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55648-152	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	VALACYCLOVIR HYDROCHLORIDE (UNII: G447S0T1VC) (ACYCLOVIR - UNII:X4HES1O11F)	VALACYCLOVIR	500 mg	
Inactive Ingredients				
	Ingredient Name		Strength	
	CELLULOSE, MICROCRYSTALLINE (UNII: OPIR32D61U)			
	CROSCARMELOSE SODIUM (UNII: M28OL1HH48)			
	FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			
	HYPROMELLOSES (UNII: 3NXW29V3WO)			
	MAGNESIUM STEARATE (UNII: 70097M6E00)			
	POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
	POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
	POVIDONE K90 (UNII: RDH86HV5Z)			
	TALC (UNII: 7SEV7J4R1U)			
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
Product Characteristics				
Color	BLUE (blue)	Score	no score	
Shape	CAPSULE (capsule-shaped)	Size	19mm	
Flavor		Imprint Code	W152	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55648-152-01	30 in 1 BOTTLE		
2	NDC:55648-152-03	90 in 1 BOTTLE		
3	NDC:55648-152-04	500 in 1 BOTTLE		
4	NDC:55648-152-05	10 in 1 CARTON		
4		10 in 1 BLISTER PACK		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA090216	05/24/2010		

VALACYCLOVIR HYDROCHLORIDE				
valacyclovir hydrochloride tablet, film coated				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55648-153	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	VALACYCLOVIR HYDROCHLORIDE (UNII: G447S0T1VC) (ACYCLOVIR - UNII:X4HES1O11F)	VALACYCLOVIR	1 g	
Inactive Ingredients				
	Ingredient Name		Strength	
	CELLULOSE, MICROCRYSTALLINE (UNII: OPIR32D61U)			
	CROSCARMELOSE SODIUM (UNII: M28OL1HH48)			
	FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			
	HYPROMELLOSES (UNII: 3NXW29V3WO)			
	MAGNESIUM STEARATE (UNII: 70097M6E00)			
	POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)			
	POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
	POVIDONE K90 (UNII: RDH86HV5Z)			
	TALC (UNII: 7SEV7J4R1U)			
	TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
Product Characteristics				
Color	BLUE (blue)	Score	2 pieces	
Shape	CAPSULE (capsule-shaped)	Size	23mm	
Flavor		Imprint Code	W153	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55648-153-02	30 in 1 BOTTLE		
2	NDC:55648-153-03	90 in 1 BOTTLE		
3	NDC:55648-153-04	500 in 1 BOTTLE		
4	NDC:55648-153-01	21 in 1 BOTTLE		
5	NDC:55648-153-06	250 in 1 BOTTLE		
6	NDC:55648-153-05	10 in 1 CARTON		
6		10 in 1 BLISTER PACK		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	

ANDA	ANDA090216	05/24/2010	
------	------------	------------	--

**Labeler** - Wockhardt Limited (650069115)

**Registrant** - Wockhardt Limited (650069115)

**Establishment**

Name	Address	ID/EI	Business Operations
Wockhardt Limited		676257570	ANALYSIS(55648-152, 55648-153) , MANUFACTURE(55648-152, 55648-153) , PACK(55648-152, 55648-153) , LABEL(55648-152, 55648-153)

Revised: 4/2013

Wockhardt Limited