

MAGNESIUM CITRATE SALINE LAXATIVE - LEMON FLAVOR- magnesium citrate solution

Universal Distribution Center LLC

Magnesium Citrate Saline Laxative Oral Solution - Lemon Flavor

Drug Facts

Active ingredient (in each fl oz)

Magnesium citrate 1.745 g

Purpose

Saline laxative

Uses:

- For relief of occasional constipation (irregularity).
- Generally produces bowel movement in ½ to 6 hours

Warnings:

Ask a doctor before use if you have

- Kidney disease • a magnesium restricted diet
- Abdominal pain, nausea, or vomiting
- Noticed a sudden change in bowel habits that persists over a period of 2 weeks
- Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

- Taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

- shake well before using • drink a full glass (8 ounces) of liquid with each dose • may be taken as a single daily dose or in divided doses
-

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

Other Information:

- **each fl oz contains:** magnesium 290 mg
- **each fl oz contains:** sodium 1 mg

store at temperatures between 46° and 86°F (8° and 30°C)

Inactive ingredients:

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Sugar Free

Alcohol content 0.025%
(From Flavoring)

MADE IN USA

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN, OR SEPARATED FROM NECKRING

Mfd for and Distributed By:
Universal Distribution Center LLC
330 Applegarth Road,
Monroe Township, NJ 08831
www.universaldc.com

Packaging

Drug Facts

store at temperatures between 46° and 86°F (8° and 30°C)

Directions:

■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours

children under 2 years of age ask a doctor

Other Information:

- each fl oz contains: magnesium 290 mg
- each fl oz contains: sodium 1 mg

Inactive ingredients: benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Mfd for and Distributed By:
Universal Distribution Center LLC
330 Applegarth Road,
Monroe Township, NJ 08831
www.universaldc.com



Magnesium Citrate

Saline Laxative Oral Solution

Lemon Flavor

Sugar Free

Alcohol content 0.025% (From Flavoring)



10 FL OZ (296mL)

Drug Facts (continued)

Active ingredient (in each fl oz) Purpose
Magnesium citrate 1.745 gSaline laxative

Uses: ■ For relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours

Warnings:

Ask a doctor before use if you have

- Kidney disease ■ a magnesium restricted diet
- Abdominal pain, nausea, or vomiting
- Noticed a sudden change in bowel habits that persists over a period of 2 weeks
- Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

- Taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN, OR SEPARATED FROM NECKRING

MAGNESIUM CITRATE SALINE LAXATIVE - LEMON FLAVOR

magnesium citrate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52000-449
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	0.059 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color		Score	
--------------	--	--------------	--

Shape		Size	
Flavor	LEMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52000-449-10	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/16/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	03/16/2026	

Labeler - Universal Distribution Center LLC (019180459)

Revised: 3/2026

Universal Distribution Center LLC