

**NOBLEAID PAIN RELIEVER- aspirin tablet, delayed release**  
**Noble Otc LLC**

-----  
**NobleAid ASPIRIN PAIN RELIEVER**

***Drug Facts***

***Active Ingredient (in each tablet)***

Aspirin 81 mg (NSAID)

***Purpose***

Pain Reliever

***Uses***

For the temporary relief of minor aches and pains

***WARNING***

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • Hives • Facial swelling • asthma (wheezing) • shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product •take more or for a longer time than directed.

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer.

**Ask a doctor before use if** •stomach bleeding •warning applies to you •you have a history of stomach problems such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease •you are taking a diuretic •you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for •gout •diabetes •arthritis

**Stop use and ask a doctor if** •an allergic reaction occurs. Seek medical help right away. •you experience any of the following signs of stomach bleeding: •feel faint •vomit blood •have bloody or black stools •have stomach pain that does not get better •pain gets worse or lasts more than 10 days •redness or swelling is present •fever lasts more

than 3 days •new symptoms occur ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

KEEP OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions:**

•Drink a full glass of water with each dose •adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours •children under 12 years: consult a doctor.

### **Other Information:**

*Store at room temperature*

### **Inactive ingredients:**

Microcrystalline Cellulose, Silicon dioxide, Talcum, Croscarmellose sodium, Stearic acid, Colloidal silicon dioxide, Titanium dioxide, Triethyl citrate, Methacrylic acid copolymer, D&C Yellow #10, FD&C Yellow #6

- **Safe pain relief**
- **Safety enteric-coated aspirin regimen**

### **ADULT LOW DOSE**

Compare to **Bayer®** Aspirin active ingredient

### **Questions or comments?**

Please call (347)378-8990 Monday Through Friday 9-5 or by Email [Info@nobleotc.com](mailto:Info@nobleotc.com)

This product is not manufactured or distributed by Bayer Healthcare, LLC owner of the Registered Trademark BAYER® Tablets

**TAMPER EVIDENT** Do not use if safety seal under cap is broken or missing.

Manufactured for

Noble OTC LLC

Perth Amboy, NJ 08861

[www.nobleotc.com](http://www.nobleotc.com)

### **Packaging**

NDC-81533-600-10

Compare to **BAYER®** Aspirin active ingredient

**Drug Facts**  
Active ingredient (in each tablet) Aspirin 81 mg (NSAID) Purpose Pain Reliever

**WARNING**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. If changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ Hives ■ Facial swelling ■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe somatic bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.  
**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer.  
**Ask a doctor before use if** ■ stomach bleeding ■ warning applies to you ■ you have a history of stomach problems such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma  
**Ask a doctor or pharmacist before use if** you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

**ASPIRIN**  
PAIN RELIEVER

✓ Safe pain relief  
✓ Safety enteric-coated aspirin regimen

**ADULT LOW DOSE**  
81 mg each tablet

**100 ENTERIC-COATED TABLETS**

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever lasts more than 3 days ■ new symptoms occur ringing in the ears or loss of hearing occurs  
**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.  
**Directions:** ■ Drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours ■ children under 12 years: consult a doctor.  
**Other Information:** Store at room temperature  
**Inactive ingredients:** Microcrystalline Cellulose, Silicon dioxide, Talcum, Croscarmellose sodium, Stearic acid, Colloidal silicon dioxide, Titanium dioxide, Triethyl citrate, Methacrylic acid copolymer, D&C Yellow #10, FD&C Yellow #6

**Questions or comments?**  
Please call (347) 378-8990 Monday Through Friday 9-5 or by Email info@nobleotc.com  
This product is not manufactured or distributed by Bayer Healthcare, LLC owner of the Registered Trademark BAYER® Tablets  
**TAMPER EVIDENT** Do not use if safety seal under cap is broken or missing.

Manufactured for Noble OTC LLC Perth Amboy, NJ 08861 www.nobleotc.com

1 95893 81505 0

NDC-81533-600-11

Compare to **BAYER®** Aspirin active ingredient

**Drug Facts**  
Active ingredient (in each tablet) Aspirin 81 mg (NSAID) Purpose Pain Reliever

**WARNING**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. If changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ Hives ■ Facial swelling ■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe somatic bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.  
**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer.  
**Ask a doctor before use if** ■ stomach bleeding ■ warning applies to you ■ you have a history of stomach problems such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma  
**Ask a doctor or pharmacist before use if** you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

**ASPIRIN**  
PAIN RELIEVER

✓ Safe pain relief  
✓ Safety enteric-coated aspirin regimen

**ADULT LOW DOSE**  
81 mg each tablet

**300 ENTERIC-COATED TABLETS**

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever lasts more than 3 days ■ new symptoms occur ringing in the ears or loss of hearing occurs  
**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.  
**Directions:** ■ Drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours ■ children under 12 years: consult a doctor.  
**Other Information:** Store at room temperature  
**Inactive ingredients:** Microcrystalline Cellulose, Silicon dioxide, Talcum, Croscarmellose sodium, Stearic acid, Colloidal silicon dioxide, Titanium dioxide, Triethyl citrate, Methacrylic acid copolymer, D&C Yellow #10, FD&C Yellow #6

**Questions or comments?**  
Please call (347) 378-8990 Monday Through Friday 9-5 or by Email info@nobleotc.com  
This product is not manufactured or distributed by Bayer Healthcare, LLC owner of the Registered Trademark BAYER® Tablets  
**TAMPER EVIDENT** Do not use if safety seal under cap is broken or missing.

Manufactured for Noble OTC LLC Perth Amboy, NJ 08861 www.nobleotc.com

1 95893 83010 4

NDC-81533-600-30

Compare to **BAYER®** Aspirin active ingredient

**Drug Facts**  
Active ingredient (in each tablet) Aspirin 81 mg (NSAID) Purpose Pain Reliever

**WARNING**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. If changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ Hives ■ Facial swelling ■ asthma (wheezing) ■ shock  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe somatic bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.  
**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer.  
**Ask a doctor before use if** ■ stomach bleeding ■ warning applies to you ■ you have a history of stomach problems such as heartburn you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma  
**Ask a doctor or pharmacist before use if** you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

**ASPIRIN**  
PAIN RELIEVER

✓ Safe pain relief  
✓ Safety enteric-coated aspirin regimen

**ADULT LOW DOSE**  
81 mg each tablet

**1000 ENTERIC-COATED TABLETS**

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ redness or swelling is present ■ fever lasts more than 3 days ■ new symptoms occur ringing in the ears or loss of hearing occurs  
**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **KEEP OUT OF REACH OF CHILDREN.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.  
**Directions:** ■ Drink a full glass of water with each dose ■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours ■ children under 12 years: consult a doctor.  
**Other Information:** Store at room temperature  
**Inactive ingredients:** Microcrystalline Cellulose, Silicon dioxide, Talcum, Croscarmellose sodium, Stearic acid, Colloidal silicon dioxide, Titanium dioxide, Triethyl citrate, Methacrylic acid copolymer, D&C Yellow #10, FD&C Yellow #6

**Questions or comments?**  
Please call (347) 378-8990 Monday Through Friday 9-5 or by Email info@nobleotc.com  
This product is not manufactured or distributed by Bayer Healthcare, LLC owner of the Registered Trademark BAYER® Tablets  
**TAMPER EVIDENT** Do not use if safety seal under cap is broken or missing.

Manufactured for Noble OTC LLC Perth Amboy, NJ 08861 www.nobleotc.com

1 95893 29018 5

## NOBLEAID PAIN RELIEVER

aspirin tablet, delayed release

### Product Information

#### Product Type

HUMAN OTC DRUG

#### Item Code (Source)

NDC:81533-600

#### Route of Administration

ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients	
Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics			
Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	l1
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81533-600-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2021	
2	NDC:81533-600-11	300 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2021	
3	NDC:81533-600-30	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/15/2021	

**Labeler** - Noble Otc LLC (041836435)