

**ICON 10000 X NPE FREE- iodine liquid
IBA Inc.**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ICON 10000 X NPE Free

ACTIVE INGREDIENT

Iodine Complex.....20.00%

Provide a full 1% available Iodine when used at the recommended dilution ratio)

Inert Ingredients

Duel emollient system 40.00%

(provides a total of 10% skin conditioners at the recommended dilution ratio)

Other Inerts..... 40.00%

(Includes Buffering Agents, Stabilizing Agents and Surfactants)

Total.....100.00%

When mixed according to label directions, this product provides highly effective levels of free Iodine for fast kill of mastitis causing organisms.

INDICATIONS AND USAGE

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment
4. Wash Teat Dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. REMOVE VISIBLE SOILS FROM THE TEATS.
2. OBSERVE FOREMILK BY STRIPPING TWO OR THREE STREAMS OF MILK INTO A STRIP CUP.
3. DIP OR SPRAY THE COW'S TEATS FULL LENGTH WITH ICON 1000 X NPE Free TEAT DIP.
- 4 WAIT APPROXIMATELY 30 SECONDS.
5. REMOVE ALL TEAT DIP BY THOROUGHLY DRYING THE TEATS AND TEAT ORIFICES WITH A CLEAN, SINGLE SERVICE PAPER TOWEL(S).
6. ATTACH MILKING UNIT

B. POST-MILKING (POST DIPPING)

Immediately after milking dip all teats full length with ICON 10000 X NPE Free Teat Dip
Ensure good coverage of all sides of the teats and teat orifices.

DILUTE 1 PART ICON 10000 X TEAT DIP PER 3 PARTS POTABLE, TEPID WATER

CAUTION: KEEP OUT OF REACH OF CHILDREN.

AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

WARNING

ICON 10000 X NPE Free IODINE TEAT DIP CONCENTRATE MUST BE DILUTED BEFORE USE.

DO NOT MIX WITH ANYTHING BUT POTABLE WATER.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

**ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES
CALL 1-800-424-9300**

Icon 10000 X NPE Free

NDC 29983-1067-9



A HIGHLY EFFECTIVE SANITIZING TEAT DIP

AN AID IN THE PREVENTION OF NEW INTRAMAMMARY INFECTIONS



IODINE TEAT DIP CONCENTRATE

Especially Formulated For Dairies Where **NPE FREE** Teat Dips Are Preferred Or Required

ACTIVE INGREDIENT:
Iodine Complex..... 20.00%
(Provides A Full 1% Available Iodine When Used At The Recommended Dilution Ratio)
INERT INGREDIENTS:
Dual Emollient System..... 40.00%
(Provides A Total Of 10% Skin Conditioners At The Recommended Dilution Ratio)
Other Inerts..... 40.00%
(Includes Buffering Agents, Stabilizing Agents and Surfactants)
TOTAL..... 100.00%

When mixed according to label directions, this product provides highly effective levels of free iodine for fast kill of mastitis causing organisms.

GENERAL RECOMMENDATIONS

- Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
- When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
- Do not use this product for cleaning or sanitizing equipment.
- Wash teat dip cups after each milking.

FIRST AID PROCEDURES

EYE CONTACT: IMMEDIATELY FLUSH EYE WITH LARGE AMOUNTS OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUES. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE.

SKIN CONTACT: FOR CONTACT WITH CONCENTRATED PRODUCTS, FLUSH THOROUGHLY WITH FRESH, TEPID WATER. DISCARD CONTAMINATED CLOTHING AND FOOTWEAR OR WASH BEFORE REUSE. FOR READY-TO-USE PRODUCTS, WASH ANY SENSITIVE AREAS.

INGESTION: DO NOT INDUCE VOMITING. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. DRINK LARGE AMOUNTS OF WATER. IF VOMITING OCCURS, KEEP AIRWAYS OPEN. KEEP HEAD LOWER THAN THE HIPS TO PREVENT ASPIRATION INTO THE LUNGS.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION FOR MEDICAL EMERGENCIES CALL 1-800-424-9300



Dilute 1 Part
ICON 10000X NPE FREE
TEAT DIP CONCENTRATE
Per 3 Parts Potable,
Tepid Water.

PATENT PENDING

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

- Remove visible soils from the teats.
- Observe foremilk by stripping two or three streams of milk into a strip cup.
- Dip, foam or spray the cow's teats full length with **DILUTED ICON 10000X NPE FREE TEAT DIP**.
- Wait approximately 30 seconds.
- Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if foaming or spraying) with a clean, single service towel(s) to avoid contamination of milk.
- Attach milking unit.

B. POST-MILKING (POST-DIPPING)

- Immediately after milking, dip or spray all teats full length with **DILUTED ICON 10000X NPE FREE TEAT DIP**. Ensure good coverage of all sides of the teats and teat orifices.

CAUTION: KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

WARNING

ICON 10000X NPE FREE TEAT DIP CONCENTRATE MUST BE DILUTED BEFORE USE. DO NOT MIX WITH ANYTHING BUT POTABLE WATER.

PROTECT THIS PRODUCT FROM FREEZING.

DO NOT RETURN DILUTED TEAT DIP TO STORAGE CONTAINER.

ALWAYS ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE: LOT NO.: NET CONTENTS: 275 GALLONS (1,040.875 LITERS) IBA STOCK # 261295 321W

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 103 GILMORE DRIVE, SUTTON, MA 01590

ICON 10000 X NPE FREE

iodine liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-1067
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	1.00 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-1067-8	238.24 kg in 1 DRUM		
2	NDC:29983-1067-9	1191.24 kg in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/06/2017	

Labeler - IBA Inc. (019494160)

Registrant - IBA (019494160)

Establishment

Name	Address	ID/FEI	Business Operations
Morgan Gallacher Inc. DBA Custom Chemical Formulators		028311595	manufacture, api manufacture, pack

Revised: 12/2023

IBA Inc.