

**PREDNISOLONE- prednisolone oral solution**  
**Chartwell RX, LLC**


-----  
**PREDNISOLONE ORAL SOLUTION, USP 15 mg per 5 mL**

**Rx only**

**DESCRIPTION**

Prednisolone oral solution contains prednisolone which is a glucocorticoid. Glucocorticoids are adrenocortical steroids, both naturally occurring and synthetic, which are readily absorbed from the gastrointestinal tract. Prednisolone is a white to practically white, odorless, crystalline powder. It is very slightly soluble in water, soluble in methanol and in dioxane; sparingly soluble in acetone and in alcohol, slightly soluble in chloroform.

The chemical name for Prednisolone is Pregna-1,4 -diene -3, 20 - dione, 11, 17, 21-trihydroxy-, (11 $\beta$ ). Its molecular weight is 360.45. The molecular formula is C<sub>21</sub>H<sub>28</sub>O<sub>5</sub> and the structural formula is:



Prednisolone oral solution contains 15 mg of prednisolone in each 5 mL. Benzoic acid, 0.1% is added as a preservative. It also contains ethyl alcohol 5% (v/v), citric acid, edetate disodium, glycerin, propylene glycol, purified water, saccharin sodium, sucrose, natural and artificial wild cherry flavor, FD&C red #40.

**CLINICAL PHARMACOLOGY**

Naturally occurring glucocorticoids (hydrocortisone and cortisone), which also have salt-retaining properties, are used as replacement therapy in adrenocortical deficiency states. Their synthetic analogs such as prednisolone are primarily used for their potent anti-inflammatory effects in disorders of many organ systems.

Glucocorticoids such as prednisolone cause profound and varied metabolic effects. In addition, they modify the body's immune responses to diverse stimuli.

## **INDICATIONS AND USAGE**

Prednisolone Oral Solution, USP is indicated in the following conditions:

### **1. Endocrine Disorders**

Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is the first choice; synthetic analogs may be used in conjunction with mineralocorticoids where applicable; in infancy mineralocorticoid supplementation is of particular importance).

1. Congenital adrenal hyperplasia
2. Nonsuppurative thyroiditis
3. Hypercalcemia associated with cancer

### **2. Rheumatic Disorders**

As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in:

1. Psoriatic arthritis
2. Rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy)
3. Ankylosing spondylitis
4. Acute and subacute bursitis
5. Acute nonspecific tenosynovitis
6. Acute gouty arthritis
7. Post-traumatic osteoarthritis
8. Synovitis of osteoarthritis
9. Epicondylitis

### **3. Collagen Diseases**

During an exacerbation or as maintenance therapy in selected cases of:

1. Systemic lupus erythematosus
2. Acute rheumatic carditis

### **4. Dermatologic Diseases**

1. Pemphigus
2. Bullous dermatitis herpetiformis
3. Severe erythema multiforme (Stevens-Johnson syndrome)
4. Exfoliative dermatitis
5. Mycosis fungoides
6. Severe psoriasis
7. Severe seborrheic dermatitis

### **5. Allergic States**

Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment:

1. Seasonal or perennial allergic rhinitis

2. Bronchial asthma
3. Contact dermatitis
4. Atopic dermatitis
5. Serum sickness
6. Drug hypersensitivity reactions

## **6. Ophthalmic Diseases**

Severe acute and chronic allergic and inflammatory processes involving the eye and its adnexa such as:

1. Allergic corneal marginal ulcers
2. Herpes zoster ophthalmicus
3. Anterior segment inflammation
4. Diffuse posterior uveitis and choroiditis
5. Sympathetic ophthalmia
6. Allergic conjunctivitis
7. Keratitis
8. Chorioretinitis
9. Optic neuritis
10. Iritis and iridocyclitis

## **7. Respiratory Diseases**

Symptomatic sarcoidosis Loeffler's syndrome not manageable by other means

1. Berylliosis
2. Fulminating or disseminated pulmonary tuberculosis when used concurrently with appropriate chemotherapy
3. Aspiration pneumonitis

## **8. Hematologic Disorders**

Idiopathic thrombocytopenic purpura in adults

Secondary thrombocytopenia in adults

Acquired (autoimmune) hemolytic anemia

Erythroblastopenia (RBC anemia)

Congenital (erythroid) hypoplastic anemia

## **9. Neoplastic Diseases**

For palliative management of:

1. Acute leukemia of childhood
2. Leukemias and lymphomas in adults

## **10. Edematous States**

To induce a diuresis or remission of proteinuria in the nephrotic syndrome, without uremia, of the idiopathic type or that due to lupus erythematosus.

## **11. Gastrointestinal Diseases**

To tide the patient over a critical period of the disease in:

1. Ulcerative colitis
2. Regional enteritis

## **12. Miscellaneous**

Tuberculous meningitis with subarachnoid block or impending block used concurrently with appropriate antituberculous chemotherapy. Trichinosis with neurologic or myocardial involvement.

In addition to the above indications prednisolone oral solution is indicated for systemic dermatomyositis (polymyositis).

## **CONTRAINDICATIONS**

Systemic fungal infections.

## **WARNINGS**

In patients on corticosteroid therapy subjected to unusual stress, increased dosage of rapidly acting corticosteroids before, during, and after the stressful situation is indicated.

Corticosteroids may mask some signs of infection, and new infections may appear during their use. There may be decreased resistance and inability to localize infection when corticosteroids are used.

Prolonged use of corticosteroids may produce posterior subcapsular cataracts, glaucoma with possible damage to the optic nerves, and may enhance the establishment of secondary ocular infections due to fungi or viruses.

Average and large doses of hydrocortisone or cortisone can cause elevation of blood pressure, salt and water retention, and increased excretion of potassium. These effects are less likely to occur with the synthetic derivatives except when used in large doses. Dietary salt restriction and potassium supplementation may be necessary. All corticosteroids increase calcium excretion.

While on corticosteroid therapy, patients should not be vaccinated against smallpox. Other immunization procedures should not be undertaken in patients who are on corticosteroids, especially on high dose, because of possible hazards of neurological complications and a lack of antibody response.

Persons who are on drugs which suppress the immune system are more susceptible to infections than healthy individuals. Chickenpox and measles, for example, can have a more serious or even fatal course in non-immune children or adults on corticosteroids. In such children or adults who have not had these diseases, particular care should be taken to avoid exposure. How the dose, route and duration of corticosteroid administration affects the risk of developing a disseminated infection is not known. The contribution of the underlying disease and/or prior corticosteroid treatment to the risk is also not known. If exposed to chickenpox, prophylaxis with varicella zoster immune globulin (VZIG) may be indicated. If exposed to measles, prophylaxis with pooled intramuscular immunoglobulin (IG) may be indicated. (See the respective package inserts for complete VZIG and IG prescribing information.) If chickenpox develops, treatment with antiviral agents may be considered.

The use of prednisolone oral solution in active tuberculosis should be restricted to those cases of fulminating or disseminated tuberculosis in which the corticosteroid is used for the management of the disease in conjunction with an appropriate antituberculous regimen.

**If corticosteroids are indicated in patients with latent tuberculosis or tuberculin reactivity, close observation is necessary as reactivation of the disease may occur. During prolonged corticosteroid therapy, these patients should receive chemoprophylaxis.**

**Use in pregnancy:** Since adequate human reproduction studies have not been done with corticosteroids, the use of these drugs in pregnancy, nursing mothers or women of childbearing potential requires that the possible benefits of the drug be weighed against the potential hazards to the mother and embryo or fetus. Infants born of mothers who have received substantial doses of corticosteroids during pregnancy should be carefully observed for signs of hypoadrenalism.

## **PRECAUTIONS**

### **General**

Drug-induced secondary adrenocortical insufficiency may be minimized by gradual reduction of dosage. This type of relative insufficiency may persist for months after discontinuation of therapy; therefore, in any situation of stress occurring during that period, hormone therapy should be reinstated. Since mineralocorticoid secretion may be impaired, salt and/or a mineralocorticoid should be administered concurrently.

There is an enhanced effect of corticosteroids on patients with hypothyroidism and in those with cirrhosis.

Corticosteroids should be used cautiously in patients with ocular herpes simplex because of possible corneal perforation.

The lowest possible dose of corticosteroid should be used to control the condition under treatment, and when reduction in dosage is possible, the reduction should be gradual.

Psychic derangements may appear when corticosteroids are used, ranging from euphoria, insomnia, mood swings, personality changes, and severe depression, to frank psychotic manifestations. Also, existing emotional instability or psychotic tendencies may be aggravated by corticosteroids.

Aspirin should be used cautiously in conjunction with corticosteroids in hypoprothrombinemia.

Steroids should be used with caution in nonspecific ulcerative colitis, if there is a probability of impending perforation, abscess or other pyogenic infections; diverticulitis; fresh intestinal anastomoses; active or latent peptic ulcer; renal insufficiency; hypertension; osteoporosis; and myasthenia gravis.

Growth and development of infants and children on prolonged corticosteroid therapy should be carefully observed.

### **Information for Patients**

Patients who are on immunosuppressant doses of corticosteroids should be warned to

avoid exposure to chickenpox or measles. Patients should also be advised that if they are exposed, medical advice should be sought without delay.

## **ADVERSE REACTIONS**

### **Fluid and Electrolyte Disturbances**

1. Sodium retention
2. Fluid retention
3. Congestive heart failure in susceptible patients
4. Potassium loss
5. Hypokalemic alkalosis
6. Hypertension

### **Musculoskeletal**

1. Muscle weakness
2. Steroid myopathy
3. Loss of muscle mass
4. Osteoporosis
5. Vertebral compression fractures
6. Aseptic necrosis of femoral and humeral heads
7. Pathologic fracture of long bones

### **Gastrointestinal**

1. Peptic ulcer with possible perforation and hemorrhage
2. Pancreatitis
3. Abdominal distention
4. Ulcerative esophagitis

### **Dermatologic**

1. Impaired wound healing
2. Thin fragile skin
3. Petechiae and ecchymoses
4. Facial erythema
5. Increased sweating
6. May suppress reactions to skin tests

### **Neurological**

1. Convulsions
2. Increased intracranial pressure with papilledema (pseudo-tumor cerebri) usually after treatment
3. Vertigo
4. Headache

### **Endocrine**

1. Menstrual irregularities
2. Development of Cushingoid state
3. Suppression of growth in children
4. Secondary adrenocortical and pituitary unresponsiveness, particularly in times of stress, as in trauma, surgery or illness

5. Decreased carbohydrate tolerance
6. Manifestations of latent diabetes mellitus
7. Increased requirements for insulin or oral hypoglycemic agents in diabetics

### **Ophthalmic**

1. Posterior subcapsular cataracts
2. Increased intraocular pressure
3. Glaucoma
4. Exophthalmos

### **Metabolic**

1. Negative nitrogen balance due to protein catabolism

To report SUSPECTED ADVERSE REACTIONS, contact Chartwell RX, LLC. at 1-845-232-1683 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

## **DOSAGE AND ADMINISTRATION**

Dosage of Prednisolone Oral Solution, USP should be individualized according to the severity of the disease and the response of the patient. For infants and children, the recommended dosage should be governed by the same considerations rather than strict adherence to the ratio indicated by age or body weight.

Hormone therapy is an adjunct to and not a replacement for conventional therapy.

Dosage should be decreased or discontinued gradually when the drug has been administered for more than a few days.

The severity, prognosis, expected duration of the disease, and the reaction of the patient to medication are primary factors in determining dosage.

If a period of spontaneous remission occurs in a chronic condition, treatment should be discontinued.

Blood pressure, body weight, routine laboratory studies, including two-hour postprandial blood glucose and serum potassium, and a chest X-ray should be obtained at regular intervals during prolonged therapy. Upper GI X-rays are desirable in patients with known or suspected peptic ulcer disease.

The initial dosage of Prednisolone Oral Solution, USP may vary from 5 mg to 60 mg per day depending on the specific disease entity being treated. In situations of less severity lower doses will generally suffice while in selected patients higher initial doses may be required. The initial dosage should be maintained or adjusted until a satisfactory response is noted. If after a reasonable period of time there is a lack of satisfactory clinical response, Prednisolone Oral Solution, USP should be discontinued and the patient transferred to other appropriate therapy. **IT SHOULD BE EMPHASIZED THAT DOSAGE REQUIREMENTS ARE VARIABLE AND MUST BE INDIVIDUALIZED ON THE BASIS OF THE DISEASE UNDER TREATMENT AND THE RESPONSE OF THE PATIENT.**

After a favorable response is noted, the proper maintenance dosage should be determined by decreasing the initial drug dosage in small decrements at appropriate time intervals until the lowest dosage which will maintain an adequate clinical response is reached. It should be kept in mind that constant monitoring is needed in regard to drug

dosage. Included in the situations which may make dosage adjustments necessary are changes in clinical status secondary to remissions or exacerbations in the disease process, the patient's individual drug responsiveness, and the effect of patient exposure to stressful situations not directly related to the disease entity under treatment. In this latter situation it may be necessary to increase the dosage of Prednisolone Oral Solution, USP for a period of time consistent with the patient's condition. If after long-term therapy the drug is to be stopped, it is recommended that it be withdrawn gradually rather than abruptly.

## **HOW SUPPLIED**

Prednisolone Oral Solution, USP containing 15 mg of Prednisolone in each 5 mL (teaspoonful) is a clear pinkish-red wild cherry flavored liquid and is supplied in 240 mL bottles (NDC 62135-250-37) and 480 mL (NDC 62135-250-47) bottles.

Pharmacist: Dispense 15 mg/5 mL Prednisolone Oral Solution, USP with suitable calibrated measuring device to assure proper measuring of dose.

## **DOSE/VOLUME CHART**

15 mg prednisolone	=	1 teaspoon
10 mg prednisolone	=	2/3 teaspoon
7.5 mg prednisolone	=	1/2 teaspoon
5 mg prednisolone	=	1/3 teaspoon

Dispense in tight, light-resistant and child-resistant container as defined in USP/NF.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. DO NOT REFRIGERATE.

Manufactured for:

Chartwell RX, LLC  
Congers, NY 10920

L71082

Rev. 09/2022

## **PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**PREDNISOLONE ORAL SOLUTION, USP 15mg per 5mL - NDC- 62135-250-37 -  
240 mL Bottle Label**



NDC 62135-250-37

# PrednisoLONE

Oral Solution, USP

**15 mg per 5 mL**  
Ethyl Alcohol 5% (v/v)

Rx Only  
240 mL

Chartwell Rx

**USUAL DOSAGE:** See accompanying prescribing information. Dispense in tight, light-resistant and child-resistant containers as defined in USP.

**Store at 20° to 25° C (68° to 77° F) [See USP Controlled Room Temperature].**

**DO NOT REFRIGERATE.**

**PHARMACIST:** Dispense with a suitable Calibrated Measuring Device.

**DESCRIPTION:** Prednisolone Oral Solution, USP contains 15 mg of prednisolone in each 5 mL (teaspoonful) and ethyl alcohol 5% (v/v). Benzoic acid 0.1% added as a preservative.

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

Manufactured By: Chartwell Pharmaceuticals  
Carmel, NY

Manufactured For: Chartwell RX, LLC  
Congers, NY 10920

GTIN 00362135250379 L71080 REV. 01 09/22



No Varnish

## PREDNISOLONE ORAL SOLUTION, USP 15mg per 5mL - NDC- 62135-250-47 - 480 mL Bottle Label

NDC 62135-250-47

# PrednisoLONE

Oral Solution, USP

**15 mg per 5 mL**  
Ethyl Alcohol 5% (v/v)

Rx Only  
480 mL

Chartwell Rx

**USUAL DOSAGE:** See accompanying prescribing information. Dispense in tight, light-resistant and child-resistant containers as defined in USP.

**Store at 20° to 25° C (68° to 77° F) [See USP Controlled Room Temperature].**

**DO NOT REFRIGERATE.**

**PHARMACIST:** Dispense with a suitable Calibrated Measuring Device.

**DESCRIPTION:** Prednisolone Oral Solution, USP contains 15 mg of prednisolone in each 5 mL (teaspoonful) and ethyl alcohol 5% (v/v). Benzoic acid 0.1% added as a preservative.

**KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.**

Manufactured by: Chartwell Pharmaceuticals Carmel, LLC.  
Carmel, NY 10512

Manufactured for: Chartwell RX, LLC.  
Congers, NY 10920

GTIN 00362135250478 L71081 REV. 01 09/22



No Varnish

### PREDNISOLONE

prednisolone oral solution

#### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62135-250
Route of Administration	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
PREDNISOLONE (UNII: 9PHQ9Y1OLM) (PREDNISOLONE - UNII:9PHQ9Y1OLM)		PREDNISOLONE	15 mg in 5 mL	
Inactive Ingredients				
Ingredient Name		Strength		
CHERRY (UNII: BUC5I9595W)				
BENZOIC ACID (UNII: 8SKN0B0MIM)				
ALCOHOL (UNII: 3K9958V90M)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
GLYCERIN (UNII: PDC6A3C0OX)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0K00R)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SUCROSE (UNII: C151H8M554)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
Product Characteristics				
Color	pink (clear pinkish-red)	Score		
Shape		Size		
Flavor	CHERRY (wild cherry)	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62135-250-37	240 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/20/2022	
2	NDC:62135-250-47	480 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/20/2022	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA040323	11/20/2022		

**Labeler** - Chartwell RX, LLC (079394054)