

GUAIFENESIN- guaifenesin tablet
Major Pharmaceuticals

Major 44-588

Active ingredient (in each immediate-release tablet)

Guaifenesin 200 mg

Purpose

Expectorant

Uses

helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Ask a doctor before use if you have

- cough accompanied by too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not exceed 6 doses in 24 hours**
- take with a full glass of water

adults and children 12 years and over	1 to 2 tablets every 4 hours
---	---------------------------------

children 6 to under 12 years	½ to 1 tablet every 4 hours
children under 6 years	ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

FD&C red #40 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid

Questions or comments?

(800) 616-2471

Principal Display Panel

MAJOR®

NDC 0904-5154-60

Immediate Release

Guaifenesin

200 mg

Expectorant

Relieves Chest Congestion

Thins and Loosens Mucus

Actual Size

100 Tablets

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Distributed by **MAJOR® PHARMACEUTICALS**

17177 N Laurel Park Drive, Suite 233, Livonia, MI 48152

Rev. 11/19 M-17 Re-Order No. 238163

50844 REV0819C58812

Drug Facts (continued)	
Directions	
<ul style="list-style-type: none"> do not exceed 6 doses in 24 hours take with a full glass of water 	<ul style="list-style-type: none"> adults and children 12 years and over: 1 to 2 tablets every 4 hours children 6 to under 12 years: 1/2 to 1 tablet every 4 hours children under 6 years: ask a doctor
Other information	
<ul style="list-style-type: none"> store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) use by expiration date on package 	
Inactive ingredients FD&C red #40 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, povidone, silicon dioxide, sodium starch glycolate, stearic acid	
Questions or comments? (800) 616-2471	
STOP PEELING	

MAJOR® NDC 0904-5154-60

Immediate Release
Guaifenesin

200 mg

Expectorant

Relieves Chest Congestion
Thins and Loosens Mucus

100 Tablets

Actual Size 

Distributed by: MAJOR® PHARMACEUTICALS
17777 N Laurel Park Drive, Suite 233, Livonia, MI 48152
Rev. 11/19 M-17 Re-order No. 238163 50844 REV0819C58812



3 090451154601 1

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient <i>(in each immediate-release tablet)</i> Guaifenesin 200 mg Expectorant
Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
Warnings Ask a doctor before use if you have <ul style="list-style-type: none"> cough accompanied by too much phlegm (mucus) persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema Stop use and ask a doctor if cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash or persistent headache. These could be signs of a serious condition. If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

PEEL HERE FOR MORE DRUG FACTS

Major 44-588

GUAIFENESIN			
guaifenesin tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-5154
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
Inactive Ingredients			
	Ingredient Name		Strength

FD&C RED NO. 40 (UNII: WZB9127XOA)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	pink (dark)	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;588
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-5154-60	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/05/2009	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/05/2009	

Labeler - Major Pharmaceuticals (191427277)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0904-5154)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0904-5154)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	pack(0904-5154)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0904-5154)

